



# Extraction rate of Valuable Compounds from Peanut Skin Waste by Ethanol-Assisted Supercritical Carbon Dioxide: Modelling and Optimization

Nicky Rahmana Putra<sup>a</sup>, Dwila Nur Rizkiyah<sup>a</sup>, Ahmad Hazim Abdul Aziz<sup>b</sup>, Zuhaili Idham<sup>a</sup>, Lailatul Qomariyah<sup>c</sup>, Mohd Azizi Che Yunus<sup>a,\*</sup>

<sup>a</sup> Centre of Lipid Engineering and Advanced Research (CLEAR), Ibnu Sina Institute for Scientific and Industrial Research, Universiti Teknologi Malaysia, 81310, UTM Johor Bahru, Malaysia; <sup>b</sup> Faculty of Food Science and Nutrition, Universiti Malaysia Sabah, 88400 Kota Kinabalu, Sabah, Malaysia; <sup>c</sup> Department of Industrial Chemical Engineering, Institut Teknologi Sepuluh Nopember, 60111 Surabaya, Surabaya, Indonesia

**Abstract** Response Surface Methodology (RSM) was employed to optimize the extraction rate of phenolic and flavonoid contents from peanut skin by supercritical carbon dioxide (ScCO<sub>2</sub>) assisted by ethanol as entrainer. The studied extraction parameters were pressure (10 to 30 MPa), temperature (40 to 70 °C), and the ratio of ethanol (2.5 to 7.5%). Brunner's and Esquivel's models were applied to evaluate the extraction rate. The best-operating conditions, in the tested range, were 30 MPa, 40 °C, and 4.64% of ethanol ratio, with a maximum extraction rate of 0.22 mg/g.sec and 0.19 mg/g.sec of the phenolic and flavonoid content, respectively. The findings concluded that higher-pressure condition has a significant impact on the extraction rate of phenolic and flavonoid.

**Keywords:** Peanut skin; Phenolic; Flavonoid; Supercritical Carbon Dioxide/Ethanol; Extraction

## Introduction

Peanut skin is a by-product of peanut-based industry and peeled out due to its astringent taste, as well as might reduce the quality of the final products [1]. Peanut (*Arachis Hypohea*) skin also comprises high-flavonoid and phenolic compounds, which could inhibit the oxidative degradation of lipids while improving the nutritional value [2]. Peanut skin extract also could act as anti-inflammatory, anti-viral, and anti-carcinogenic; which could prevent cancers [3].

Supercritical carbon dioxide (ScCO<sub>2</sub>) extraction as green technology is preferable to extract peanut skin due to the application of high-purity and non-organic solvent. ScCO<sub>2</sub> offers a higher quantity of bioactive compounds extraction and high solubility of interest compounds (catechin, phenolic and flavonoid) compared to the conventional method (soxhlet extraction) [2, 4, 5]. Nevertheless, one of the drawbacks of ScCO<sub>2</sub> is the limited interest compounds, as it only applies to the non-polar. The addition of ethanol into ScCO<sub>2</sub> is one of the novel approaches to extract polar compounds, especially samples that contain flavonoid contents [6]. The advantage of using SC-CO<sub>2</sub> is it provides high amounts of bioactive compounds at a low critical temperature, which is cost-effective and suitable for thermo-sensitive compounds.

**\*For correspondence:**

r-azizi@utm.my

**Received:** 4 May 2021

**Accepted:** 20 April 2022

© Copyright Putra *et al.*

This article is distributed under the terms of the

[Creative Commons](#)

[Attribution License](#), which

permits unrestricted use

and redistribution provided

that the original author and

source are credited.

The application and optimization of ScCO<sub>2</sub> on phenolic and flavonoid extraction is widely studied. However, there are limited studies to determine the extraction rate of phenolic and flavonoid content in the ScCO<sub>2</sub> [7-10]. There are several ways to analyze the extraction rate data experimentally, Brunner's and Esquivel's models as empirical models are commonly used to correlate the experimental data to obtain the extraction rate [11, 12]. Response Surface methodology has been used to determine the parameter effects to enhance the extraction rate of phenolic and flavonoid. As a result, the optimum parameter could be obtained while predicting maximum responses [12].

Therefore, this study aimed to optimize the extraction parameters for high extraction rate of phenolic and flavonoid content from peanut skin by ethanol-assisted ScCO<sub>2</sub>.

## Materials and methods

### *Sample preparations*

Peanut skin as a waste of peanut butter industry (G-Tech, Johor Bahru) was dried for 4 hours and sieved to 355-425  $\mu\text{m}$  at 60 °C before kept in a refrigerator at -10 °C. The moisture content of the sample was reduced to less than 10% (Libherr EFL 3505, USA).

### *Chemical*

CO<sub>2</sub> (99.99% purity) was purchased from Kras Instrument, Malaysia. Gallic acid and quercetin were bought from and Sigma-Aldrich (St. Louis, USA). Ethanol (99.86% purity), methanol (99.86% purity) HPLC grade, Folin-Ciocalteu, n-hexane (99.86% purity), Na<sub>2</sub>CO<sub>3</sub>, Al<sub>2</sub>NO<sub>3</sub> and CH<sub>3</sub>COOK were obtained from Fisher Scientific (Atlanta, USA).

### *ScCO<sub>2</sub> Extraction assisted by ethanol*

The system of ScCO<sub>2</sub> consists of: 50 mL extraction vessel (dimensions: 1.4 cm internal diameter, 33 cm long), CO<sub>2</sub> pump (Supercritical 24, Japan), back pressure regulator (Jasco BP 2080, Japan), and oven (Mettler, Germany). The schematic diagram was shown in previous studies in Chiller's temperature was set to 6 °C. Next, about 5  $\pm$  0.005 g of peanut skin was measured and put into an extraction vessel before kept in the chiller. Then, the heater of the back-pressure regulator (Jasco BP 2080, Japan) was set at 50 °C. Liquid CO<sub>2</sub> and ethanol (99.86% purity) were later controlled at a constant flow rate of 3 mL/min. The extraction time was 180 minutes and the extracted yield was recorded and collected every 30 minutes before stored in a refrigerator (Libherr EFL 3505, USA) at -10°C for further analysis.

### *Total Phenolic Content (TPC) Analysis*

Analysis of total phenolic content was performed according by Rosli, Roslan [13]. 5 mL of Folin-Ciocalteu reagent was mixed with 50 mL of H<sub>2</sub>O while 3 g of Na<sub>2</sub>CO<sub>3</sub> was mixed with 40 mL of distilled water before sonicated for five minutes. Next, 1 mg of extract was diluted to 1 mL of ethanol and mixed by 5 mL of Folin-Ciocalteu mixture. The solution was then let to suspend by resting for ten minutes in room temperature. Later, 4 mL of Na<sub>2</sub>CO<sub>3</sub> mixture was added into the solution and placed in room temperature for 30 minutes. A Spectrophotometer UV-Vis (Jasco, Japan) was used to measure the absorbance with a wavelength of 760 nm. Meanwhile, for a standard curve, gallic acid was prepared according to a method by Rosli, Roslan [13], and results were expressed as gallic acid equivalent (mg/g sample)

### *Total Flavonoid Content (TFC) Analysis*

Analysis of total flavonoid content was performed according to Silva, Favaro-Trindade [14]. 1 mg of extract was diluted by 1 mL of ethanol. Next, in a different test tube, 1 mL ethanoic extract was mixed with 0.2 mL of Al<sub>2</sub>NO<sub>3</sub> 10%. Later, 0.2 mL of 1.0 M CH<sub>3</sub>COOK was added to the solution. The solution was rested for 40 min in room temperature. The wavelength of a spectrophotometer UV-Vis (Jasco, Japan) was set at 415 nm. The results were expressed as quercetin equivalent (mg/g sample)

### *Empirical modeling*

Brunner's and Esquivel's model were used to fitting the extraction rate data of phenolic and flavonoid

content. Esquivel model has two adjustable parameters ( $Y_2$  and  $k_2$ ) as shown in Eq. 1 [15]:

$$P_t = Y_2 \left( \frac{t}{k_2 + t} \right) \tag{1}$$

where  $P_t$  is the total phenolic or flavonoid content (mg/g),  $Y_2$  is the predicted total phenolic or flavonoid content (mg/g),  $k_2$  are the adjustable parameters (sec), and  $t$  is extraction time (sec). Therefore, the extraction rate can be obtained by  $Y_2/k_2$  (mg/g.sec).

Brunner's model also has two adjustable parameters ( $Y_2$  and  $k_2$ ) that represents a specific solution of Fick's law, as shown in Eq.2 [16]:

$$P_t = Y_2(1 - e^{-k_2t}) \tag{2}$$

where  $P_t$  is the total phenolic or flavonoid content (mg/g),  $Y_2$  is the predicted total phenolic or flavonoid content (mg/g),  $1/k_2$  is the adjustable parameter and  $t$  is extraction time (sec). Therefore, the extraction rate can be obtained by  $Y_2/k_2$  (mg/g.sec).

### Response Surface Methodology

In this study, pressure ( $X_1$ ), temperature ( $X_2$ ), and the ratio of ethanol ( $X_3$ ) were chosen to optimize the extraction rate of phenolic and flavonoid content during supercritical CO<sub>2</sub> extraction. Table 1 shows the Factors and Levels of the extraction, and experiment were designed using Box-Behnken Design (BBD) with one central point data. Design Expert 6.0.4 study as statistical software was applied for this Finally, the response surfaces of the parameters were analyzed by Analysis of Variance (ANOVA). The general second-order coefficients as shown in Equation 3.

$$Y = \beta_0 + \sum_{i=1}^k \beta_i X_i + \sum_{i=1}^k \beta_{ii} X_i^2 + \sum_{\sum_{i<j}^k} \beta_{ij} X_i X_j + \epsilon \quad X_2 X_k + \epsilon \tag{3}$$

**Table 1** Range of parameters and responses applied in ScCO<sub>2</sub> extraction assisted by ethanol.

Parameters	Unit	Range	Increment	Responses
Pressure, $X_1$	MPa	10 - 30	$\Delta X_1 = 10$	Extraction rate of TPC, $Y_1$ Extraction rate of TFC, $Y_2$
Temperature, $X_2$	°C	40 - 70	$\Delta X_2 = 15$	
Ratio of ethanol, $X_3$	%	2.5 - 7.5	$\Delta X_3 = 2.5$	

### Statistical Analysis

The determination of the most suitable model was based on the average absolute relative deviation (AARD) and the coefficient of determination ( $R^2$ ). A lower value of AARD and a high value of  $R^2$  show that the model successfully fits the experimental data. The equations of AARD and  $R^2$ , are shown as in Eq 3 and 4, respectively.

$$AARD (\%) = \frac{100}{n} \sum_{i=1}^n \left| \frac{E_{model} - E_{exp}}{E_{exp}} \right| \tag{4}$$

Based on Eq 3,  $n$  is the number of data points,  $E_{exp}$  is the value of experimental extraction rate data, and  $E_{model}$  is the calculated extraction data by using Brunner's or Esquivel's model.

$$R^2 = 1 - \frac{\sum_i (EM_i - ED_i)^2}{\sum_i (x_i - \bar{x})^2} \tag{5}$$

Based on Eq 4,  $\sum_i (EM_i - ED_i)^2$  is the residual data (i.e. an error between the model and the experimental data). Meanwhile,  $\sum_i (x_i - \bar{x})^2$  is the variance of the data.

**Table 2** Calculated parameters for the Applied Mathematical Models (modified Brunner's and Esquivel's Models) to fit the Total Phenolic Contents (TPC).

Run	X <sub>1</sub> (MPa)	X <sub>2</sub> (C)	X <sub>3</sub> (%)	Yield TPC (mg/g)	Brunner's Model					Esquivel's Model				
					k <sub>2</sub> (sec)	Y <sub>2</sub> (g)	Y <sub>2</sub> /k <sub>2</sub> (mg/g.sec)	AARD (%)	R <sup>2</sup>	k <sub>2</sub> (sec)	Y <sub>2</sub> (g)	Y <sub>2</sub> /k <sub>2</sub> (mg/g.sec)	AARD (%)	R <sup>2</sup>
1	30	55	2,5	279.87 ± 2.4	5452	438	0,080	6,497	0,986	286	0,000340	0,097	4,124	0,995
2	20	55	5	183.26 ± 3.1	1996	218	0,109	0,849	1,000	182	0,000447	0,081	2,064	0,999
3	10	55	7,5	112.22 ± 2.2	585492	7183	0,012	22,376	0,950	144	0,000157	0,023	12,357	0,983
4	20	70	2.5	166.26 ± 2.6	8079	309	0,038	3,164	0,996	203	0,000180	0,037	2,772	0,997
5	20	40	2,5	165.94 ± 2.7	4032	234	0,058	6,525	0,990	176	0,000278	0,049	4,955	0,995
6	20	70	7,5	88.57 ± 1.4	1252	99	0,079	1,784	0,998	88	0,000614	0,054	1,958	0,996
7	10	70	5	236.71 ± 1.3	5160	348	0,067	1,950	0,999	253	0,000230	0,058	3,467	0,998
8	10	40	5	122.19 ± 1.8	8600	229	0,027	14,048	0,989	152	0,000163	0,025	8,721	0,991
9	30	70	5	202.72 ± 2.6	4243	293	0,069	7,137	0,980	206	0,000409	0,084	4,877	0,993
10	20	40	7.5	102.52 ± 2.1	728	110	0,150	0,487	1,000	100	0,000823	0,083	1,653	0,998
11	10	55	2,5	127.77 ± 3.4	7076	211	0,030	1,326	0,999	100	0,000823	0,083	21,759	0,878
12	30	55	7,5	220.85 ± 1.4	904	243	0,269	1,133	0,999	217	0,000762	0,165	0,879	1,000
13	30	40	5	287.33 ± 3.4	1412	328	0,233	2,405	0,997	286	0,000575	0,164	0,680	1,000
Average							0,098	4,299	0,993			0,078	4,619	0,989

**Table 3** Calculated parameters for the Applied Mathematical Models (modified Brunner's and Esquivel's Models) to fit the Total Flavonoid Contents (TFC).

Run	X <sub>1</sub> (MPa)	X <sub>2</sub> (C)	X <sub>3</sub> (%)	Yield TFC (mg/g)	Brunner's Model					Esquivel's Model				
					k <sub>2</sub> (sec)	Y <sub>2</sub> (g)	Y <sub>2</sub> /k <sub>2</sub> (mg/g.sec)	AARD (%)	R <sup>2</sup>	k <sub>2</sub> (sec)	Y <sub>2</sub> (g)	Y <sub>2</sub> /k <sub>2</sub> (mg/g.sec)	AARD (%)	R <sup>2</sup>
1	30	55	2,5	477.91 ± 5.2	4150	683	0,165	6,510	0,990	491	0,000339	0,167	4,125	0,995
2	20	55	5	208.51 ± 2.1	4985	329	0,066	9,569	0,980	205	0,000440	0,090	2,070	0,999
3	10	55	7,5	42.79 ± 1.4	218806	1177	0,005	22,414	0,952	62	0,000156	0,010	12,376	0,983
4	20	70	2.5	49.40 ± 1.1	218806	1177	0,005	22,016	0,955	62	0,000156	0,010	6,410	0,996
5	20	40	2,5	353.26 ± 4.5	4881	534	0,109	6,472	0,988	363	0,000339	0,123	4,125	0,995
6	20	70	7,5	338.90 ± 3.4	1255	405	0,323	1,773	0,998	357	0,000613	0,219	1,959	0,996
7	10	70	5	35.32 ± 2.4	3554	44	0,012	3,729	0,996	38	0,000224	0,009	3,381	0,998
8	10	40	5	66.06 ± 0.9	17645	188	0,011	13,284	0,983	82	0,000163	0,013	8,722	0,991
9	30	70	5	189.84 ± 2.2	4008	270	0,067	7,156	0,981	203	0,000316	0,064	5,706	0,990
10	20	40	7.5	159.11 ± 1.1	713	170	0,238	0,567	1,000	158	0,000623	0,098	2,653	0,995
11	10	55	2,5	12.06 ± 0.3	7073	20	0,003	1,327	0,999	16	0,000623	0,010	40,782	0,917
12	30	55	7,5	91.62 ± 1.5	901	101	0,112	1,134	0,999	90	0,000768	0,069	0,981	1,000
13	30	40	5	353.83 ± 3.5	1421	405	0,285	2,404	0,997	351	0,000574	0,202	0,680	1,000
Average							0,098	8,037	0,985			0,085	6,015	0,991

## Results and Discussions

### Modelling

Brunner's and Esquivel's models were preferable for the extraction rate analysis as they require less adjustable parameters, and the models are easily fitted to experimental data [11]. Table 2 shows that the results of Brunner's and Esquivel's model fitted the recovery TPC and TFC from peanut skin by ScCO<sub>2</sub> assisted by ethanol. Meanwhile, Figure 1 and 2 show the example of Brunner's and Esquivel's model fitted the obtained data at a pressure of 10 to 30 MPa, temperature of 40 to 70 °C, and ethanol ratio of 2.5 to 7.5 %.

Based on the average coefficients of determination ( $R^2$ ) and average absolute relative deviation (AARD), the most suitable model was determined and presented in Table 2 and Table 3, respectively. The highest average coefficients of determination (i.e.  $R^2 \geq 1$ ) and the lowest average absolute relative deviation (AARD  $\leq 10\%$ ) presented the most suitable mathematical model. Table 2 indicates that Brunner's model gave better fitting for TPC recovery; thus, proved that it could provide more accurate data on the extraction rate of TPC compared to Esquivel's model. However, Table 3 shows the Esquivel's model suggested the best fitting for the extraction rate of TFC. Therefore, in this study, the calculated TPC and TFC extraction rate using Brunner's and Esquivel's model, respectively, were then applied in RSM for optimization. The maximum calculated extraction rate of TPC and TFC were later set to obtain the optimum parameters. Furthermore, the Brunner and Esquivel model is suitable to fit the experimental data of high-pressure instead of low-pressure condition as shown in Table 2 and 3. This is similar studies to the recovery of peanut skin oil that the models gives the low percentage of error to correlate the experimental data at high-pressure condition (30 MPa) [12]. The models also gives the similar results to give low percentage of error to correlate the waste of palm-fiber to obtain the extraction rate of carotenoid and tocopherol at high-pressure condition (30 MPa) [11].

### Statistical Analysis

The fixing the parameters studies were based on the preliminary studies, reviewing previous studies and the capacity of the ScCO<sub>2</sub> system. The limitation of parameters is that the maximum operating pressure has been restricted to 30 MPa due to the capacity of our ScCO<sub>2</sub> extraction system. Furthermore, the maximum temperature condition was restricted to 70 °C due to the prevention of degradation of flavonoid content [2]. Furthermore, the ratio of modifier was restricted to 7.5% due to the prevention the changing supercritical phase to subcritical phase [17].

The quadratic models have successfully to describe the experimental analysis due to the satisfactory coefficient ( $R^2$ ) of the quadratic model was over than 0.8, and the  $p$ -value was below than 0.05, as shown in Table 2 and Table 3, respectively. Furthermore, the final model is shown in Equations 6 and 7 and Table 2 and 3. Effects of the treatments variables, their interactions, and coefficients on the response variables were obtained by analysis of variance (ANOVA). This is the optimum conditions were 30 MPa, 40 °C and ratio of entrainer (ethanol) of 4.64% with a maximum extraction rate of the phenolic and flavonoid content of 0.22 mg/g.sec and 0.19 mg/g.sec, respectively. Furthermore, the pressure,  $X_1$  effect is the most significant effect on the extraction rate of phenolic due to the lowest value of  $p$ -value (0.0001). The parameter of  $X_1^2$ ,  $X_2^2$ ,  $X_3^2$  and  $X_2 X_3$  are not significant parameters on recovery of TPC due to high  $p$ -value ( $p$ -value  $>0.05$ ). In the recovery of TFC, pressure and quadratic pressure are only the significant factor based on  $p$  value ( $p$ -value  $< 0.05$ ).

$$Y_1 = 0.11 + 0.064 X_1 - 0.027 X_2 + 0.038 X_3 + 0.00288 X_1^2 - 0.013 X_2^2 - 0.015 X_3^2 - 0.051 X_1 X_2 + 0.051 X_1 X_3 - 0.013 X_2 X_3 \tag{6}$$

$$Y_2 = 0.09 + 0.058 X_1 - 0.023 X_2 + 0.00458 X_3 - 0.04 X_1^2 + 0.022 X_2^2 + 0.013 X_3^2 - 0.033 X_1 X_2 + 0.024 X_1 X_3 - 0.071 X_2 X_3 \tag{7}$$





Figure 1 Brunner's and Esquivel's model fitted the experimental data of total phenolic content





**Figure 2** Brunner's and Esquivel's model fitted the experimental data of total flavonoid content



**Table 4.** ANOVA table for the responses.

Source	Sum of square	df	Mean Square	F	p-value
Extraction rate of TPC (mg/g.sec)					
Regression	0.074	9	0.0081	37.57	<0.0001
Residual	0.0015	7	0.00021		
Total	0.75	16			
Extraction rate of TFC (mg/g.sec)					
Regression	0.067	9	0.0073	6.44	0.011
Residual	0.0081	17	0.0012		
Total	0.75	26			


**Table 5.** Coefficient of polynomial extraction rate of TPC and TFC.

Extraction rate, (mg/g.s)	Coefficient, (β)		p-value	Extraction rate, (mg/g.s)	Coefficient, (β)		p-value
	TPC, Y <sub>1</sub>				TFC, Y <sub>2</sub>		
Intercept	0.11			Intercept	0.090		
X <sub>1</sub>	0.064	< 0.0001		X <sub>1</sub>	0.058	0.0020	
X <sub>2</sub>	-0.027	0.0014		X <sub>2</sub>	-0.023	0.0947	
X <sub>3</sub>	0.038	0.0002		X <sub>3</sub>	0.00458	0.7165	
X <sub>1</sub> <sup>2</sup>	0.00288	0.7009		X <sub>1</sub> <sup>2</sup>	-0.040	0.0472	
X <sub>2</sub> <sup>2</sup>	-0.013	0.1073		X <sub>2</sub> <sup>2</sup>	0.022	0.2349	
X <sub>3</sub> <sup>2</sup>	-0.015	0.0830		X <sub>3</sub> <sup>2</sup>	0.013	0.4501	
X <sub>1</sub> X <sub>2</sub>	-0.051	0.0002		X <sub>1</sub> X <sub>2</sub>	-0.033	0.0917	
X <sub>1</sub> X <sub>3</sub>	0.051	0.0002		X <sub>1</sub> X <sub>3</sub>	-0.024	0.1940	
X <sub>2</sub> X <sub>3</sub>	-0.013	0.1267		X <sub>2</sub> X <sub>3</sub>	0.071	0.0041	
	R <sup>2</sup> = 0.972				R <sup>2</sup> =0.8923		

**Parameter Effects**

Figure 3 shows that increased in pressure from 10 to 30 MPa had significantly enhanced the extraction rate of TPC at constant temperatures of 40 °C and 70 °C. This is because the density, solubility, and diffusivity effect are dominant in the extraction process [4]. The high-pressure condition will enhance the density of solvents; thus, high interaction of the solvent molecules will then increase the dissolution of the solute [17] [18]. As a result, the increase in density enhances the solubility and diffusivity [19]. Furthermore, Figure 3 shows the increase in TPC with pressure is more evidenced at 40 C than at 70 C due to the vapor pressure and density. At pressure of 10 MPa and temperature 70 °C, the vapor pressure effect was occurred in this condition, thus the recovery of TPC was high. Contractively in pressure of 30 MPa and temperature of 70 °C, the effect of reducing of solubility was occurred instead of vapour pressure in this condition, therefore the extraction rate of TPC was not significantly increased.

Increase in the rate of ethanol from 2.5 to 7.5% also could enhance the extraction rate of TFC at a constant temperature of 55 °C and pressure of 10 MPa, as shown in Figure 4(b). The addition of ethanol increases the polarity of carbon dioxide; thus, the flavonoid – a polar compound, is easily soluble. However, an increase in the rate of ethanol will slightly decrease the extraction rate of TFC at constant pressure and temperature of 30 MPa and 55 °C, respectively (Figure 3(b)). This is due to ethanol could prevent the ScCO<sub>2</sub> from diffusing into the peanut skin and increasing the rate of ethanol would reduce the mixture density. The compactness of material inside of the vessel was also occurred due to in the high pressure condition, thus the yield of flavonoid content will decrease. Maran, Manikandan [32] also found that extraction pressure and entrainer flow rate give significant effect to obtain high flavonoid contents from tea (*Camellia sinensis L.*) leaves due to high solubility and polarity of solvent.



**Figure 3** Response of the effect of variables on extraction rate of TPC (mg/g.sec) at (a) constant ratio of ethanol 5% and (b) constant temperature 55 °C.

The recovery of TPC from fermented orange pomace by supercritical carbon dioxide also has similar results, whereby pressure was found to be the most significant factor to enhance the TPC recovery [20]. Adil, Cetin [21] also found that pressure could give a significant effect compared to other parameters for enhancing the recovery of phenolic contents from apple and peach pomaces. Guo-qing, Hao-ping [22] found that high pressure encourages the bulk density of the fluid, which increased the solubility. However, as pressure increases further, the fluid is less compressible, thus has an insignificant impact in bulk density. However, Duba and Fiori [23] revealed contradictive results by which the pressure over 35 MPa has no significant effect on the recovery of flavanol/phenolic compounds. Machmudah, Zakaria [24] also found that the selectivity of ScCO<sub>2</sub> is also weakened by high-pressure condition (i.e. 40 MPa), as all the studied compounds will be extracted.


An increase in temperature (40 to 70 °C) at a constant pressure of 10 MPa slightly enhanced the extraction rate of TPC at constant rate of ethanol 5% from 0.027 to 0.067 mg/g.sec. The sublimation pressure is dominant during the low-pressure condition; thus, the TPC is easily diluted in the mixture

solvent [18]. High temperature condition (70 °C) also escalates the diffusivity, thus, the solvating power of solvents will be enhanced to extract phenolic content. However, at constant pressure of 30 MPa, enhancement of temperature decreased the extraction rate of TPC from 0.233 mg/g.sec to 0.069 mg/g at constant rate of ethanol 5%. as shown in Figure 3(a). This is due to a high-temperature could decrease the density of the solvent; thus, the solubility of solutes and diffusivity effect will be alleviated [25, 26]. Therefore, the solvation power of solvent will reduce to carry out the phenolic contents. Meanwhile, Figure 3(b) shows that increase in the ratio of ethanol (2.5 to 7.5%) does not give a significant impact on the enhancement of the TPC extraction rate at constant pressures of 10 and 30 MPa. This is contradictive that phenolic compounds are easily diluted/dissolved in a polar solvent compared to the nonpolar solvents. Hasmida, Liza [27] also found that the effect of entrainer does not gives the significant effect to enhance the recovery of phenolic content in the extraction of *Quercus infectoria* galls. Molino *et al.* [29] also found the contradictive results that effectiveness of ethanol as an entrainer for astaxanthin extraction was not significant due to the selectivity of solvents. This is contradictive result with the recovery of tocopherol and carotenoids from wasted palm-fiber that increase of entrainer ratio (5 to 7.5 %) enhance the yield [11]. This is due to the swelling matrix of raw material by ethanol; thus, the supercritical carbon dioxide is easily to penetrate the raw material. On the other hand, the presence of ethanol was found to be effective for the extraction of lutein, as an enhancement was observed both in terms of recovery and purity. Thus, the phenolic was unsuccessful extracted due to the ethanol is not enough to swell peanut skin.

A similar trend of TPC extraction rate is shown as in Figure 4(a). Increase in pressure (10 to 30 MPa) encouraged TFC rate at a constant ethanol ratio of 5% and temperatures of 40 and 70 °C. Molino, Mehariya [28] found that the density of solvents could play a significant role in enhancing flavonoid recovery. Putra, Rizkiyah [29] also reported that the increase in pressure could be resulting in the enhanced extraction of polar and high-molecular-mass compounds such as polyphenols (i.e. the major antioxidants in plant extracts) suggesting the significance of operating at lower temperature and higher pressure. Ouédraogo, Dicko [30] also found that to enhance the recovery of flavonoids from *Odontonema strictum* leaves with high antioxidant activity by increasing of pressure (200 Bar).

Besides that, increase of temperature from 40 to 70 °C also could decrease the extraction rate of TFC from 0.185 to 0.041 mg/g.sec, at a constant pressure of 30 MPa, 5% ratio of ethanol and temperature 40 to 70 °C, as shown in Figure 4(a). Due to the low density of ScCO<sub>2</sub>, the alleviation of solubility relies on the increase in temperature. On the other hand, temperature affects the volatility of the solute; therefore, it is difficult to predict the outcome. The similar results with the recovery of flavonoid from *Odontonema strictum* leaves that increase temperature condition (55 to 65 °C) decrease the flavonoid extract [30]. A higher temperature (65 °C) accelerate mass transfer to improve the extraction yield by enhancement of the solubility of the solute. In contrast, increase in temperature (40 to 60 °C) increased the solvating power when applying ScCO<sub>2</sub> assisted by ethanol in recovery of flavonoid content from brewer's spent grain. The predominance of the vapor pressure effect depended on the high-temperature condition to enhance the extraction rate of flavonoid content [31].

Increase in the rate of ethanol from 2.5 to 7.5% also could enhance the extraction rate of TFC at a constant temperature of 55 °C and pressure of 10 MPa, as shown in Figure 4(b). The addition of ethanol increases the polarity of carbon dioxide; thus, the flavonoid – a polar compound, is easily soluble. However, an increase in the rate of ethanol will slightly decrease the extraction rate of TFC at constant pressure and temperature of 30 MPa and 55 °C, respectively (Figure 3(b)). This is due to ethanol could prevent the ScCO<sub>2</sub> from diffusing into the peanut skin and increasing the rate of ethanol would reduce the mixture density. The compactness of material inside of the vessel was also occurred due to in the high pressure condition, thus the yield of flavonoid content will decrease. Maran, Manikandan [32] also found that extraction pressure and entrainer flow rate give significant effect to obtain high flavonoid contents from tea (*Camellia sinensis L.*) leaves due to high solubility and polarity of solvent.



**Figure 4** Response of the effect of variables on extraction rate of TFC (mg/g.sec) at (a) constant ratio of ethanol 5% and (b) constant temperature 55 °C.

## Conclusion

In this research, Brunner's model gives better fitting for TPC recovery to obtain its extraction rate, as compared to Esquivel's. In contrast, Esquivel's model gives better fitting for TFC recovery. Therefore, the calculated TPC extraction rate in Brunner's model and TFC extraction rate in Esquivel's model were used for optimization. The quadratic polynomial showed that the model was successfully fitted to the extraction rate of TPC and TFC. Based on the optimization study using RSM, the best conditions were reported to be at 30 MPa, 40 °C and rate of ethanol of 4.64%, with a maximum phenolic and flavonoid extraction rate content of 0.22 mg/g.sec and 0.19 mg/g.sec, respectively. Nevertheless, this study found that increasing pressure to a certain level would enhance the extraction rate of phenolic and flavonoid contents.

## Acknowledgement

The authors acknowledged the Centre of Lipid Engineering and Applied Research (CLEAR), Universiti Teknologi Malaysia (UTM) for providing research facilities to conduct this study. We also would like to thank professional development research Universiti Teknologi Malaysia grant (R.J130000.7113.05E53) to support this work.

## References

- [1] Putra, N.R., et al., *Effect of particle size on yield extract and antioxidant activity of peanut skin using modified supercritical carbon dioxide and soxhlet extraction*. Journal of Food Processing and Preservation, 2018. **42**(8): p. e13689.
- [2] Putra, N.R., et al., *Recovery and solubility of flavonoid and phenolic contents from Arachis Hypogea in supercritical carbon dioxide assisted by ethanol as cosolvent*. Journal of Food Processing and Preservation, 2020. **44**(10): p. e14768.
- [3] Chanda, S., D. Bhayani, and D. Desai, *Polyphenols and flavonoids of twelve Indian medicinal plants*. The bioscan, 2013. **8**(2): p. 595-601.
- [4] Putra, N.R., M.A. Che Yunus, and S. Machmudah, *Solubility model of arachis hypogea skin oil by modified supercritical carbon dioxide*. Separation Science and Technology, 2019. **54**(5): p. 731-740.
- [5] Putra, N.R., et al., *Comparison extraction of peanut skin between CO2 supercritical fluid extraction and soxhlet extraction in term of oil yield and catechin*. Pertanika Journal of Science & Technology, 2018. **26**(2).
- [6] Radfar, S. and S.M. Ghoreishi, *Experimental extraction of L-Carnitine from oyster mushroom with supercritical carbon dioxide and methanol as co-solvent: Modeling and optimization*. Journal of Supercritical Fluids, 2018. **140**: p. 207-217.
- [7] Hrnčič, M.K., et al., *Extracts of White and Red Grape Skin and Rosehip Fruit: Phenolic Compounds and their Antioxidative Activity*. Acta Chimica Slovenica, 2019.
- [8] Garcia-Mendoza, M., et al., *Extraction of phenolic compounds and anthocyanins from juçara (Euterpe edulis Mart.) residues using pressurized liquids and supercritical fluids*. Journal of Supercritical Fluids, 2017. **119**: p. 9-16.
- [9] Castro-Vazquez, L., et al., *Bioactive flavonoids, antioxidant behaviour, and cytoprotective effects of dried grapefruit peels (Citrus paradisi Macf.)*. Oxidative medicine and cellular longevity, 2016. **2016**.
- [10] Bimakr, M., et al., *Optimization of Supercritical Carbon Dioxide Extraction of Bioactive Flavonoid Compounds from Spearmint (Mentha spicata L.) Leaves by Using Response Surface Methodology*. Food and Bioprocess Technology, 2012. **5**(3): p. 912-920.
- [11] Putra, N.R., et al., *Recovery of valuable compounds from palm-pressed fiber by using supercritical CO<sub>2</sub> assisted by ethanol: modeling and optimization*. Separation Science and Technology, 2019: p. 1-14.
- [12] Putra, N.R., et al., *Extraction of peanut skin oil by modified supercritical carbon dioxide: Empirical modelling and optimization*. Separation Science and Technology, 2018. **53**(17): p. 2695-2703.
- [13] Rosli, N.L., et al. *Phytochemical analysis and antioxidant activities of Trigona Apicalis propolis extract*. in *AIP Conference Proceedings*. 2016. AIP Publishing.
- [14] Silva, F., et al., *Physicochemical properties, antioxidant activity and stability of spray-dried propolis*. J. ApiProd. ApiMed. Sci., 2011. **3**: p. 94-100.
- [15] Esquivel, M., M. Bernardo-Gil, and M. King, *Mathematical models for supercritical extraction of olive husk oil*. The Journal of Supercritical Fluids, 1999. **16**(1): p. 43-58.
- [16] Brunner, G., *Gas extraction: an introduction to fundamentals of supercritical fluids and the application to separation processes*. Vol. 4. 2013: Springer Science & Business Media.
- [17] Machmudah, S., et al., *Extraction of astaxanthin from Haematococcus p luvialis using supercritical CO<sub>2</sub> and ethanol as entrainer*. Industrial & engineering chemistry research, 2006. **45**(10): p. 3652-3657.
- [18] Putra, N.R., et al., *Formulation and evaluation of a new semi-empirical model for solubility of plant extracts in supercritical carbon dioxide assisted by ethanol as co-solvent*. Chemical Engineering Communications, 2020: p. 1-9.
- [19] Peng, W.L., et al., *Optimization of process variables using response surface methodology for tocopherol extraction from Roselle seed oil by supercritical carbon dioxide*. Industrial Crops and Products, 2020. **143**: p. 111886.
- [20] Espinosa-Pardo, F.A., et al., *Extraction of phenolic compounds from dry and fermented orange pomace using supercritical CO<sub>2</sub> and cosolvents*. Food and Bioprocess Processing, 2017. **101**: p. 1-10.
- [21] Adil, I.H., et al., *Subcritical (carbon dioxide+ ethanol) extraction of polyphenols from apple and peach pomaces, and determination of the antioxidant activities of the extracts*. The journal of supercritical fluids, 2007. **43**(1): p. 55-63.
- [22] Guo-qing, H., et al., *Optimization of conditions for supercritical fluid extraction of flavonoids from hops (Humulus lupulus L.)*. Journal of Zhejiang University Science B, 2005. **6**(10): p. 999.
- [23] Duba, K. and L. Fiori, *Supercritical CO<sub>2</sub> extraction of grape seeds oil: scale-up and economic analysis*. International Journal of Food Science & Technology, 2019. **54**(4): p. 1306-1312.
- [24] Machmudah, S., et al., *Lycopene extraction from tomato peel by-product containing tomato seed using supercritical carbon dioxide*. Journal of Food Engineering, 2012. **108**(2): p. 290-296.
- [25] Ahmed, Z., et al., *Extraction and Modeling of Algerian Rosemary Essential Oil Using Supercritical CO<sub>2</sub>: Effect of Pressure and Temperature*. Energy Procedia, 2012. **18**: p. 1038-1046.
- [26] Hasmdia, et al., *Solubility correlation of gall (Quercus infectoria) extract in supercritical CO<sub>2</sub> using semi-*



- empirical equations*. Asia-Pacific Journal of Chemical Engineering, 2017. **12**(5): p. 790-797.
- [27] Hasmida, M., et al., *Total phenolic content and antioxidant activity of quercus infectoria galls using supercritical CO2 extraction technique and its comparison with soxhlet extraction*. Pertanika Journal of Science & Technology, 2015. **23**(2): p. 287-295.
- [28] Molino, A., et al., *Extraction of astaxanthin and lutein from microalga Haematococcus pluvialis in the red phase using CO2 supercritical fluid extraction technology with ethanol as co-solvent*. Marine drugs, 2018. **16**(11): p. 432.
- [29] Putra, N.R., et al., *Recovery and solubility of flavonoid and phenolic contents from Arachis Hypogea in supercritical carbon dioxide assisted by ethanol as cosolvent*. Journal of Food Processing and Preservation, 2020: p. e14768.
- [30] Ouédraogo, J.C.W., et al., *Enhanced extraction of flavonoids from Odontonema strictum leaves with antioxidant activity using supercritical carbon dioxide fluid combined with ethanol*. The Journal of Supercritical Fluids, 2018. **131**: p. 66-71.
- [31] Spinelli, S., et al., *Supercritical carbon dioxide extraction of brewer's spent grain*. The Journal of Supercritical Fluids, 2016. **107**: p. 69-74.
- [32] Maran, J.P., et al., *Box-Behnken design based multi-response analysis and optimization of supercritical carbon dioxide extraction of bioactive flavonoid compounds from tea (Camellia sinensis L.) leaves*. Journal of Food Science and Technology, 2015. **52**(1): p. 92-104.