

ORIGINAL ARTICLE

Pullulan Production Utilizing Various Carbon Sources by *Aureobasidium melanogenum* DSM 2404 as Biotherapeutic Tool in Biomedical Applications

*Daniel Joe Dailin^{1,2}, Luo Zaini Mohd Izwan Low², Siti Fatimah Zaharah Mohd Fuzi³, Tong Woei Yenn⁴, Nor Hasmaliana Abdul Manas^{1,2}, Samina Mehnaz⁵, Hesham El Enshasy^{1,2,6}

¹ Institute of Bioproduct Development, Universiti Teknologi Malaysia, 81310, Skudai, Johor, Malaysia.

² Bioprocess and Polymer Engineering Department, Faculty of Chemical and Energy Engineering, Universiti Teknologi Malaysia, 81310, Skudai, Johor, Malaysia.

³ Department of Technology & Natural Resources, Faculty of Applied Science & Technology, UTHM, Pagoh Campus, 84600 Panchor, Johor.

⁴ Universiti Kuala Lumpur, Institute of Medical Science Technology (UniKL MESTECH), Clinical Laboratory Section, A1, 1, Jalan TKS 1, Taman Kajang Sentral, 43000 Kajang, Selangor, Malaysia.

⁵ School of Life Sciences, Forman Christian College (A Chartered University), Lahore 54600, Pakistan.

⁶ Bioprocess Development Department, City for Scientific Research and Technology Applications (CSAT), New Burg Al Arab, Alexandria, Egypt.

ABSTRACT

Introduction: Pullulan is a biodegradable biopolymer made up of maltotriose subunits that are water-soluble. It has a direct glucan structure comprising α -1,4 and α -1,6 linkages in a 2:1 ratio, which gives it exceptional physical and structural properties, making it a suitable choice for application in biomedical applications. Despite its potential benefits, the price of production is expensive, and productivity is low, which are significant drawbacks. Therefore, the objective of this work is to identify the ideal carbon source for *Aureobasidium melanogenum* DSM 2404 to produce high pullulans. **Methods:** In shake flask studies, the effects of different carbon sources, such as maltose, glucose, sucrose, lactose, and xylose, on cell growth and pullulan generation by *Aureobasidium melanogenum* DSM 2404 were examined. **Results:** The highest pullulan production was obtained from sucrose (13.38 g L⁻¹) compared to other types of carbon sources used in the fermentation medium. The maximal pullulan production rate of 0.112 [g⁻¹ L⁻¹ h⁻¹] was obtained in sucrose culture. This suggests that sucrose provides favorable conditions for the microorganisms to produce pullulan at a faster rate compared to other sugars. **Conclusion:** Sucrose was found to be the most efficient carbon source for the synthesis of pullulan using *Aureobasidium melanogenum* DSM 2404 generating 13.38 g L⁻¹ of pullulan.

Malaysian Journal of Medicine and Health Sciences (2023) 19(SUPP9): 263-268. doi:10.47836/mjmhs.19.s9.36

Keywords: Pullulan; Production; Carbon sources; *Aureobasidium melanogenum*; Biotherapeutic

Corresponding Author:

Daniel Joe Dailin, PhD
Email: jddaniel@utm.my
Tel: +607-5534369

INTRODUCTION

Polysaccharides, often known as a type of natural macromolecular polymer typically consist of chains with either straight or branched glycosidic linkages to interconnect more than ten monosaccharides. Its molecular weight (Mw) can be hundreds or a billion times higher [1]. There are three types of polysaccharides such as capsular polysaccharides (CPSs), lipopolysaccharides (LPSs) and xopolysaccharides

(EPSs) [2]. The majority of CPSs are linked to bacteria's pathogenicity and virulence enhancing factors [3]. While for LPSs, most Gram negative bacteria's outer membrane contains LPSs, and some bacteria create LPSs as a virulence factor [4,5]. Lastly, EPSs are produced outside of a cell (extracellular) and this polymer is mostly high Mw [6]. Currently, there are many important EPSs for industrial applications such as kefirin [7], xanthan [8] and pleuran [9]. Microbial polysaccharides have gained significant attention due to their unique properties and wide range of applications in various industries.

One of the most studied and industrialized EPSs named pullulan. It has garnered significant attention

from researchers and industries alike due to its unique properties. During the fermentation of *Aureobasidium pullulans*, Bauer [10] first examined pullulan. Later, Bernier [11] successfully extract and identify pullulan from the culture solutions of *A. pullulans*. According to Bender's investigations [12], which showed that it was exclusively composed of α -D-glucans with a preponderance of α -(1 \rightarrow 4) linkages and had a positive optical rotation, the term "pullulan" was given. It is highly soluble in water, forming clear solutions, and can form films with excellent oxygen and moisture barrier properties. Pullulan films are also flexible, transparent, and have good mechanical strength.

The pullulan market has experienced significant growth in recent years due to its versatile properties and wide range of applications. Pullulan's unique characteristics, such as film-forming ability, solubility, and biocompatibility, have made it a sought-after ingredient in various industries. According to market analysts, from 2017 the pullulan market is anticipated to develop at a compound annual growth rate (CAGR) of about 2.2 percent, beginning with a value of US\$129 million and reaching US\$130 million by 2023[13]. Additionally, according to Adroit Research Market [14] the COVID-19 pandemic has caused the pullulan market size to generally be evaluated to be worth US\$ 68 million in 2022 and to increase to US\$ 89 million by 2029 with a CAGR of 4.6% during the operational time window 2022-2028. Pharmaceutical firms currently control the largest industries, accounting for about 40.74 percent of the global market share in 2015. The largest market was Japan, with a market share of 667 million tonnes, followed by China (20.65%) and the USA (29.65%) [13]. Overall, the pullulan market is expected to continue growing as industries recognize its potential in various applications. The demand for natural and sustainable ingredients, coupled with the expanding food, pharmaceutical, and packaging sectors, is likely to drive the market's future growth.

The applications of pullulan in biomedical industries are extensive [15]. Pullulan is inert and has non-toxic, non-immunogenic, and biocompatible characteristics that make it broadly used in the biomedical field [16]. Pullulan has a high concentration of hydroxyl groups which could be used as a bioactive polymer and dextran-based blood plasma substitute [17]. Moreover, it can also be shaped into pills or tablets size for the use of pharmaceutical and nutraceutical purposes, such as NPcaps capsules from Capsugel® company, US [18]. Due to pullulan wide applications, it is required to identify the best production strategy for the high production of pullulan. It is worth noting that pullulan production strategies may vary depending on the specific application and industry requirements. Therefore, a

thorough understanding of the production process, continuous monitoring, and optimization efforts are necessary to develop the most efficient strategy for high pullulan production. Identifying the best production strategy for high-yield pullulan production is crucial to meet the increasing demand for this versatile polysaccharide. Several factors need to be considered when developing an optimal production strategy, including the choice of microbial strain, fermentation conditions, and downstream processing. By comparing various types of carbon sources, this study sought to identify the best carbon source for *Aureobasidium melanogenum* DSM 2404 to produce pullulan. The study would typically involve selecting a range of carbon sources to evaluate their impact on pullulan production. Common carbon sources tested for pullulan production include maltose, glucose, sucrose, lactose, and xylose. Each carbon source has a different composition and influence pullulan production differently.

MATERIALS AND METHODS

Microorganism

The research employed *Aureobasidium melanogenum* DSM 2404 (formerly *Aureobasidium pullulans* var. *melanogenum*), which was obtained in freeze-dried form from the Leibniz Institute DSMZ – Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH (German Collection of Microorganisms and Cell Cultures GmbH).

Preparation of Inoculum and production medium

One mL of a functioning cell bank containing the fungal culture was added to a 250 mL Erlenmeyer flask that contained 50 mL of sterile seed media (SM) to create the inoculum[19]. The SM was made up of the following components per liter of distilled water: sucrose (30.0 g), $(\text{NH}_4)_2\text{SO}_4$ (0.6 g), yeast extract (0.4 g), K_2HPO_4 (5.0 g), $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ (0.2 g), NaCl (1.0 g) and MnCl_2 (0.01 g), with the pH adjusted to 5.5 before sterilization for 20 minutes at 121°C. Separately, sucrose was sterilized for 20 minutes at 110°C and added to the Erlenmeyer flask containing the 50 mL sterile SM after cooling. The inoculated flask was then incubated for 48 hours at 200 rpm and 28°C in a rotary shaker incubator. After 48 hours of incubation, 5% of the inoculum was transferred into the production medium, which contained the following components per liter of distilled water: maltose (50.00 g), yeast extract (10.00 g), peptone (20.00 g), $(\text{NH}_4)_2\text{SO}_4$ (0.60 g), K_2HPO_4 (5.00 g), $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ (0.60 g), and NaCl (1.00 g).

Effect of Different Types of Carbon Sources

In order to find the optimal carbon source for *Aureobasidium melanogenum* DSM 2404 to produce high pullulans, various carbon sources including maltose, glucose, sucrose, lactose, and xylose were

examined. The production medium composed of (in g L⁻¹): yeast extract, 10.00; peptone, 20.00; (NH₄)₂SO₄, 0.60; K₂HPO₄, 5.00; MgSO₄·7H₂O, 0.60; NaCl, 1.00; was used in this study. Separately, each carbon source was sterilized for 20 minutes at 110°C, which later was transferred into the production medium. After inoculating the production medium with 5% of the inoculum, for 48 hours at 200 rpm and 28°C, the flask was incubated in a rotating shaker incubator.

Cell Dry Weight Determination

Samples in all flasks containing 50 ml each were withdrawn after 72 hours of cultivation in a centrifugation tube. The culture broth was then centrifuged in 50 ml falcon tubes at 7000 rpm for 15 minutes to settle the cells. Gravimetric methods were used to determine the cell dry weight.

Crude EPS Determination

Crude EPS was precipitated and separated from the supernatant. Then it underwent further analysis using method given by [20]. A 50 mL conical centrifuge tube was filled with 10 mL of supernatant and two volumes (20 mL) of pre-chilled 95 percent ethanol to extract the EPS. The ethanol-supernatant mixture was then chilled at 4°C overnight to precipitate the EPS. The ethanol-supernatant combination was centrifuged the following day at 27°C for 10 minutes at 7000 rpm. After removing the supernatant, the EPS was dried at 80°C overnight before being weighed.

RESULTS

Two monosaccharides (glucose and xylose) and three disaccharides (maltose, sucrose, and lactose) were studied for their ability to support the production of CDW and pullulan. The results of CDW and pullulan production after five days

of cultivation was presented in Figure 1. Control indicates cultivation medium without any C source. For CDW production, the highest yield was obtained from glucose (13.08 g L⁻¹), continued via maltose (11.73 g L⁻¹), sucrose (9.94 g L⁻¹), lactose (5.73 g L⁻¹), and xylose (5.51 g L⁻¹). For pullulan production, the highest yield was obtained from sucrose (13.38 g L⁻¹), continued via maltose (12.13 g L⁻¹), glucose (5.70 g L⁻¹), xylose (2.35 g L⁻¹), and lactose (1.71 g L⁻¹). The highest cell productivity of 1.35 g g⁻¹ was achieved by using sucrose as the sole C source, while the lowest was 0.30 g g⁻¹ from lactose. The pH readings are ranging from 5.09 to 6.77.




Figure 1 : The results of CDW and pullulan production after five days of cultivation.

Table I : Kinetic parameters for cell growth and pullulan production during batch cultivation of *Aureobasidium melanogenum* in shake flask culture (data were taken after 120 hours cultivation, and n=3)

Carbon Source	X _{max} [g L ⁻¹]	P _{max} [g L ⁻¹]	dx/dt [g ⁻¹ L ⁻¹ h ⁻¹]	dp/dt [g ⁻¹ L ⁻¹ h ⁻¹]	Y _{p/x} [g g ⁻¹]
Control	3.51±0.18	1.61±0.02	0.029	0.013	0.459
Maltose	11.73±0.35	12.13±0.1	0.098	0.101	1.034
Glucose	13.08±0.11	5.70±0.03	0.109	0.048	0.436
Sucrose	9.93±0.05	13.38±0.18	0.082	0.112	1.346
Lactose	5.72±0.04	1.71±0.02	0.048	0.014	0.299
Xylose	5.51±0.01	2.40±0.07	0.046	0.020	0.427

Notes: X_{max}: maximal cell dry weight; P_{max}: maximal pullulan production; dx/dt: growth rate; dp/dt: production rate; Y_{p/x}: Specific production.

To comprehend the impact of the carbon source on the kinetics of cell development and pullulan formation, Table 1 summarized the influence of carbon source on the overall primary and secondary kinetic parameters during cultivation at the shake flask level.

According to table 1, the cell development rate and production rate were highly influenced by the type of sugar used during fermentation. The maximal pullulan production rate of $0.112 \text{ [g}^{-1} \text{ L}^{-1} \text{ h}^{-1}]$ was obtained in sucrose culture whereas, the lowest value of 0.014 was in lactose culture. This suggests that sucrose provides favorable conditions for the microorganisms to produce pullulan at a faster rate compared to other sugars. In contrast, lactose resulted in the lowest pullulan production rate of 0.014. On the other hand, a maximal growth rate of $0.109 \text{ [g}^{-1} \text{ L}^{-1} \text{ h}^{-1}]$ was attained in glucose culture followed by maltose culture. This implies that glucose and maltose are more efficiently utilized by the microorganisms for their growth during fermentation compared to other sugars.

DISCUSSION

Aureobasidium sp. was found to be able to utilize various C sources [20, 21]. Most of the studies found that the pullulan yield was higher when using sucrose as a C source [20, 22, 23]. Sucrose is the ideal C source for pullulan synthesis in *A. pullulans* CGMCC1234, according to Sheng et al [24]. In his research, he studied the relationships between different C sources and the β -fructofuranosidase enzyme's activity. In reduced concentrations of sucrose, this enzyme catalyzes sucrose to glucose and fructose. Furthermore, this enzyme can aid in kestose synthesis by combining fructose with sucrose at a higher sucrose concentration [25]. Sheng and associates found that the activity of β -fructofuranosidase was highest when sucrose was employed as the carbon source, and the lowest value was noted when the strain was grown in media containing different carbon sources. The extracellular fluid's osmotic stress can be reduced by increasing the level of this enzyme, which promotes *A. pullulans*' growth rate and causes it to release more pullulan in the initial stages of fermentation. Besides that, sucrose is a disaccharide composed of glucose and fructose [26]. It has a relatively high carbon content, which makes it a suitable substrate for microbial growth and polysaccharide production [27]. Pullulan is a polysaccharide consisting of repeating maltotriose units (three glucose units), and sucrose provides the necessary carbon atoms for the biosynthesis of pullulan [28]. Sucrose is relatively inexpensive compared to other carbon sources, such as glucose or maltose [29]. Its low cost and ready availability make it an attractive choice for industrial-scale

pullulan production, as it helps keep the production costs down.

However, higher pullulan production can be also be achieved from other C sources such as glucose [30, 31, 32]. Glucose, a monosaccharide, can also be utilized as a carbon source for pullulan production [33]. It can be obtained from various sources, such as corn syrup or glucose syrup. Some pullulan-producing microorganisms, such as *Aureobasidium pullulans*, have the ability to metabolize glucose efficiently for pullulan synthesis [34]. Elevated levels of glucosyltransferase, UDPG-pyrophosphorylase and α -phosphoglucose mutase activities remained observed when *A. pullulans* Y68 was cultured in the presence of glucose [14]. These enzymes were found to be involved in the pullulan synthesis process. Although glucose act as a potential C source to produce high CDW, the pullulan production from glucose is lower than sucrose. Hence, sucrose is used as a potential C source and undergoes further optimization.

The maximal specific production of pullulan of $1.34 \text{ [g g}^{-1}]$ was obtained in sucrose culture followed by maltose culture. This indicates that sucrose and maltose promote higher pullulan production relative to the amount of biomass formed by the microorganisms. This finding showed similar result from previous in which high sucrose and maltose concentration was used to support high pullulan production [35].

These results suggest that different sugars provide varying conditions for pullulan production. Sucrose seems to be particularly effective in promoting both pullulan production rate and specific production, while glucose and maltose favor microbial growth rate. Lactose, on the other hand, appears to be less favorable for pullulan production based on the lower production rate observed. It is important to note that these results are specific to the study or experiment mentioned and may vary depending on the microbial strain, fermentation conditions, and other factors. Further research and experimentation are necessary to confirm and generalize these findings in different contexts

CONCLUSION

This study investigate the effect of various carbon sources on *Aureobasidium melanogenum* DSM 2404 growth and its production of pullulan in shake flasks. According to the findings, sucrose was the best carbon source for pullulan synthesis as demonstrated by *Aureobasidium melanogenum* DSM 2404's output of 13.38 g L^{-1} after 120 hours of growth. On the other hand, the maximal biomass of 13.08 g L^{-1} was obtained in glucose culture. Further studies are now

going in our laboratories for in-depth study of the biosynthesis regulation using mixed substrate at the bioreactor level for bioprocess industrialization of this important polysaccharide.

ACKNOWLEDGMENT

The authors would like to thank the Ministry of Higher Education, Malaysia through FRGS grant No. (FRGS/1/2020/TK0/UTM/02/16).

REFERENCES

- Adroit Market Research. Pullulan market size, share, trends, and forecast 2021-2028; 2020.
- Al-Tabakha MM. HPMC capsules: current status and future prospects. *J Pharm Pharm Sci.* 2010; 13(3):428-442. doi: 10.18433/j3k881.
- Bauer R. Physiology of *Dematiom pullulans* de Bary. *Zentralbl Bacteriol Parasitenkd Infektionskr Hyg Abt2.* 1938;98:133-167.
- Bender H, Lehmann J, Wallenfels K. Pullulan, an extracellular glucan from *Pullularia pullulans*. *Biochim Biophys Acta.* 1959;36:309–316. doi: 10.1016/0006-3002(59)90172-6.
- Bernier B. The production of polysaccharides by fungi active in the decomposition of wood and forest litter. *Can J Microbiol.* 1958;4,195–204. doi: 10.1139/m58-020
- Bozoudi D, Tsaltas D. The multiple and versatile roles of *Aureobasidium pullulans* in the vitivinicultural sector. *Fermentation.* 2018;4(4):85. doi: 10.3390/fermentation4040085
- Caroff M, Novikov A. Lipopolysaccharides: structure, function and bacterial identifications. *OCL.* 2020;27:31. doi: 10.1051/ocl/2020025
- Chen G, Wang J, Su Y, Zhu Y, Zhang G, Zhao H, et al. Pullulan production from synthetic medium by a new mutant of *Aureobasidium pullulans*. *Prep Biochem Biotechnol.* 2017;47(10):963-969. doi: 10.1080/10826068.2017.1350979
- Chi Z, Liu NN, Jiang H, Wang QQ, Chen JT, Liu GL, et al. Relationship between β -D-fructofuranosidase activity, fructooligosaccharides and pullulan biosynthesis in *Aureobasidium melanogenum* P16. *Int J Biol Macromol.* 2019;125:1103-1111. doi: 10.1016/j.ijbiomac.2018.12.141
- Choudhury AR, Saluja P, Prasad GS. Pullulan production by an osmotolerant *Aureobasidium pullulans* RBF-4A3 isolated from flowers of *Caesulia axillaris*. *Carbohydr Polym.* 2011;83:1547–52. doi: 10.1016/j.CARBPOL.2010.10.003
- Coltelli MB, Danti S, De Clerck K, Lazzeri A, Morganti P. Pullulan for advanced sustainable body- and skin-contact applications. *J. Funct. Biomater.* 2020;11(1):20. doi: 10.3390/jfb11010020.
- Dailin DJ, Selvamani S, Michelle K, Jusoh, YMM, Chuah LF, et al. Production of high-value added exopolysaccharide by biotherapeutic potential *Lactobacillus reuteri* strain. *Biochem Eng J.* 2022;188:108691. doi: 10.1016/j.bej.2022.108691
- Dailin DJ, Low LZMI, Malek RA, Wan Azelee NI, Abdul Manas NH, Keat HC, et al. Pullulan, a biopolymer with potential applications in pharmaceutical and cosmeceutical: A review. *Biosci Res.* 2019;16(3): 2604-2616.
- Duan X, Chi Z, Wang L, Wang X. Influence of different sugars on pullulan production and activities of α -phosphoglucose mutase, UDPG-pyrophosphorylase and glucosyltransferase involved in pullulan synthesis in *Aureobasidium pullulans* Y68. *Carbohydr Polym.* 2008;73(4): 587-593. doi: 10.1016/j.carbpol.2007.12.028
- Elmi A, Nasher F, Dorrell N, Wren B, Gundogdu O. Revisiting *Campylobacter jejuni* virulence and fitness factors: role in sensing, adapting, and competing. *Front Cell Infect Microbiol.* 2021;10, 607704. doi: 10.3389/fcimb.2020.607704
- Haghighatpanah N, Mirzaee H, Khodaiyan F, Kennedy JF, Aghakhan A, Hosseini SS, et al. Optimization and characterization of pullulan produced by a newly identified strain of *Aureobasidium pullulans*. *Int J Biol Macromol.* 2020;152:305-313. doi: 10.1016/j.ijbiomac.2020.02.226.
- Ma ZC, Fu WJ, Liu GL, Wang ZP, Chi ZM. High-level pullulan production by *Aureobasidium pullulans* var. *melanogenum* P16 isolated from mangrove system. *Appl Microbiol Biotechnol.* 2014;98(11):4865-4873. doi: 10.1007/s00253-014-5554-5.
- Maftoun P, Malek R, Abdel-Sadek M, Azi R, Enshasy HE. Bioprocess for semi-industrial production of immunomodulator polysaccharide Pleuran by *Pleurotus ostreatus* in submerged culture. *J Sci Ind Res.* 2013;72:655-662.
- Maldonado RF, S6-Correial, Valvano MA. Lipopolysaccharide modification in Gram-negative bacteria during chronic infection. *FEMS Microbiol Rev.* 2016;40(4):480–493. doi: 10.1093/femsre/fuw007.
- Uzcan E, Sargin S, Guksungur Y. Comparison of pullulan production performances of air-lift and bubble column bioreactors and optimization of process parameters in air-lift bioreactor. *Biochem Eng J.* 2014;92:9-15. doi: 10.1016/j.bej.2014.05.017
- Manzoor A, Dar AH, Pandey VK, Shams R, Khan S, Panesar PS, et al. Recent insights into polysaccharide-based hydrogels and their potential applications in food sector: A review. *Int J Biol Macromol.* 2022;213:987-1006. doi: 10.1016/j.ijbiomac.2022.06.044
- Nordin NZ, Rashidi AR, Dailin DJ, Malek R, Azelee NIW, Manas NH. Xanthan biopolymer in pharmaceutical and cosmeceutical applications: critical review. *Biosci Res,* 2020;17(1): 205-220.

23. Sharma H, Pal J, Neelam DK. Bacterial Extracellular Polymers: A Review. *J Pure Appl Microbiol.* 2021;15(3):1072-1082. doi: 10.22207/JPAM.15.3.28
24. Sheng L, Tong Q, Ma M. Why sucrose is the most suitable substrate for pullulan fermentation by *Aureobasidium pullulans* CGMCC1234? *Enzyme Microb Technol.* 2016;92: 49–55. doi: 10.1016/j.enzmictec.2016.06.016
25. Sheng L, Zhu G, Tong Q. Effect of uracil on pullulan production by *Aureobasidium pullulans* CGMCC1234. *Carbohydr Polym.* 2014;101:435-437. doi: 10.1016/j.carbpol.2013.09.063
26. Chiarello, E., Di Nunzio, M., Picone, G., Antonelli, G., Capozzi, F., & Bordoni, A. (2022). Insight on Glucose and Fructose Absorption and Relevance in the Enterocyte Milieu. *Nutrients*, 14(3), 517.
27. Tzavaras, D., Papadelli, M., & Ntaikou, I. (2022). From milk kefir to water kefir: Assessment of fermentation processes, microbial changes and evaluation of the produced beverages. *Fermentation*, 8(3), 135.
28. Wani, S. M., Masoodi, F. A., Mir, S. A., & Khanday, F. A. (2023). Pullulan production by *Aureobasidium pullulans* MTCC 1991 from apple pomace and its characterization. *Food Bioscience*, 51, 102254.
29. Mohammed, S., & Ray, L. (2022). Polyhydroxyalkanoate recovery from newly screened *Bacillus* sp. LPPI-18 using various methods of extraction from Loktak Lake sediment sample. *Journal of Genetic Engineering and Biotechnology*, 20(1), 1-20.
30. Singh RS, Kaur N, Kennedy JF. Pullulan production from agro-industrial waste and its applications in food industry: A review. *Carbohydr Polym.* 2019;217:46-57. doi: 10.1016/j.carbpol.2019.04.050
31. Umapathi A, Kumawat M, Daima HK. Engineered nanomaterials for biomedical applications and their toxicity: a review. *Environ. Chem. Lett.* 2021;20:445-468. doi: 10.1007/s10311-021-01307-7
32. Yildiz H, Karatas N. Microbial exopolysaccharides: Resources and bioactive properties. *Process Biochem.* 2018;72:41-46. doi: 10.1016/j.procbio.2018.06.009
33. Li, X., Zhao, S., Chen, L., Zhou, Q., Qiu, J., Xin, X., Zhan, Y., Yuan, W., Tian, C., Yang, J., & Yu, X. (2023). High-level production of pullulan from high concentration of glucose by mutagenesis and adaptive laboratory evolution of *Aureobasidium pullulans*. *Carbohydrate Polymers*, 302, 120426.
34. Chi, Z., Kong, C. C., Wang, Z. Z., Wang, Z., Liu, G. L., Hu, Z., & Chi, Z. M. (2022). The signaling pathways involved in metabolic regulation and stress responses of the yeast-like fungi *Aureobasidium* spp. *Biotechnology Advances*, 55, 107898.
35. Yang, J., Li, X., Zhao, S., Yuan, W., Zhou, Q., Zhang, Y., Qiu, J., Wang, J., Zhu, Q., Yang, X., Jiang, X., Tian, C., & Chen, L. (2023). Light calcium carbonate improves pullulan biosynthesis by *Aureobasidium pullulans* under high concentration of sugar. *Food Chemistry*, 415, 135760.