



## Journal of Advanced Research in Fluid Mechanics and Thermal Sciences

Journal homepage:  
[https://semarakilmu.com.my/journals/index.php/fluid\\_mechanics\\_thermal\\_sciences/index](https://semarakilmu.com.my/journals/index.php/fluid_mechanics_thermal_sciences/index)  
ISSN: 2289-7879



# Drying Behavior of Moringa Oleifera Leaves using Tray Dryer

Nabilah Abdul Samad<sup>1</sup>, Yanti Maslina Mohd Jusoh<sup>1</sup>, Nor Alafiza Yunus<sup>1</sup>, Ida Idayu Muhamad<sup>1</sup>, Daniel Joe Dailin<sup>2</sup>, Noorazwani Zainol<sup>2</sup>, Dayang Norulfairuz Abang Zaidel<sup>3,4,\*</sup>

<sup>1</sup> Faculty of Chemical and Energy Engineering, Universiti Teknologi Malaysia, 81310 Johor Bahru, Malaysia

<sup>2</sup> Institute of Bioproduct Development, Universiti Teknologi Malaysia, 81310, Johor Bahru, Malaysia

<sup>3</sup> Department of Chemical and Environmental Engineering, Malaysia-Japan International Institute of Technology, Universiti Teknologi Malaysia, Jalan Sultan Yahya Petra, 54100 Kuala Lumpur, Malaysia

<sup>4</sup> Innovation Centre in Agritechology for Advanced Bioprocessing, Universiti Teknologi Malaysia, Pagoh Campus, 84600 Pagoh, Johor, Malaysia

### ARTICLE INFO

#### Article history:

Received 16 December 2022

Received in revised form 10 April 2023

Accepted 18 April 2023

Available online 7 May 2023

#### Keywords:

Moringa oleifera; drying; tray dryer; color; protein; iron; total phenolic content analysis

### ABSTRACT

Moringa oleifera leaves have been dried in a tray drier to investigate the drying behaviors of Moringa oleifera leaves at three different temperatures for their moisture content, color, iron and protein content, as well as total phenolic content. Three levels of temperatures were chosen varying from 50, 60 and 70 °C in this study. Drying time reduced significantly with the rise in temperature. The final moisture content of dried leaves at 50, 60 and 70 °C was 4.89, 4.79, and 4.59 % and reached equilibrium moisture content at 4, 2, and 1.5 hours. For the water activity, all the dried leaves give a low amount of water activity from 0.0912 to 0.1838 at 70 - 50 °C where the higher the temperature the lower the water activity. This indicates that the lower the temperature, the slower the drying time required to decrease moisture content to below 5 % and a smaller amount of water activity. This ability to preserve its taste and nourishment. For the color changes, the value for greenness (-a) has no significant difference but notable differences were detected between the color parameters of fresh and dried leaves for whiteness L value and yellowness b. This shows that drying under a tray dryer has the same color as the fresh *M. oleifera* leaves. There was no notable difference in the value for protein but a significant difference ( $P < 0.05$ ) in the value for iron contents of the fresh and dried leaves. The total phenolic content (TPC) obtained from this study was 46.97 - 55.78 mg GAE/g FW for drying at 50 to 70 °C. Results from this study discover a great option for large-scale drying using method.

## 1. Introduction

*Moringa oleifera* Lam. is among Moringa species which is the most frequently cited use for therapeutic and nutrition purposes in most countries due to the plant containing an appropriate number of amino acids along with high quantities of essentials nutrients such as carbohydrates, protein, vitamins, calcium, and minerals as well as its ability to withstand in drought resistance. In addition, the plant is known to be a multipurpose tree as all parts of the plant starting from the roots,

\* Corresponding author.

E-mail address: [dnorulfairuz@utm.my](mailto:dnorulfairuz@utm.my)

<https://doi.org/10.37934/arfmts.105.2.182191>

seed, woods, pods, oil, fruits, flower, and leaves have their own benefits and it is also used widely for dealing in malnutrition as it gives cheaper nutrition and contains natural antioxidant properties [1]. Taking 3 oz full of *M. oleifera* leaf powder able to give enough iron and calcium required for pregnant and breastfeeding women [2]. In addition, according to Trigo *et al.*, [3], *M. oleifera* leaf powder is among the greatest natural sources of iron and calcium, as well as a multivitamin supplement with high levels of amino acids and other minerals.

Drying is a vital procedure in removing moisture content from the products by maintaining the aroma of raw herbs, nutritional characteristics, and appearance, as maximum as possible while preserving and extending its storage quality in the quality of flavour and free from microorganisms [4,29]. The first step in the food production is the drying method, especially when considering herbs as the initial step in every food processing industry. Different drying approaches may affect a notable difference in the composition of physicochemical in a dried sample [5]. Plenty of techniques applied by the prior researchers for drying *M. oleifera* leaves such as shade drying, oven drying, sunlight, microwave drying, freeze-drying and tray dryer [6-10].

The tray dryer is a closed insulated compartment in which trays are placed on top of one another in which hot air is supplied by a fan or natural flow [11]. These methods had gained popularity with scholars as a drying process for food, medicine, herbs, chemicals, dye, and material, especially on the large scale, as the dryer is a simple and cost-effective design [12]. Besides, it is also able to establish a great condition of the final sample and is capable to reduce the drying time without affecting the condition of the sample as food sample, since the food is spread out on trays to a proper volume and thickness, it may be dried uniformly as an airflow dispersion across the trays [11].

The prior research on the drying of *M. oleifera* leaves using a tray dryer was completed by Aznury *et al.*, [10] and investigated the drying effect on different temperatures and drying efficiency without taking to account the changes behaviours after drying. The aims of this research were to:

- i. determine the influence of tray dryer at several temperatures (50, 60 and 70 °C) on the moisture and water activity of *M. oleifera* leaves;
- ii. to analyze the differences in the color, iron, and protein contents of the sample after drying; and
- iii. examine total phenolic content for *M. oleifera* leaves by utilizing microwave-assisted extraction (MAE) technique.

## 2. Materials and Methods

### 2.1 Materials

*Moringa Oleifera* Lam. leaves were obtained from Hadham Enterprise, Johor, Malaysia same as the previous study by Samad *et al.*, [9].

### 2.2 Drying of *Moringa Oleifera* Leaves

The drying procedure steps were conducted by referring to previous study by Samad *et al.*, [9]. *M. oleifera* leaves were cleaned with water and then dusted with a towel to remove contaminants from the surface of the leaves and to measure its initial moisture content. The leaves had been dried using a tray-drying method that consist of ten trays (dimension is 41.5cm x 41.5 cm x 3.0 cm for each tray). To obtain *M. oleifera* leaves powder (MOLP), three different temperatures (50, 60, and 70 °C) were investigated. 30 g of fresh leaves sample was spread thoroughly on a square tray to be used for all drying methods with triplicates to report the dry weight until the moisture content (MC) reached less than 5% (Geankoplis, 1993) and then ground using an electric grinder (Waring Commercial

Blender 8011S Model HGB2WTS3, USA) and sieved using a stainless-steel sieve with a diameter of 355  $\mu$ m aperture. The MOLP was then placed in a closed container in a dry and dark cupboard for future usage.

### 2.3 Extraction of *Moringa Oleifera* Using MAE

The extraction procedure was performed via MAE using MAS-II Plus Microwave Synthesis/Extraction Reaction Workstation (SINEO Microwave Chemistry Technology Co. Ltd., China) referring to optimise result done by Sin *et al.*, [13] with slight modification. The 1.5 g powdered *M. oleifera* sample was mixed with 100 mL of 35 % ethanol and placed in a Schott bottle to be soaked and then poured into 4 neck flasks to be extracted in MAE at temperature (65 °C), sample-to-solvent ratio (0.015) and ethanol concentration (35%) for 5 min while the microwave power was fixed at 375 W.

After MAE extraction, the *M. oleifera* leaves extract (MOLE) was set at room temperature to let it cool and centrifuged at the conditions of 15 min, 5000 rpm and 5 °C. Later, the supernatant was dried to evaporate the remaining ethanol solvent by using a rotary evaporator at 60°C in a water bath until all the solvent is evaporated and then dried using a freeze drier (Alpha 1-2-LDplus, Germany). The MOLE obtained were set aside at 5 °C. The MOLE was analysed for TPC.

### 2.4 Colour Analysis

Colour analysis was completed following the procedure from previous study by Ali *et al.*, [8] and been recorded using CIE Standard Illuminant C at CIE L\* a\* b\* color space coordinates

$\Delta E^*_{ab}$  was defined as shown Eq. (1) to signify the amount of the color change between two objects (dried leaves and fresh leaves) as follows in Potisate and Phoungchandang [14]:

$$\Delta E^*_{ab} = \sqrt{(\Delta L^*)^2 + (\Delta a^*)^2 + (\Delta b^*)^2} \quad (1)$$

where L\* is the brightness of a color; a\* is the greenness of a color when negative; b\* is the yellowness of a color when positive.

### 2.5 Protein and Iron Analysis

The protein and iron content were analysed referring to previous study Samad *et al.*, [9] by the association of Official Analytical Chemists (AOAC) method.

### 2.6 Total Phenolic Content (TPC) Analysis

The total phenolic content (TPC) of MOLE was evaluated using a modified Folin-Ciocalteu (FC) method [15]. 600  $\mu$ L of water and 10  $\mu$ L of the sample (50mg in 50ml of 35% ethanol) or standard solution of gallic acid were mixed, to which 50  $\mu$ L of undiluted FC reagent was then added. After 10 min, 150  $\mu$ L of 2 g in 100 mL sodium carbonate solution (Na<sub>2</sub>CO<sub>3</sub>) was added and the volume was filled up to 1 mL with water. Incubate the solution for 2 h at room temperature of 28  $\pm$  1 °C and read absorbance at 760 nm by using EPOCH Microplate Spectrophotometer (BioTek Instrument Inc., USA). The result was expressed as the equivalent of milligrams of gallic acid per gram of Fresh Weight (mg GAE/g FW).

The standard calibration curve was expressed in TPC concentration of the sample equivalent to a milligram of gallic acid per gram (mg GAE/g) with Gallic acid hydrate. Based on the sample concentration, leaves powder dry weight and extract volume were used for calculating TPC yield, according to Eq. (2) by referring to Siddiqui *et al.*, [16]:

$$\text{TPC Yield} = C_1 \times \frac{V_{\text{extract}} (\text{ml})}{m_{\text{plant extract}} (\text{g})} \quad (2)$$

Where  $C_1$  is concentration of gallic acid obtained from the calibration curve in mg/ml,  $V$  is the volume of extract in ml, and  $m$  is the mass of the plant extract (g).

## 2.7 Experimental Design

The drying procedure steps were conducted at three levels of temperatures varying from 50, 60 and 70 °C until the moisture content (MC) reached less than 5%. Then, ground using an electric grinder and sieved to obtain *M. oleifera* leaves powder (MOLP). The MOLP obtained was analysed further for moisture content, water activity, protein and iron content including color analysis.

The extraction procedure was performed via Microwave-assisted Extraction (MAE) at condition 1.5 g powdered *M. oleifera* sample was mixed with 100 mL of 35 % ethanol and placed in a Schott bottle to be soaked and then poured into 4 neck flasks to be extracted in MAE at temperature (65 °C), sample-to-solvent ratio (0.015) and ethanol concentration (35%) for 5 min while the microwave power was fixed at 375 W. After MAE extraction, the *M. oleifera* leaves extract (MOLE) was analysed for TPC.

## 2.8 Data Analysis

The results obtained were documented in average  $\pm$  standard deviation (STDEV) for triplicates and statistically evaluated and performed utilizing Microsoft Excel. A difference of  $P < 0.05$  level was counted as statistically significant. Besides, the t-test was calculated to determine the statistically significant of the result obtained.

## 3. Results and Discussion

### 3.1 Moisture Content & Water Activity

Table 1 tabulates the moisture content (MC), water activity ( $A_w$ ) and drying time for MOLP at temperature limits for 50 – 70 °C. The initial moisture content for the fresh leaves was 80.34 %. Table 1 shows that the drying time increased when the temperature decreased. The greater the temperature, the quicker the drying time required to obtain the moisture content below 5 %. The final MC of dried leaves at 50, 60 and 70 °C were about 4.89, 4.79, and 4.59 % and achieved equilibrium moisture content at 4, 2, and 1.5 hours. For the water activity, all the dried leaves give a low amount of water activity from 0.0912 – 0.1838 at 70 - 50 °C where the higher the temperature the lower the water activity. Hence, it is free from microorganisms like mould, yeast, and bacteria.

**Table 1**

Moisture content and water activity  
after tray dryer method

T (°C)	Final MC (%)	Time (h)	Aw
50	4.89	4	0.1838
60	4.79	2	0.1208
70	4.59	1.5	0.0912

This indicates that the lesser the temperature, the slower the drying time required to obtain the MC to below than 5 %. By decreasing the MC capable to preserve their taste and nourishment along with the increase its shelf life [8,17]. This finding demonstrates that MC was significantly affected by temperature and drying time.


Comparable trend on the properties of drying has been stated by prior studies in using tray dryer for green herbs, for example, *M. oleifera* leaves, mint leaves, white mulberry leaves, curry leaves and green tea leaves [10,18,19,20,21]. Earlier research on the drying of *M. oleifera* leaves using conventional drying technique by Ali *et al.*, [7] showed around 8, 5.75 and 2 hours at 40, 50 and 60°C in an oven dryer which is more slower drying time contrasted to this research. This shows that a tray dryer technique gives a favourable option for drying many *M. oleifera* leaves at a considerably quicker drying time.

### 3.2 Protein and Iron Content

From Figure 1(a) illustrated 7.55 % of fresh MOLP in the protein content was and a bit higher as compared to the amount was reported by Trigo *et al.*, [3] which was 6.7 %. After the drying process, lots of moisture was reduced to less than 5%, and the amount of protein increased to 27.01, 25.92 and 27.08 % at 50, 60 and 70 °C. There was no notable difference ( $P > 0.05$ ) in the protein content of the dried MOLP at different temperatures. This shows that even at high or low temperatures, the amount of protein content would be almost similar. However, in this study found that, at low temperatures (50 °C) it takes four times longer than compared to dry at 70 °C. But protein is very sensitive to high temperatures as denaturation of protein cells might influence it at high temperatures. This can be supported by a study from Razzak *et al.*, [22] where the protein content decreased from 24.67 to 19.89 % at 60 – 80 °C using an oven dryer.

The protein content in this study was acceptable with the finding by Ali *et al.*, [8] where the amount of protein obtained was 29.8, 28.8 and 28.6 % at 40, 50 and 60 °C respectively using oven drying method while a study by Samad *et al.*, [9] obtained from 28.15 – 28.75 % at 300 - 1000 W using microwave drying method. The difference in protein content may be because of the diverse locality and moment of harvested [6]. This amount of protein content indicated a great advantage in therapeutic and nutrition purposes because of a good amount of nutrient preservation.

According to Figure 1(b) the amount of fresh MOLP was 14.46 mg/kg similar to previous study by Samad *et al.*, [9] and after the drying the amount of iron improved to 44.41, 47.54 and 53.54 mg/kg at 50, 60 and 70 °C due to moisture change. There is a substantial difference ( $P < 0.05$ ) between 60 and 70 °C. A higher amount of iron for fresh leaves was reported by a previous study by Ali *et al.*, [8] at 18.5 mg/kg and the result for oven drying at 40, 50 and 60 °C were 207.7, 216.6 and 209.6 mg/kg. The differences could be regarding of locality of the plantation of *M. oleifera* and the time been harvested [6].



**Fig. 1.** Content for tray drying of MOLP at 50, 60 and 70 °C (a) Protein, (b) Iron

The improved amount of protein and iron after drying were caused by the variation in nutrient concentration of internal composition within the *M. oleifera* leaves as the moisture was decreased to less than 5% [23]. This implies that finding the best drying method for *M. oleifera* leaves which have a great amount of iron and protein content after drying is necessary as it can improve the nutrition amount as well as can maintain its advantageous qualities [24].

### 3.3 Color Measurement

Appearance of a dried leaves is an important marker in terms of quality part to get a customer approval for the end-product. The tabulate data in Table 2 shows the color analysis of fresh and dried MOLP for three temperatures (50, 60 and 70 °C). The original values of L\*, a\* and b\* for fresh *M. oleifera* leaves was 41.25, -3.95 and 3.33. The brightness values L\* decreased a little as the drying temperature rise from 50 to 70 °C. There was a notable difference to the changes in L\* and b\* values as the drying temperature increases. This small difference in L\* may resulted in the leaves appearance to turn into dark green. Similar effect has been stated by Rudra *et al.*, [25] where the L\* value declined after drying at all drying temperatures during oven drying. Chlorophyll degradation is one of the causes that the changes in brightness value of dark green in *M. oleifera* leaves occur [7].

**Table 2**

Effect of different tray dryer temperature on the color of *M. oleifera* leaves; L\* is the brightness of a color; a\* is the greenness of a color when negative; b\* is the yellowness of a color when positive

Drying Condition	FRESH	50 (°C)	60 (°C)	70 (°C)
L*	49.93 ± 1.32 <sup>a</sup>	47.80 ± 1.69 <sup>b</sup>	45.44 ± 1.22 <sup>a</sup>	46.44 ± 1.44 <sup>b</sup>
a*	-3.95 ± 0.25 <sup>a</sup>	-2.78 ± 0.08 <sup>b</sup>	-2.38 ± 0.31 <sup>b</sup>	-2.016 ± 0.60 <sup>b</sup>
b*	3.33 ± 0.18 <sup>a</sup>	2.77 ± 0.28 <sup>b</sup>	1.82 ± 0.20 <sup>b</sup>	1.92 ± 0.10 <sup>a</sup>
ΔE*ab		2.4933	4.9876	4.2317

In this study, all the drying process stop at MC less than 5% hence, study by Chen *et al.*, [30] support that at lower moisture the color of the sample is similar to fresh sample by referring to the amount of ΔE\*. However, from this study it found that the ΔE\*ab obtained at 60 °C has the highest color change matched to the fresh leaves while 50 °C has the smallest which was a 2.49 color difference. This shows that the changes in color values were minimum in lower temperature as the

drying received lower heat thermal. Furthermore, study by Ali *et al.*, [8] confirm this by stating that low  $\Delta E^*ab$  levels are preferable in dry food items when compared to the fresh leaves.


From this research, there are slight variations in the color as darkening occurred, although the color nearly to the initial fresh *M. oleifera* leaves as there is just a little reduction in the value of color parameter by using tray drying technique. This shows that the tray drying technique at 50 °C able to sustain a perfect and acceptable green color as the original leaves.

### 3.4 Total Phenolic Content

The study by Rahim *et al.*, [26] stated that antioxidants that originated from plants have been known as free radical scavengers. Moreover, researchers found that some medicinal plants consist of multiple classes of natural antioxidants, such as phenolic acids, flavonoids, and tannins, which have more strong antioxidant activity than dietary plants [27].

MOLE contain abundant phenolic compounds such as quercetin and kaempferol which belongs to the flavonoid group [14]. This is due to the free radical scavenging capability being enabled by the hydroxyl groups; therefore, the TPC could be used as a basis for rapid screening of antioxidant activity. The TPC of the MOLE was measured based on the standard calibration curve of  $y = 0.3406x - 0.003$ ,  $R^2 = 0.9948$ . From Figure 2, the range of TPC obtained from this study was from 51.87 – 54.81 mg GAE/g FW for drying at 50 - 70 °C. There is no significant difference in temperatures for drying MOLE using a tray dryer. The amount of TPC obtained in this study is lower with the study reported by Abd Hashib *et al.*, [31] for drying pineapple using spray dried was 58 mg GAE/100g fruits compared with fresh pineapple.

The decreased value for TPC was due to the thermal processing of different temperature as the bounding between phytochemical were released from the matrix. Hence, it causes a synergistic effect with other phytochemicals, for instance, phenolics and flavonoids might also affect. Besides, study shows that MOLE have a strong source of cytokinin which can distinguish signals and gives a lot of phenolics and other plant secondary metabolites [28]. However, the previous studies by Potisate and Phoungchandang [14] were found to be lower which the study from 5 varieties at the range of 7 – 33 mg GAE/g FW. The differences could be regarding of locality of the plantation of *M. oleifera* and the time been harvested [6]. Hence, it can conclude that MOLE are rich in total phenolic contents and can act as natural antioxidants.



**Fig. 2.** The total phenolic content for tray drying of MOLE at 50, 60 and 70 °C

#### 4. Conclusions

Drying using tray dryer method was studied at three different temperatures varying from 50, 60 and 70 °C. The drying time decreased significantly with the increase in drying temperature. The final MC of dried MOLP at 50, 60 and 70 °C were 4.89, 4.79, and 4.59 % and reached equilibrium MC at 4, 2, and 1.5 hours. The water activity obtained was from 0.0912 – 0.1838 at 70 - 50 °C. This result indicates that, the lower the temperature, the slower the drying time required to decrease moisture content to below 5 % and a lower amount of water activity. By lowering the moisture content enable to preserve their taste and nourishment. For the color changes, the value for greenness (-a) has no significant difference but significance differences were detected between the color parameters of fresh and dried leaves for brightness (L) value and yellowness (b). This shows that drying using a tray dryer has the same color as the fresh *M. oleifera* leaves. The protein contents for MOLP obtained were 27.01, 25.92 and 27.08 % while the iron contents were 44.41, 47.54 and 53.54 mg/kg at 50, 60 and 70 °C respectively. The TPC of MOLE obtained from this study was 46.97 - 55.78 mg GAE/g FW for drying at 50 - 70 °C. Results from this study discover a great option for large-scale drying using method tray dryer on *M. oleifera* leaves as a fast-drying technique.

#### Acknowledgement

The authors acknowledged Malaysia-Japan International Institute of Technology, Universiti Teknologi Malaysia Kuala Lumpur for the funding and Ministry of Education Malaysia, Universiti Teknologi Malaysia for the financial support from Research Grant Q.K130000.3843.21H95. Continuous encouragement and laboratories support during the work by the management and laboratory technician of Faculty of Chemical and Energy Engineering.

#### References

- [1] Feihrmann, Andresa C., M. G. Nascimento, C. Z. Belluco, P. Fioroto, L. Cardozo-Filho, and L. C. Tonon. "Evaluation of the Effect of Antioxidant Moringa Oleifera." *Chemical Engineering Transactions* 57 (2017): 1993-1998.
- [2] Gopalakrishnan, Lakshmipriya, Kruthi Doriya, and Devarai Santhosh Kumar. "Moringa oleifera: A review on nutritive importance and its medicinal application." *Food science and human wellness* 5, no. 2 (2016): 49-56. <https://doi.org/10.1016/j.fshw.2016.04.001>
- [3] Trigo, Carla, Maria Luisa Castello, Maria Dolores Ortola, Francisco Jose Garcia-Mares, and María Desamparados Soriano. "Moringa oleifera: An unknown crop in developed countries with great potential for industry and adapted to climate change." *Foods* 10, no. 1 (2020): 31. <https://doi.org/10.3390/foods10010031>
- [4] Elzaawely, A. A., and S. Tawata. "Effect of extraction and drying methods on the contents of kava pyrones and phenolic compounds in *Alpinia zerumbet* leaves." *Asian Journal of Plant Sciences* 10, no. 8 (2011): 414-418. <https://doi.org/10.3923/ajps.2011.414.418>
- [5] Capecka, Ewa, Anna Mareczek, and Maria Leja. "Antioxidant activity of fresh and dry herbs of some Lamiaceae species." *Food chemistry* 93, no. 2 (2005): 223-226. <https://doi.org/10.1016/j.foodchem.2004.09.020>
- [6] Anjorin, Toba Samuel, Pius Ikokoh, and Simon Okolo. "Mineral composition of *Moringa oleifera* leaves, pods and seeds from two regions in Abuja, Nigeria." *International Journal of Agriculture and Biology* 12, no. 3 (2010): 431-434.
- [7] Ali, M. A., Y. A. Yusof, N. L. Chin, M. N. Ibrahim, and S. M. A. Basra. "Drying kinetics and color analysis of *Moringa oleifera* leaves." *Agriculture and Agricultural Science Procedia* 2 (2014): 394-400. <https://doi.org/10.1016/j.aaspro.2014.11.055>
- [8] Ali, M. A., Y. A. Yusof, N. L. Chin, and M. N. Ibrahim. "Processing of *Moringa* leaves as natural source of nutrients by optimization of drying and grinding mechanism." *Journal of Food Process Engineering* 40, no. 6 (2017): e12583. <https://doi.org/10.1111/jfpe.12583>
- [9] Samad, Nabilah Abdul, Dayang Norulfairuz Abang Zaidel, Ida Idayu Muhamad, Yanti Maslina Mohd Jusoh, and Nor Alafiza Yunus. "Influence of microwave drying on the properties of *Moringa oleifera* leaves." *Chemical Engineering Transactions* 89 (2021): 469-474.



- [10] Aznury, Martha, Ahmad Zikri, Nova Rachmadona, and Muhammad Farhan Saputra. "Analyzing Effect of Temperature on Drying Moringa (*Moringa Oleifera*) Leaves Using Photovoltaic Tray Dryer." *Journal of Hunan University Natural Sciences* 48, no. 10 (2021).
- [11] Omofoyewa, Mufliah Gbemisola, Lekan Taofeek Popoola, and Abdulwahab Giwa. "Optimization of drying parameters of moringa seeds in a tray dryer using Box-behnken technique of RSM." In *International Journal of Engineering Research in Africa*, vol. 32, pp. 86-99. Trans Tech Publications Ltd, 2017. <https://doi.org/10.4028/www.scientific.net/JERA.32.86>
- [12] Misha, Suhaimi, S. Mat, M. H. Ruslan, Kamaruzzaman Sopian, and Elias Salleh. "Review on the application of a tray dryer system for agricultural products." *World applied sciences journal* 22, no. 3 (2013): 424-433.
- [13] Sin, Kang, Wilheliza A. Baraoidan, and Pag-asa D. Gaspillo. "Microwave-assisted extraction of phenolic compounds from *Moringa oleifera* Lam. leaves using response surface methodology as optimization tool." *Philipp. Agric. Sci* 97, no. 1 (2014): 36-42.
- [14] Potisate, Yuparat, and Singhanat Phoungchandang. "Microwave drying of *Moringa oleifera* (Lam.) leaves: drying characteristics and quality aspects." *Asia-Pacific Journal of Science and Technology* 20, no. 1 (2015): 12-25.
- [15] Rodríguez-Pérez, Celia, Bienvenida Gilbert-López, Jose Antonio Mendiola, Rosa Quirantes-Piné, Antonio Segura-Carretero, and Elena Ibáñez. "Optimization of microwave-assisted extraction and pressurized liquid extraction of phenolic compounds from *Moringa oleifera* leaves by multiresponse surface methodology." *Electrophoresis* 37, no. 13 (2016): 1938-1946. <https://doi.org/10.1002/elps.201600071>
- [16] Siddiqui, Nazish, Abdur Rauf, Abdul Latif, and Zeenat Mahmood. "Spectrophotometric determination of the total phenolic content, spectral and fluorescence study of the herbal Unani drug Gul-e-Zoofa (*Nepeta bracteata* Benth)." *Journal of Taibah university medical sciences* 12, no. 4 (2017): 360-363. <https://doi.org/10.1016/j.jtumed.2016.11.006>
- [17] Geankoplis, Christie J. "Transport Processes and Unit Operations: PTR Prentice Hall." *City, State, Country* (1993).
- [18] Raviteja, G., P. S. Champawat, S. K. Jain, and S. Chavan. "Drying characteristics of mint leaves in tray dryer." *International Journal of Current Microbiology and Applied Sciences* 8, no. 3 (2019): 543-551. <https://doi.org/10.20546/ijcmas.2019.803.066>
- [19] Phoungchandang, S., L. Tochip, and V. Srijesdaruk. "White mulberry leaf drying by tray and heat pump dehumidified dryers." *World J Agric Sci* 4, no. 5 (2008): 844-851.
- [20] Jain, Priti, Nidhi Khanna, M. K. Pathak, and Rajendra Singh Thakur. "Drying Kinetics of Curry Leaves under Thin Layer Tray Drying" *International Journal of Innovative Research in Science, Engineering and Technology* 6, no. 11 (2017): 21729–21733.
- [21] Shomali, Abbas, and Behrooz Abbasi Souraki. "Experimental investigation and mathematical modeling of drying of green tea leaves in a multi-tray cabinet dryer." *Heat and Mass Transfer* 55 (2019): 3645-3659. <https://doi.org/10.1007/s00231-019-02662-6>
- [22] Razzak, A., Roy, K. R., Sadia, U., Mominul, H., Suvro, T., Sikder, M. B. H., Zzaman, W. "Effect of drying condition on physicochemical and antioxidant properties of dried *Moringa* leaf powder." *Food Research* 5, no 6 (2021): 165–171. [https://doi.org/10.26656/fr.2017.5\(6\).759](https://doi.org/10.26656/fr.2017.5(6).759)
- [23] Pati, G. D., I. L. Pardeshi, and K. J. Shinde. "Drying of green leafy vegetables using microwave oven dryer." *Journal Ready to Eat Food* 2, no. 1 (2015): 18-26.
- [24] Yashaswini, J. P., N. Venkatachalapathy, Kulbhushan Sharma, R. Jaganmohan, and Akash Pare. "Effect of microwave vacuum drying on nutritional composition of moringa (*Moringa oleifera*) leaves." *International Journal of Chemical Studies* 9, no. 1 (2021): 2790-2795. <https://doi.org/10.22271/chemi.2021.v9.i1am.11648>
- [25] Rudra, Shalini Gaur, Harjinder Singh, Santanu Basu, and U. S. Shivhare. "Enthalpy entropy compensation during thermal degradation of chlorophyll in mint and coriander puree." *Journal of Food Engineering* 86, no. 3 (2008): 379-387. <https://doi.org/10.1016/j.jfoodeng.2007.10.020>
- [26] Rahim, MS Azman Abdul, Jailani Salihon, Mashitah Mohd Yusoff, and M. Rizal Martua Damanik. "Antioxidative Activity and Phenols Content in Five Tropical Lamiaceae Plants." *Journal of Tropical Resources and Sustainable Science (JTRSS)* 1, no. 2 (2013): 49-54. <https://doi.org/10.47253/jtrss.v1i2.641>
- [27] Wong, Chi-Chun, Hua-Bin Li, Ka-Wing Cheng, and Feng Chen. "A systematic survey of antioxidant activity of 30 Chinese medicinal plants using the ferric reducing antioxidant power assay." *Food chemistry* 97, no. 4 (2006): 705-711. <https://doi.org/10.1016/j.foodchem.2005.05.049>
- [28] Farooq, B., and B. Koul. "Comparative analysis of the antioxidant, antibacterial and plant growth promoting potential of five Indian varieties of *Moringa oleifera* L." *South African Journal of Botany* 129 (2020): 47-55. <https://doi.org/10.1016/j.sajb.2018.12.014>
- [29] Manuwa, S. I., A. M. Sedara, and F. A. Tola. "Design, fabrication and performance evaluation of moringa (*oleifera*) dried leaves pulverizer." *Journal of Agriculture and Food Research* 2 (2020): 100034. <https://doi.org/10.1016/j.jafr.2020.100034>

- [30] Chen, Bo-Lin, Guan-Syun Lin, and Wei-Mon Yan. "Effect of Different Drying Conditions on Vacuum Freezing and Microwave-Combined Drying of Pineapple. " *Journal of Advanced Research in Experimental Fluid Mechanics and Heat Transfer* 8, no 1 (2023): 1–5.
- [31] Abd Hashib, Syafiza, Umami Kalthum Ibrahim, Asdarina Yahya, and Norazah Abd Rahman. "The Comparison of Bioactive Compounds and Antioxidant Activity of Fresh Pineapple and Pineapple Powder." *Journal of Advanced Research in Applied Sciences and Engineering Technology* 17, no. 1 (2019): 54-60.