

ICCCP 2012: 5-6 May 2012, Kuala Lumpur, Malaysia

Review on the Extraction of Biomolecules by Biosurfactant Reverse Micelles

S. H. Mohd-Setapar^{a,*}, S. N. Mohamad-Aziz^a, N. H. Harun^a, C. Y. Mohd-Azizi^a

Center of Lipid Engineering Applied Research (CLEAR), Faculty of Chemical Engineering, Universiti Teknologi Malaysia, 81310 UTM Skudai, Johor, Malaysia

Abstract

Surfactant reverse micelles have been widely applied in biotechnology areas for decades. Due to its unique criteria in protecting the bioactivities of biomolecules, the attempts have typically been done for recovery, separation and purification of many biological products such as proteins, enzymes, and antibiotics. Surfactant is a key role in stabilizing the transfer process into reverse micelles phase. Its application in reverse micelles extraction has so far been confined to chemical surfactant which synthetically derived from petroleum. By replacing synthetic surfactant with natural biosurfactant in reverse micelles, it may contribute generating products into a green application as well as helping the future to diminish the side effect of using a toxic surfactant to the environment. Biosurfactant which synthesized by microorganism will be a potential candidate as an environmental-friendly stabilizer due to its environmental acceptability. Production of bioactive molecules will be greener as well as meets the worldwide biotechnology needs.

© 2012 Published by Elsevier B.V. Selection and/or peer review under responsibility of Asia-Pacific Chemical, Biological & Environmental Engineering Society Open access under [CC BY-NC-ND license](https://creativecommons.org/licenses/by-nc-nd/4.0/).

Keywords: Extraction, Mixing, Optimization, Affinity, Reverse Micelles, Antibiotics

1. Introduction

The biotechnological process has grown since early 1980s and becoming more important nowadays because it offers numerous advantages over to chemical synthesis. In separating the biological products, the constraint is the maintenance of the producing environments which are susceptible to chemical and physical

* Corresponding author. Tel.: +607-5535496; fax: +607-5581463.

E-mail address: sitihamidah@cheme.utm.my

changes with respect to time. These diluted complex molecules and multicomponent mixtures are also prone to be sensitive to temperature, pH, and concentration throughout the process [1]. Liquid-liquid extraction is a versatile technology and very useful in the separation of compounds such as biomolecules, chemicals and nanoparticles. The process transfer is motivated by the differences in polarity and hydrophobic or hydrophilic character of the feed components [2]. Liquid-liquid extraction by reverse micelles has successfully been demonstrated to be capable of selectively solubilise the biomolecules such as proteins [3], amino acids [4] and enzymes [5]. It greatly received attention in recent years due to its ability to solubilise the biomolecules while retaining their bioactivities and native structures in the organic environment [6]. The distribution of biomolecules between the organic media and aqueous phase in reverse micelles system is largely driven by the operational parameters such as water content, aqueous pH, ionic strength, type of salt, type of solvent, concentration and type of surfactant [7]. Surfactant applications in reverse micelles extraction have so far been confined to chemical surfactants which synthetically derived from petroleum, thus may causing the environmentally and toxically problems. Therefore, biosurfactant which synthesized by variety of microorganism will be a potential candidate as environmental-friendly development in modern bioseparation technology. In this short review, we discuss the potential applications of biosurfactant in bioseparation as an alternative to the synthetic surfactant. With its ability as stabilizer in reverse micelles system, we can look forward to biosurfactant as green molecules of the future.

2. Biomolecules Separation

The typical production of biomolecules mainly involves two main steps, i.e. fermentation section, and subsequently the downstream process. The manufacturing of biomolecules initiated with a highly specialized fermentation process containing very dilute and complex mixtures which end up with low products concentration [8]. Then, the products quality depends strongly throughout the downstream processing which comprises a large, energy-intensive separation and purification steps.

2.1. Reverse Micelles System

Reverse micelles consist of spheroidal aggregates of surfactant molecules which spontaneously formed in the organic solvent media [9]. As shown in Fig. 1, the system is forming when the polar head groups of surfactant molecules are attracted by aqueous core and oriented towards inside, and a non-polar part of hydrocarbon chain is attracted by organic phase and point towards outside [10]. This is due to the aggregates generally constituted in ternary surfactant-water-oil mixtures which can be referred as a phase region of Winsor II system [11]. The inner core of reverse micelles essentially is a nano-sized droplet of water act as a host of solubilised molecules. Water and hydrophilic molecules such as biological active compounds can be solubilised inside the cores, which stabilized by surfactant layer shielding the water phase from the oil continuum [9].




Fig. 1. A typical structure of reverse micelle

2.2. Reverse Micelles Properties and Benefits

The reverse micelles extraction has a great potential to be implemented in the recovery, separation and purification of the targeted biomolecules due to it can be implemented only under moderate conditions. The main advantage is the selective solubilisation of biomolecules inside reversed micelles which provides a logical aqueous environment protecting the bioactivities of the biomolecules [12]. Reverse micelles have successfully been carried out by many researches due to its system has a lower viscosity and shorter phase separation time [13]. Furthermore, reverse micelles are thermodynamically stable, optically transparent, low costs, energy savings, and can easily be operated continuously at steady state, which are beneficial both for large-scale production and for quality control [13].

2.3. Factors Affecting Biomolecules Transfer

The partitioning of biomolecules between the micellar phase and an aqueous phase is dependent upon the environments of both bulk aqueous phase and organic phase. Factors that affect the phase transfer include water content, pH values, ionic strength, temperature, surfactant type and concentration, and type of solvent [14]. The fundamental studies of the factors concluding selective separation of biomolecules are essential to establish correlations between physicochemical properties of the biomolecules and shows below:

1) Water content is described by the water to the surfactant molar ratio ($[H_2O]/[surfactant]$) renowned as w_o or water in oil [15]. This parameter indicates the structure and size of the micelles, and the number of surfactant molecules per micelle with the assumption of that all the surfactant is located in the micellar wall. Reverse micelles is formed when the surfactant aggregates contain a small amount of water, i.e. once w_o is below 15 [10].

2) The water phase pH of the solution determines the net charge of biomolecules. The pH must be at a level that generates a biomolecule net charge opposite in sign to the surfactant headgroup, so that there is an attraction between the biomolecule surface and the polar headgroups on the internal surface of the reversed micelle. Studies on the pH value effect on protein extraction by the reversed micelle process indicate that the difference between the pH and the isoelectric point (pI) of the protein must oscillate between 1 and 2 points for higher extraction efficiency [16].

3) In reverse micelles structures, repulsion occurs between the surfactant molecules which caused by the surfactant charge and ions of the opposite charge present in the micellar water phase. These ions control the repulsion force of the surfactant headgroups and consequently the micellar sizes. Increases in the ionic strength of the biomolecules feed solution can be expected to reduce the interaction between the biomolecule and surfactant head groups, hence decreasing the biomolecules solubilization. This is due the screening effect, that is, the formation of an electrostatic shield that reduces the intensity of the electrostatic interaction between biomolecule and surfactant as explained by Andrews [17].

4) Changes in temperature have drastic effects on the physicochemical properties of the reverse micellar system. Marcozzi [18] observed a significant increase in α -chymotrypsin recovery when carrying out backward extraction at 38°C, and when Glatz [19] conducted the backward extraction of glucoamylase at 35°C, the recovery of enzymatic activity increased from 40% to 90%.

5) Solvent has known to influence the biomolecules transfer between aqueous phase and organic phase. The solvents that commonly be used in reverse micelles are organic and immiscible in water [15, 20]. They included isooctane, n-octane, heptanes, cyclohexane, benzene, kerosene, and chloroform. However, Ruth and his researchers [21] reported the possibility of creating the reverse micellar environment with addition of

several different polar but does not support for large micellar size.

6) Surfactant type and concentration is the most important variables for biomolecules solubilisation by reverse micelle. Biomolecule solubilisation in the organic phase is increased by increasing the surfactant concentration [22]. Choosing an appropriate surfactant concentration in the organic phase can change the selectivity of the reverse micellar phase for a targeted biomolecule [23].

3. Biosurfactant Potential Applications

Biosurfactant is a natural microbial surfactant which abundantly produced from natural resources by variety of microorganism [24]. It is a surface-active amphiphatic molecule of biological origin which omnipresent in biological systems and possess similar mechanisms as synthetic surfactant. Biosurfactant contains hydrophobic and hydrophilic moieties that grant the ability to accumulate between fluid phases, thus reducing surface and interfacial tension at the surface and respectively [25]. Biosurfactant have gained considerable interest recently in a wide range of industrial applications, especially in food and pharmaceutical sector. The reason for their popularity as high value microbial products is primarily because of their specific action, lower toxicity, higher biodegradability, stable at extremes of temperature, pH, salinity and widespread applicability, and their unique structures which provide new properties that classical surfactants may lack [26].

As proclaimed by Goto [27], it is important to insure the chosen biosurfactant has no negative impact on the stability or activity of the product since biosurfactant may bind to proteins and other bioactive molecules. Thus, from the classification properties, the most known biosurfactant found out are glycolipids, which consist of long-chain aliphatic acids or hydroxyaliphatic acids. Among the glycolipids, the best known are rhamnolipids, trehalolipids and sophorolipids [28]. For instance, sophorolipids (SLs) is one of the most important members of glycolipids biosurfactant is produced by yeast and derived from plant. SLs consist of a dimeric carbohydrate sophorose linked to a long-chain hydroxy fatty acid, and have a mixture of acidic and lactonic forms with at least 6 to 9 different hydrophobic sophorosides. Schippers [29] revealed that CMC and the solubilization ratio of the SLs were found to be in a good range compared to synthetic surfactant.

4. Conclusion

Since the toxicity of chemically synthesized surfactant may causing the serious health and environmental problems, replacement by alternative biosurfactant in reverse micelles extraction is a green movement in biotechnology application. Sophorolipids is the potential candidate of biosurfactant which produced by yeast, i.e. non-pathogenic molecule. With good range of critical micelle concentration and solubilization ratio, it should be further emphasized for biomolecules extraction in downstream processing. Consequently, this will contribute to diminish the side effects of releasing toxically chemical surfactant to the environment as well as promoting and commercializing the development of new green technology in bioseparation industry to the country and worldwide.

Acknowledgements

The authors gratefully appreciate the support of the Ministry of Science, Technology & Innovation, Malaysia (MOSTI) for the National Science Foundation scholarship (NSF) awarded to Siti Norazimah Bt Mohamad Aziz. The authors also acknowledged for financial support from Ministry of Higher Education (MOHE) and University Teknologi Malaysia (Vote No: Q.J130000.7125.00H02)

References

- [1] Wethearly LR. *Engineering processes for bioseparations*. Butterworth-Heinemann Ltd; 1994.
- [2] Frank TC, Dahuron L, Holden BS, Prince WD, Seibert AF, Wilson LC. *Liquid-liquid Extraction and other liquid-liquid operations and equipment, Perry's Chemical Engineers' Handbook*. McGraw-Hill; 2008.
- [3] Shen CW, Yu T. Protein separation and enrichment by counter-current chromatography using reverse micelle solvent systems. *J. Chromatography A* 2007;**1151**:164-168.
- [4] Dövyap Z, Bayraktar E, Mehmetoğlu, V. Amino acid extraction and mass transfer rate in the reverse micelle system. *Enz Microb Tech* 2006;**38**:557-562.
- [5] Debnath S, Das D, Das PK. Unsaturation at the surfactant head: influence on the activity of lipase and horseradish peroxidase in reverse micelles. *Biochem Biophys Res Com* 2007;**356**:163-168.
- [6] Yu YC, Chu Y, Ji JY. Study of the factors affecting the forward and back extraction of yeast-lipase and its activity by reverse micelles. *J Colloid Interf Sci* 2003;**267**:60-64.
- [7] Zhao X, Li Y, He X, Zhong N, Xu Z, Yang L. Study of the factors affecting the extraction of soybean protein by reverse micelles. *Mol Biol Rep* 2009;**37**:669-675.
- [8] Naoe K, Noda K, Kawagoe M, Imai M. Higher order structure of proteins solubilized in AOT reverse micelles. *Coll Surf B* 2004; **38**:179-185.
- [9] Dekker M, Leser ME. *The use of reverse micelles for the separation of proteins, Highly Selective Separations in Biotechnology*. 1st Ed. Chapman & Hall. 1994.
- [10] Malik MA, Mohammad YW, Mohd AH. Microemulsion method: A novel route to synthesize organic and inorganic nanomaterials. *Arab J of Chemistry* 2010;1-21.
- [11] Paul BK, Moulik MSP. Uses and applications of microemulsions. *Current Science* 2001;**80**(8):990-1001.
- [12] Tingyue G. *Liquid-liquid Partitioning methods for bioseparations, handbook of Bioseparations*. 2nd ed. New York: Academic Press; 2000.
- [13] Priscilla GM, Andre ML, Francilene AH, Angela FJ, Thereza CVP, Perola OM, Carlota OR, Adalberto PJ. Liquid-liquid extraction of biomolecules: an overview and update of the main techniques. *J of Chem Technol & Biotechnol* 2008;**83**:143-157.
- [14] Rodrigues EMG, Pessoa Jr A, Milagres AMF. Screening of variables in xylanase recovery using bdbac reversed micelles. *Appl Bioche. Biotechnol* 1999a;**779**-788.
- [15] Luisi PL, Giomini M, Pileni MP, Robinson BH. Reversed micelles as hosts for proteins and small molecules. *Biochim Biophys Acta* 1988; **47**:209-246.
- [16] Pessoa Junior A, Vitolo M. Separation of inulinase from *Kluyveromyces marxianus* using reversed micellar extraction. *Biotechnol Techn* 1997;**11**(6):421-422.
- [17] BA Andrews, Haywood K. Effect of pH, ion type and ionic strength on partitioning of proteins in reverse micelle systems. *J of Chrom A* 1994;**668**:55-60.
- [18] Marcozzi G, Correa N, Luisi PL, Caselli M. Protein extraction by reversed micelles: a study of the factors affecting the forward and backward transfer of α -Chymotrypsin and its activity. *Biotechnol Bioeng* 1991;**38**:1239-1246.
- [19] Glatz CE. Extraction of charged fusion proteins in reversed micelles: comparison between different surfactant systems. *Biotechnol Prog* 1995;**11**:260-264.
- [20] QL Chang, JY Chen. Reversed micellar extraction of trypsin:effect of solvent on the protein transfer and activity recovery. *Biotechnol Bioeng* 1995b;**46**:172-174.
- [21] Ruth ER, Joel RK, Erik PU, Nancy EL. Novel Reverse micelles partitioning nonaqueous polar solvents in a hydrocarbon continuous phase. *J Phys Chem B* 1997;**101**:8292-8297.
- [22] Hentsch M, Menoud P, Steiner L, Flaschel E, Renken A. Optimization of the surfactant (AOT) concentration in a reverse micellar extraction process. *Biotechnology Techniques* 1992;**6**(4):359-364.
- [23] Pires MJ, Aires-Barros MR, Cabral JMS. Liquid-liquid extraction of proteins with reversed micelles. *Biotechnol Prog* 1996; **12**:290-301.

- [24] Kitamoto D, Isoda H, Nakahara T. Functions and potential applications of glycolipid biosurfactants from energy-saving materials to gene delivery carriers. *J Biosci Bioeng* 2002;**94**:187-201.
- [25] Karanth NGK, Deo PG, Veenanadig NK. Microbial production of biosurfactants and their importance. *Curr Sci* 1999; **77**:116–123.
- [26] Desai JD, Banat IM. Microbial production of surfactants and their commercial potential. *Microbiol Mol Biol Rev* 1997;**61**:47–64.
- [27] Goto M, Ono T, Nakashio F, Hatton TA. Design of suitable for protein extraction by reversed micelles. John Wiley & Sons, Inc 1997:26-32.
- [28] Muthusamy K, Gopalakrishnan S, Ravi TK, Sivachidambaran P. Biosurfactants: Properties, commercial production and application. *Current Science* **94**(6): 736-747.
- [29] Schippers C, Gessner K, Müller T. Microbial degradation of phenanthrene by addition of a sophorolipid mixture. *J Biotechnol* 2000; **83**:189-198.