

INTEGRAL EQUATION APPROACH FOR COMPUTING GREEN'S  
FUNCTION ON UNBOUNDED SIMPLY CONNECTED REGION

SHEIDA CHAHKANDI NEZHAD

A dissertation submitted in partial fulfillment of the  
requirements for the award of the degree of  
Master of Science (Mathematics)

Faculty of Science  
Universiti Teknologi Malaysia

JUNE 2013

## ABSTRACT

This research is to compute the Green's function on an unbounded simply connected region by conformal mapping and by solving an exterior Dirichlet problem. The exact Green's function is found by using Riemann mapping and Möbius transform. The Dirichlet problem is then solved using a uniquely solvable Fredholm integral equation on the boundary of the region. The kernel of this integral equation is the generalized Neumann kernel. The method for solving this integral equation is by using the Nyström method with the trapezoidal rule to discretize it into a system. The linear system is solved by the Gaussian elimination method. As an examination of the proposed method, several numerical examples for some various test regions are presented. These examples include a comparison between the numerical result and the exact solutions.

## ABSTRAK

Kajian ini adalah untuk mengira fungsi Green pada rantau terkait ringkas tak terbatas dengan kaedah pemetaan konformal dan dengan kaedah menyelesaikan masalah Dirichlet luaran. Fungsi tepat Green boleh ditemui dengan menggunakan pemetaan Riemann dan penjelmaan Möbius. Masalah Dirichlet kemudiannya diselesaikan menggunakan persamaan kamiran Fredholm yang mempunyai penyelesaian unik di sempadan rantau ini. Inti persamaan kamiran ini adalah inti Neumann teritlak. Kaedah untuk menyelesaikan persamaan kamiran ini ialah dengan menggunakan kaedah Nyström dengan petua trapezoid untuk diskritkannya kepada sebuah sistem. Sistem linear diselesaikan dengan kaedah penghapusan Gauss. Untuk mengkaji kaedah yang dicadangkan, contoh-contoh berangka bagi beberapa rantau ujian dibentangkan. Contoh-contoh ini termasuk perbandingan antara keputusan berangka dan penyelesaian tepat.

## TABLE OF CONTENTS

CHAPTER	TITLE	PAGE
	<b>REPORT STATUS DECLARATION</b>	
	<b>SUPERVISOR'S DECLARATION</b>	
	<b>TITLE PAGE</b>	i
	<b>DECLARATION</b>	ii
	<b>ACKNOWLEDGEMENTS</b>	iii
	<b>TABLE OF CONTENTS</b>	iv
	<b>ABSTRACT</b>	v
	<b>ABSTRAK</b>	vi
	<b>LIST OF TABLES</b>	viii
	<b>LIST OF FIGURES</b>	ix
	<b>LIST OF APPENDICES</b>	xi
<b>1</b>	<b>RESEARCH FRAMEWORK</b>	<b>2</b>
	1.1 Introduction	2
	1.2 Background of the problem	5
	1.3 Statement of the problem	7
	1.4 Objectives of the study	7
	1.5 Scope of the study	8

<b>2</b>	<b>LITERATURE REVIEW</b>	<b>9</b>
2.1	Introduction	9
2.2	Review of Previous Work	9
2.3	Unbounded Simply Connected Region	11
2.4	The Dirichlet Problem	12
2.5	Integral Equation	13
2.6	The Generalized Neumann Kernel	14
<b>3</b>	<b>THE GREEN'S FUNCTION</b>	<b>17</b>
3.1	Computing Green's function by conformal mapping	17
3.2	Computing Green's function by exterior Dirichlet problem	21
3.3	Integral Equation for the Exterior Dirichlet problem	23
<b>4</b>	<b>NUMERICAL IMPLEMENTATION</b>	<b>25</b>
4.1	Introduction	25
4.2	Discretization of the Integral Equation and Computing the Green's Function	25
4.3	Numerical Examples	30
<b>5</b>	<b>CONCLUSION AND FUTURE WORK</b>	<b>40</b>
5.1	Summary	40
5.2	Suggestions for Further Research	41
	<b>REFERENCES</b>	<b>43</b>
	Appendices A-C	46-51

## LIST OF TABLES

TABLE NO.	TITLE	PAGE
4.1	The error $\ G(z, z_0) - G_n(z, z_0)\ _\infty$ for example 4.1	32
4.2	The error $\ G(z, z_0) - G_n(z, z_0)\ _\infty$ for example 4.2	35
4.3	The error $\ G(z, z_0) - G_n(z, z_0)\ _\infty$ for example 4.3	38

## LIST OF FIGURES

FIGURE NO.	TITLE	PAGE
2.1	Unbounded simply connected region	11
3.1	Ellipse (a=5)	19
3.2	Oval of Cassini (a=1.06)	20
3.3	(a) Bounded region $\Omega$ , (b) Unbounded region $\Omega^-$	21
3.4	Unbounded region $\Omega^-$	22
4.1	The test region $\Omega_1^-$ for Example 1	31
4.2	Green's function for $\Omega_1^-$ in contours form	33
4.3	Green's function for $\Omega_1^-$ in 3D form	33
4.4	The test region $\Omega_2^-$ for Example 2 with a=3	34
4.5	Green's function for $\Omega_2^-$ in contours form	36
4.6	Green's function for $\Omega_2^-$ in 3D form	36
4.7	The test region $\Omega_3^-$ for Example 3 with a=1.06	37
4.8	Green's function for $\Omega_3^-$ in contours form	39
4.9	Green's function for $\Omega_3^-$ in 3D form	39

**LIST OF APPENDICES**

<b>APPENDIX</b>	<b>TITLE</b>	<b>PAGE</b>
A	Computer program for Example 1	46
B	Computer program for Example 2	48
C	Computer program for Example 3	51



## CHAPRER 1

### RESEARCH FRAMEWORK



George Green (1793–1841)

## 1.1 Introduction

Green's functions are presented by the British mathematician George Green (1793-1841), who first developed this concept in 1830s. In the modern linear partial differential equations, Green's functions are analyzed largely from the point of view of fundamental solutions. In the following sections, Green's functions are described in one-dimension and two-dimension space.

### Green's function in one-dimension

Green's function in one-dimension has several applications related to boundary value problems in ordinary differential equations. In this section the Green's function is introduced in the context of a simple one-dimensional problem.

Consider the differential equation in the standard form (Jeffrey, 2001)

$$\frac{d^2y}{dx^2} + a(x)\frac{dy}{dx} + b(x)y = f(x) \quad (1.1)$$

which is defined over the interval  $a \leq x \leq b$ .

Now let  $y_1(x)$  and  $y_2(x)$  be two linearly independent solutions of the homogeneous differential equation, with  $y_1(x)$  such that at  $x = a$  it satisfies the homogeneous boundary condition

$$k_1y_1(a) + k_2y_1'(a) = 0, \quad (1.2)$$

and  $y_2(x)$  such that at  $x = b$  it satisfies the homogeneous boundary condition

$$k_1y_2(b) + k_2y_2'(b) = 0. \quad (1.3)$$

The solution of equation of (1.1) can be written as

$$y(x) = \int_a^x \frac{y_1(t)y_2(x)}{W(t)} f(t)dt + \int_x^b \frac{y_2(x)y_1(t)}{W(t)} f(t)dt, \quad (1.4)$$

where

$$W(x) = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} = y_1 y_2' - y_1' y_2.$$

The solution (1.4) can be written as

$$y(x) = \int_a^b G(x, t) f(t) dt, \quad (1.5)$$

where the function  $G(x, t)$  is called the Green's function for differential equation (1.1) described over the interval  $a \leq x \leq b$ . This function is defined as

$$G(x, t) = \begin{cases} \frac{y_1(t)y_2(x)}{W(t)}, & a \leq t \leq x \\ \frac{y_2(x)y_1(t)}{W(t)}, & x \leq t \leq b. \end{cases} \quad (1.6)$$

The Green's function in (1.5) has the following properties (Jeffrey, 2001):

1. The piecewise defined Green's function  $G(x, t)$  satisfies the differential equation in the respective intervals  $a \leq x \leq t$  and  $t \leq x \leq b$ .
2.  $G(x, t)$  is continuous function of  $x$  for  $a \leq x \leq b$ .
3.  $G(x, t)$  satisfies the homogeneous boundary conditions.
4. The function  $G_x(x, t)$  is continuous for  $a \leq x \leq t$  and  $t \leq x \leq b$ , but it is discontinuous across where it experiences the jump

$$G_x(x, x_+) - G_x(x, x_-) = -1,$$

where  $G_x$  is derivative of  $G$  with respect to  $x$ .

### Green's function in two-dimensions

In this section, we illustrate the use of Green's function in two-dimensions to the boundary value problems in partial differential equations which arise in a wide class of problems in engineering and mathematical physics.

The concept of Green's functions is intimately tied to the Dirac delta function. The Dirac delta function  $\delta(x - \xi, y - \eta)$  in two-dimensions is presented by (Rahman, 2007):

$$I. \delta(x - \xi, y - \eta) = \begin{cases} \infty, & x = \xi, y = \eta \\ 0 & \text{Otherwise} \end{cases}$$

$$II. \iint_{\Omega_\varepsilon} \delta(x - \xi, y - \eta) dx dy = 1, \Gamma_\varepsilon: (x - \xi)^2 + (y - \eta)^2 < \varepsilon^2$$

$$III. \iint_{\Omega} f(x, y) \delta(x - \xi, y - \eta) dx dy = f(\xi, \eta)$$

for arbitrary continuous function  $f(x, y)$  in the region  $\Omega$ .

The application of Green's function in two-dimension can best be described by considering the solution of the Dirichlet problem

$$\nabla^2 u = h(x, y) = 0 \quad \text{in two-dimensional region } \Omega$$

$$u = f(x, y) \quad \text{on the boundary } \Gamma,$$

$$\text{where } \nabla^2 = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}.$$

The Green's function denoted by  $G(x, y; \xi, \eta)$  for the Dirichlet problem involving the Laplace operator is defined as the function which satisfies the following properties:

i-  $\Delta G = \delta(x - \xi, y - \eta)$  in  $\Omega$ ,  $G = 0$  on  $\Gamma$

ii-  $G$  is symmetric, that is,  $G(x, y; \xi, \eta) = G(\xi, \eta; x, y)$

iii-  $G$  is continuous in  $(x, y; \xi, \eta)$  but  $\frac{\partial G}{\partial n}$  the normal derivative has a

discontinuity at the point  $(\xi, \eta)$  which is specified by the equation

$$\lim_{\varepsilon \rightarrow 0} \int_{\Gamma_\varepsilon} \frac{\partial G}{\partial n} ds = 1,$$

where  $n$  is the outward normal to the circle

$$\Gamma_\varepsilon: (x - \xi)^2 + (y - \eta)^2 = \varepsilon^2.$$

**Theorem 1.1** (Rahman, 2007)

The solution of the Dirichlet problem  $\nabla^2 u = h(x, y)$  in  $\Omega$  with the boundary condition  $u = f(x, y)$  on  $\Gamma$  is given by

$$u(x, y) = \iint_{\Omega} G(x, y; \xi, \eta) h(\xi, \eta) d\xi d\eta + \int_{\Gamma} f \frac{\partial G}{\partial n} ds, \quad (1.7)$$

where  $G$  is the Green's function and  $n$  denotes the outward normal to the boundary  $\Gamma$  of the region  $\Omega$ .

**1.2 Background of the problem**

In 1828 George Green (1793–1841) published an Essay on the Application of Mathematical Analysis to the Theory of Electricity and Magnetism. Green's essay remained relatively unknown until it was published between 1850 and 1854. In 1877 Carl Neumann considered the concept of Green's functions in his study of the Laplace's equation.

With the function's success in solving Laplace's equation, other equations began also to be solved using Green's functions. In the case of the heat equation, in 1888 Hobson derived the free-space Green's function for one, two and three dimensions. Indeed, Sommerfeld would be the great champion of Green's functions at the turn of the 20th century because he presented the modern theory of Green's function as it applies to the heat equation (Duffy, 2001).

As mentioned, Green's functions have become a fundamental mathematical technique for solving boundary value problems and other important equation in applied mathematics. Properties of Green's functions for bounded region have been investigated in detail by many authors.

We next give a Green's function on unbounded simply connected region.

Consider the Green's function  $G(x, y; \xi, \eta)$  that satisfies

$$\frac{\partial^2 u(x, y)}{\partial x^2} + \frac{\partial^2 u(x, y)}{\partial y^2} = -\delta(x - \xi, y - \eta), \quad (x, y) \in \Omega \quad (1.8)$$

with boundary condition as

$$u(x, y) = 0, \quad (x, y) \in \Gamma. \quad (1.9)$$

Here  $\Omega$  can be bounded or unbounded region and  $\delta(x, y)$  is the Diract delta-function.

If  $\Omega$  is the unbounded half-plane  $\{-\infty < x < \infty, y > 0\}$ , then the classical Green's function is given by (Hon et al., 2010)

$$G(x, y; \xi, \eta) = \frac{1}{4\pi} \ln \sqrt{\frac{(x - \xi)^2 + (y + \eta)^2}{(x - \xi)^2 + (y - \eta)^2}}. \quad (1.10)$$

Green's function for the disk  $|w| < 1$  is given by

$$G(w, s) = -\frac{1}{2\pi} \ln \left| \frac{w - s}{1 - \bar{s}w} \right|, \quad (1.11)$$

where  $s$  is inside the unit disk and is a pole of  $G$ .

For arbitrary bounded simply connected regions  $\Omega$ , Green's function can now be found by the method of conformal transplantation. Let  $w = f(z)$  map  $\Omega$  conformally onto  $|w| < 1$  and  $\Omega \cup \partial\Omega$  continuously onto  $|w| \leq 1$ . If  $G$  is the Green's function for  $\Omega$ , the function

$$(w, s) \rightarrow g(f^{[-1]}(w), f^{[-1]}(s))$$

has all properties of Green's function for the unit disk, and hence must agree with (1.10). It follows that the desired Green's function for  $\Omega$  is given by (Henrici, 1986)

$$G(z, z_0) = -\frac{1}{2\pi} \ln \left| \frac{f(z) - f(z_0)}{1 - f(z)f(z_0)} \right|. \quad (1.12)$$

In general the Green's function for  $\Omega$  can be expressed by

$$G(z, z_0) = u(z) - \frac{1}{2\pi} \ln |z - z_0|, \quad z, z_0 \in \Omega, \quad (1.13)$$

where  $u$  is the unique solution of the interior Dirichlet problem

$$\begin{cases} \nabla^2 u(z) = 0, & z \in \Omega, \\ u(\eta(t)) = \ln|\eta(t) - z_0|, & \eta(t) \in \Gamma. \end{cases} \quad (1.14)$$

Nasser (2007) has developed a new method for solving the interior and exterior Dirichlet problem in simply connected regions with smooth boundaries. His method is based on two uniquely Fredholm integral equations of the second kind with the generalized Neumann kernel. Recently, his method has been used by Alagele (2012) for computing Green's function on bounded simply connected region only. This research wishes to extend the work by Alagele (2012) for computing Green's function on an unbounded simply connected region by getting a unique solution of the exterior Dirichlet problem using integral equation approach with the generalized Neumann kernel.

### **1.3 Statement of the problem**

This research is to compute the Green's function on an unbounded simply connected region by conformal mapping and by solving an exterior Dirichlet problem via an integral equation with the generalized Neumann kernel.

### **1.4 Objectives of Study**

The objectives of this research are:

- i. To investigate the properties of Green's function for unbounded simply connected region and its connection with the exterior Dirichlet problem and conformal mapping.
- ii. To compute the Green's function on an unbounded simply connected region by using conformal mapping method.
- iii. To compute Green's function on an unbounded simply connected region by solving an exterior Dirichlet problem via an integral equation with the generalized Neumann kernel.
- iv. To create a numerical technique for solving the boundary integral equation using MATHEMATICA.
- v. To compute and graph Green's functions on several test regions.

## **1.5 Scope of Study**

There are Dirichlet problems and Green's function for bounded and unbounded multiply connected regions. The main concern of this research is the evaluation of Green's function on the unbounded simply connected region with smooth boundary. The boundary integral equation method which involves the generalized Neumann kernel and conformal mapping method are considered for a computing Green's function.



## REFERENCE

- Ahlfors, L.V. (1979). *Complex Analysis*. International Student Edition.  
Singapore: McGraw-Hill.
- Ali Hassan Mohamed Murid (1997). *Boundary Integral Equation Approach  
For Numerical Conformal Mapping*. Ph.D. Thesis. Universiti  
Teknologi Malaysia, Skudai.
- Alagele, M. M. A. (2012). *Integral Equation Approach for computing Green's  
Function on Simply Connected Region*. M.Sc Dissertation, Universiti  
Teknologi Malaysia, Skudai.
- Amano, K. (1994). A charge simulation method for the numerical conformal  
mapping of interior, exterior and doubly-connected domains. *J. Comp. Appl.  
Math.* 53, pp. 353-370.
- Asmar, N. H. (2002). *Applied Complex Analysis with Partial Differential  
Equations*. New Jersey: Prentice Hall Inc.
- Atkinson, K.E. (1997). *The Numerical Solution of Integral Equations of the  
Second Kind*. Cambridge: Cambridge University Press.
- Bayin, S. S. (2006). *Mathematical Methods in Science and Engineering*. New  
York: John Wiley.

- Crowdy, D. and Marshall, J. (2007). Green's functions for Laplace's equation in multiply connected domains. *IMA Journal of Applied Mathematics*. 72, pp. 278-301.
- Duffy, D. G. (2001). *Green's Functions with Applications*. New York: Chapman & Hall/CRC.
- Embree, M. and Trefethen, L. N. (1999). Green's Functions For Multiply Connected Domains Via Conformal Mapping. *SIAM Rev.* 41, pp. 745-761.
- Henrici, P. (1986). *Applied and Computational Complex Analysis*, Vol. 3. New York: John Wiley.
- Hon, Y. C, Li, M. and Melnikov, Y. A. (2010). Inverse source identification by Green's function. *Engineering Analysis with Boundary Elements*. 34, pp. 352-358.
- Jeffrey, A. (2002). *Advanced Engineering Mathematics*. Florida: Harcourt press.
- Rahman, M. (2007). *Integral Equation and their Application*. Canada: WIT press.
- Mahanty, J. (1974). *The Green Function Method in Solid State Physics*. New Delhi: Affiliated East-West pr.
- Mikhlin, S. G. (1957). *Integral Equations, English Translation of Russian edition 1948*. Armstrong: Pergamon Press.
- Nasser, M. M. S. (2007). Boundary Integral Equations with the Generalized Neumann Kernel for the Neumann Problem. *MATEMATIKA*. 23, pp. 83-98.
- Nasser, M. M. S. (2009). The Riemann-Hilbert Problem and the Generalized Neumann Kernel on Unbounded Multiply Connected Regions. *The University Research Journal*. 20, pp. 47-60.

- Nurul Akmal Binti Mohamed (2005). Numerical Computation of the Riemann Mapping Function Using Integral Equation Method and Cauchy's Integral Formula. B.Sc Dissertation, Universiti Teknologi Malaysia, Skudai.
- Nasser, M. M. S, Murid, A. H. M, Ismail M., and Alejaily, E. M A. (2011). Boundary integral equations with the generalized Neumann kernel for Laplace's equation in multiply connected regions. *Applied Mathematics and Computation*. 217, pp. 4710-4727.
- Wegmann, R. (2001). Constructive solution of a certain class of Riemann–Hilbert problems on multiply connected circular regions. *Journal of Computational and Applied Mathematics*. 130, pp. 139-161.
- Wegmann, R. Murid, A. H. M, and Nasser, M. M. S. (2005). The Riemann–Hilbert problem and the generalized Neumann kernel. *Journal of Computational and Applied Mathematics*. 2005. 182, pp. 388-415.
- Wegmann, R. and Nasser, M. M. S. (2008). *The Riemann–Hilbert problems and the generalized Neumann kernel on multiply connected regions*. J. Comp. Appl. Math. Vol. 214, pp. 36-57.