

MEASUREMENT OF RADON-222 AND RADIUM-226 IN WELL WATER
OF KELANTAN

NURUL ASHIKIN BINTI SIDI AHMAD

A thesis submitted in fulfillment of the
requirements for the award of the degree of
Master of Science (Physics)

Faculty of Science
Universiti Teknologi Malaysia

OGOS 2013

ABSTRACT

The Environmental Protection Agency (EPA) of USA sets the Maximum Contamination Level (MCL) for ^{222}Rn in drinking water to be 11.1000 Bq/l and for ^{226}Ra to be 0.1100 Bq/l. This study was conducted to determine the levels of ^{222}Rn and ^{226}Ra in private well water in the state of Kelantan. A total of 25 samples of well water were collected from various districts in the state. The levels of ^{222}Rn and ^{226}Ra dissolved in well water were measured using the liquid scintillation counting technique. ^{222}Rn activity concentrations were found within the range of 0.0491 to 4.3127 Bq/l for both unfiltered and filtered samples. Meanwhile, activity concentration of ^{226}Ra measured is between 0.0278 to 1.9670 Bq/l for both unfiltered and filtered samples, exceeding the MCL value. The annual effective doses due to the intake of natural radionuclide from well water for three different age groups were estimated based on the amount of water consumed. The average annual effective dose of ^{226}Ra calculated for babies in the range of 0.12 to 1.79 $\mu\text{Sv/y}$. The average annual effective dose of ^{226}Ra for children was in the range of 0.11 to 0.34 $\mu\text{Sv/y}$ while, for the adults, it was in the range of 0.11 to 0.33 $\mu\text{Sv/y}$ for ^{226}Ra . The annual effective dose intake for the babies was found to be higher than other groups but within the limit recommended by World Health Organization (WHO). The doses for the other two groups are below the recommended value.

ABSTRAK

Agensi Perlindungan Persekitaran atau *Environmental Protection Agency (EPA)* dari Amerika Syarikat menetapkan aras pencemaran maksimum atau *Maximum Contamination Level (MCL)* untuk ^{222}Rn di dalam air minuman adalah 11.1000 Bq/l dan untuk ^{226}Ra adalah 0.1100 Bq/l. Kajian ini dijalankan untuk mengenalpasti aras ^{222}Rn dan ^{226}Ra dalam air perigi persendirian di negeri Kelantan. Sebanyak 25 sampel air perigi telah dikumpul dari pelbagai daerah di negeri tersebut. Aras ^{222}Rn dan ^{226}Ra terlarut dalam air perigi diukur dengan menggunakan teknik pembilang sintilator. Kepekatan aktiviti bagi ^{222}Rn adalah dalam julat 0.0491 sehingga 4.3127 Bq/l untuk sampel tidak dituras dan dituras. Sementara itu, aktiviti bagi ^{226}Ra , kepekatan yang diukur adalah antara 0.0278 sehingga 1.9670 Bq/l untuk kedua-dua sampel yang tidak dituras dan dituras, iaitu melebihi nilai MCL. Dari sampel, boleh anggar dos tahunan yang berkesan untuk pengambilan radionuklid semulajadi dalam air perigi bagi tiga kumpulan umur yang berbeza telah dianggarkan berdasarkan jumlah air yang digunakan. Purata pengambilan dos tahunan berkesan bagi ^{226}Ra yang dikira untuk bayi adalah dalam lingkungan 0.12 sehingga 1.79 $\mu\text{Sv/y}$. Bagi kanak-kanak pula, purata dos tahunan berkesan adalah dalam lingkungan 0.11 sehingga 0.34 $\mu\text{Sv/y}$. Manakala purata pengambilan dos tahunan berkesan untuk orang dewasa dalam lingkungan 0.11 sehingga 0.33 $\mu\text{Sv/y}$. Pengambilan dos tahunan berkesan untuk bayi didapati lebih tinggi daripada kumpulan lain tetapi masih berada dalam had seperti yang disyorkan oleh *World Health Organization (WHO)*. Dos untuk kedua-dua kumpulan yang lain adalah di bawah nilai yang disyorkan.

TABLE OF CONTENTS

CHAPTER	TITLE	PAGE
	DECLARATION	ii
	DEDICATION	iii
	ACKNOWLEDGEMENTS	iv
	ABSTRACT	v
	ABTRAK	vi
	TABLE OF CONTENTS	vii
	LIST OF TABLES	x
	LIST OF FIGURES	xi
	LIST OF SYMBOLS	xii
	LIST OF ABBREVIATIONS	xiv
	LIST OF APPENDICES	xv
1	INTRODUCTION	1
	1.0 Introduction	1
	1.2 Statement of Problem	3
	1.3 Objectives of the Research	5
	1.4 Statement of Hypothesis	5
	1.5 Scope of Study	6
	1.6 Thesis Outlines	6

2	LITERATURE REVIEW	7
2.1	Definition of Radiation and Radioactivity	7
2.2	Type of Ionizing Radiation	9
2.2.1	Alpha Particles	9
2.2.2	Beta Particles	10
2.2.3	Gamma Ray	12
2.3	Natural Radioactivity	13
2.3.1	Uranium	17
2.3.2	Radium	18
2.3.3	Radon	19
2.4	Annual Dose Limit (ADL)	21
2.5	Radioactivity and Related Annual Effective Dose	22
2.6	Liquid Scintillation Counter (LSC)	24
2.7	Routes of Exposure and Health Effect	24
2.8	Maximum Contaminant Level (MCL)	28
2.8.1	MCL for Radium	28
2.8.2	MCL for Radon	29
3	METHODOLOGY	30
3.1	Location Identification	31
3.2	Sample Collection	34
3.3	Sample Preparation	39
3.3.1	Preparation of Liquid Scintillation	40
3.3.2	Labeling of Sample	43
3.3.3	Filtration	44
3.3.4	Shaking	45
3.3.5	Storage	46
3.3.6	Counting	47
3.4	Detection Method	47
3.4.1	Liquid Scintillation Counter (LSC)	48
3.4.2	Global Positioning System (GPS)	49

3.5	Calculation	50
3.5.1	Radioactivity of ^{226}Ra and ^{222}Rn	50
3.6	Estimation of Annual Effective Dose	56
4	ANALYSIS & DISCUSSION	58
4.1	Radioactivity of Radon and Radium	58
4.2	Estimation of Annual Effective Dose	64
5	CONCLUSION & RECOMMENDATIONS	68
5.1.	Conclusions	68
5.2	Suggestions	70
	REFERENCES	71
	APPENDICES A-K	78

LIST OF TABLES

TABLE NO.	TITLE	PAGE
2.1	Alpha emitting of radioactive elements	10
2.2	Beta emitting of radioactive elements	11
2.3	Gamma emitting of radioactive elements	12
2.4	Mass abundance and half-life : Natural Uranium	18
2.5	Physical properties of ²²² Rn	20
2.6	The recommended dose limits in Basic Safety Radiation Protection	22
2.7	The average worldwide effective dose in select Canadian cities compared to worldwide average.	23
2.8	Health effects from the contaminant of radionuclide in drinking water	
2.9	Previous internationally studies average of radon concentration	27
2.10	The MCLs established by USEPA and CDPH	29
3.1	List of Sampling Station and the coordinate	37
3.2	Spike amount of Radium-226 and it activity	51
3.3	The dose conversion factors for ingestion of radionuclide in water	57
4.1	Radon activity in unfiltered and filtered water samples (Bq/l)	59
4.2	Radium activity in unfiltered and filtered water samples (Bq/l)	60
4.3	Average of annual effective dose (mSv/y) for different age groups	65

LIST OF FIGURES

FIGURE NO.	TITLE	PAGE
2.1	The types of radiation in the electromagnetic spectrum	8
2.2	Natural Decay Series: Uranium-238	14
2.3	Natural Decay Series: Thorium- 232	15
3.1	Sampling locations selected from five districts in Kelantan	32
3.2	The yellow pin in map shows the selected sample location	33
3.3	Manual water collections from well-water	35
3.4	The collection of groundwater by pipes system	36
3.5	The water samples collected	39
3.6	PPO, POPOP and distilled water used to prepare the Liquid Scintillation or cocktail	40
3.7	Preparation of Liquid Scintillator	41
3.8	Preparation of standard solution, Radium- 226	42
3.9	Polyethylene bottle were weight and were labeling according to the sampling station.	44
3.10	Water samples were filtered through 0.4 μ m membrane filter paper and connected to water pump	45
3.11	Upper organic of scintillator transfer into plastic vials	46
3.12	Liquid Scintillation Counter	48
3.13	Global Positioning System (GPS)	49
4.1	The radon activities obtained by LSC for unfiltered and filtered water samples	62
4.2	The radium activities obtained by LSC for unfiltered and filtered water samples	63

LIST OF SYMBOLS

A	Alfa
β	Beta
A	Half life
A_o	Original activity
A_t	Current activity
Bq	Becquerel
Bq/l	Becquerel per litre
C	Concentration of radionuclide
C_A	Concentration in Air
C_o	Original Radon concentration in water
C_T	Concentration in Toluene
C_w	Concentration in Water
D	Annual effective dose
D_T	Diffusion coefficient for Radon between Air and Toluene
D_w	Diffusion coefficient for Radon between Air and Water
E	Conversion factor
HNO_3	Nitric Acid
I	Annual intake of drinking water
pCi/l	Pico Curie per little
POPOP	[1, 4-Bis (5-phenyl-2-oxazoly)]
PPO	(2, 5-diphenyloxazole).
Ra	Radium
Rn	Radon
R_N	Total activity of Radon
S	Second

Sv/y	Sievert per year
T	Elapsed time
Th	Thorium
U	Uranium
V _A	Volume of Air
V _t	Volume of Toluene
V _w	Volume of Water
Z	Atomic number
²¹⁴ Bi	Bismuth-214
²²² Rn	Radon-222
²²⁶ Ra	Radium-226
²²⁸ Ac	Actinium-228
²¹² Pb	Lead-212
²³⁴ U	Uranium-234
²³⁵ U	Uranium-235
²³⁸ U	Uranium-238
⁴⁰ K	Potassium-40

LIST OF ABBREVIATIONS

ADL	Annual Dose Limit
AMCL	Alternate Maximum Contaminant Level
ANM	Agency Nuclear Malaysia
BSRP	Basic Safety Radiation Protection
CPM	Cycles per Minute
DLR	Detection Limit for the purpose of Reporting
DPM	Disintegrations per Minute
F	Filtered
GPS	Global Positioning System
KLTN	Kelantan
LSC	Liquid Scintillation Counter
MCL	Maximum Contaminant Level
MMMP	Multi-Media Mitigation Program
ND	Non Detectable
NIRS	National Inorganics and Radionuclide Survey
NORM	Naturally Occurring Radioactive Material
NURE	National Uranium Resource Evaluation
SSI	Swedish Radiation Protection Institute
UF	Unfiltered
UNSCEAR	United Nations Scientific Committee on the Effect of Atomic Radiation
USEPA	United States Environmental Protection Agency
WHO	World Health Organisation
yr	Year

LIST OF APPENDICES

APPENDIX	TITLE	PAGE
A	Temperature recorded for each sample	78
B	The table of density of water	80
C	The calculation to get volume of each bottle	81
D	The mean efficiency for each channel (Region A, Region B and Region C)	87
E	The calculation to get the average value for each cycle	90
F	Table of time separation and the elapsed time	97
G	Radon distribution in Toluene (Dt)	101
H	Radon distribution Between Air and Water (Dw)	103
I	The calculation to get ^{222}Rn and ^{226}Ra activity concentration in water samples	105
J	Table of annual effective dose (mSv/y) intake from radionuclides (^{226}Ra) for different age groups.	136
K	Determination of radium and radon concentration in well water using liquid scintillation method	138

CHAPTER 1

INTRODUCTION

1.0 Introduction

Most sources of drinking water for rural people, especially those living far inland and are still lagging behind in development, most of them get the source of drinking water from groundwater and surface water. Some of the drinking waters are obtained from groundwater sources such as springs, wells and boreholes. Meanwhile, surface water source is obtained from rivers, lakes and streams. Normally, drinking water from groundwater sources has higher concentrations of radon and radium than surface water.

Radon is a radioactive nuclide which is quite short-lived, half of a given amount decay is only 3.82 days. It is also a chemically inert gas and highly soluble in water. The presence of radon in the environment it as the most dangerous radioactive elements than the others. It also do not combine with any particular element and thus move freely in the ground. It is produced continuously in rocks and minerals. Radon is produced through α -decay of Radium-226 and it is a daughter product of the Uranium-238 natural decay series. Radon-222 is easily soluble in water under pressure, especially well water since it is one of the sources of groundwater. High

concentrations of ^{222}Rn are more worrisome to the water users, especially its effects on health to the residents. By drinking water from groundwater, it will increase the risk of human exposure to radiation effects. Almost 90% of stomach cancer deaths caused by inhaling radon released to the indoor air from water. Only about 10% of the deaths were from cancers of internal organs, mostly the stomach, caused by ingestion of radon in water (USEPA, 1999). Cothorn *et al.* (1986) proved that there was a risk of death of about 1-7% from lung cancer for the people of the United States. Most cancer associated with indoor radon levels in buildings due to groundwater resources.

In most countries, the level of radon concentrations measured in the water supply used by the people can usually reached 20 Bq/l. But there are some cases in some countries where it is above 100 Bq/l. However, the results of epidemiological studies found no association between radon content in drinking water and cancer of the digestive system and other systems until now. The World Health Organisation (WHO) has recommended guidelines for the quality of drinking water with the amount of radon in public drinking water sources to be above 100 Bq/l (WHO, 2009).

^{226}Ra is parent of ^{222}Rn and it is an alpha emitter with a half-life of 1600 years. It can be found in groundwater and it supplies a continuous source of ^{222}Rn . The concentration of ^{226}Ra can reach up to a few Bq/l and its present in the system depends on the availability of water or rock. Most people are concerned about the presence of ^{226}Ra in drinking water as it is the main source of water in their daily activities. United State Environmental Protection Agency (USEPA) has reported that the Maximum Contaminant Level (MCL) for the combination of the two radionuclide, $^{226}\text{Ra}+^{228}\text{Ra}$ reported as 0.185 Bq/l. However, ^{226}Ra in drinking water have been reported to be as much as 0.110 Bq/l (USEPA, 2000). Known, ^{228}Ra is owned by Thorium-232 series and it decays by beta emission. ^{228}Ra is controlled by the low solubility of thorium and it is transported on solids. Meanwhile ^{226}Ra is owned by ^{238}U series and it decays by alpha emission. ^{226}Ra can also be transported by water flow by long-term transport of some fertilizers into the waterways. If ^{226}Ra

is present within large amounts in drinking source, it is dangerous as it is the toxic chemical (Zhuo *et al.*, 2001). Among some natural radionuclide present in water resources, only ^{222}Rn and ^{226}Ra can be dangerous to the people who use the water sources in daily activities especially when it is used as drinking water.

When the concentration of ^{226}Ra is at a high level, it will increase health risks, especially its potential as bones seeker. This is a worrying situation especially when it involves the ^{226}Ra content in private well water, where the people who use the water are less concerned about the presence of dangerous radionuclide such as ^{226}Ra in the water. In addition, the rural people are less concerned with the treatment to remove radionuclide in view of the costs payable by them. A survey was conducted by the Swedish Radiation Protection Institute (SSI) indicated that the maximum concentration of radium in groundwater in Sweden was up to 2.5 Bq/l (SSI, 1996).

There are various methods practiced to analyze the ^{222}Rn and ^{226}Ra content in water sources. However, the most widely used method is to use the Liquid Scintillation Counter (LSC). This is because, conventional LSC has an advantage in terms of performance, especially if it involves counting large numbers of samples and analysis of samples can be run in a short period of time. In addition, it can also be used to measure low concentrations of ^{222}Rn .

One of the aims of this study is to measure the concentrations of ^{222}Rn and ^{226}Ra in groundwater from well water by using liquid scintillation techniques. There are many studies that have been conducted using this method, especially for the analysis of aqueous samples of water taken by the many authors (for example Schönhofer, 1989; Beata Kozłowska *et al.*, 1999; Chalupnik *et al.*, 1992; Escobar *et al.*, 1996; Suomela, 1993).

This study focuses on natural radioactivity content in well water samples, collected from several districts in Kelantan. The determinations of natural radionuclide in well water are useful to study the environmental sanitation and human health for those who use the well water for their daily activities.

1.2 Statement of Problem

In rural areas, groundwater is more likely to be the source of drinking water, cooking, cleaning, bathing, and also agricultural activities. Nowadays, most people who use sources of groundwater are not aware of the hazards of radioactive sources from groundwater resources. They use the water source to carry out daily activities, without knowing the presence of radioactive sources in the water. Groundwater is being contaminated due to agricultural runoff or disposal of liquid waste. ^{226}Ra maybe mobile in surficial environment, specifically in the reducing environment but at high level of ^{226}Ra it can cause health risks. Meanwhile ^{222}Rn has the ability to migrate through the pores of the media through by diffusion process, such as through the rock materials. In this study, the method used for the evaluation of ^{222}Rn and ^{226}Ra levels in well water is using LSC.

1.3 Objectives of the Research

The objectives of the study are:

- i) To measure the radioactivity concentration of ^{222}Rn and ^{226}Ra in well water samples collected from Kelantan.
- ii) To estimate the annual effective dose due to the intake of natural radionuclide from well water for three different age groups.

1.4 Statement of Hypothesis

Following are the hypothesis of this study:

- i) There exist of radioisotope ^{222}Rn and ^{226}Ra in groundwater from Kelantan.
- ii) Due to the nature of soil and rock in Kelantan, the activity concentration of ^{226}Ra is probably high.
- iii) Babies are among those who have a higher annual effective dose from ^{226}Ra than children and adults.

1.5 Scope of Study

A sampling of groundwater for two isotopes of the ^{238}U series, ^{222}Rn and ^{226}Ra was conducted in 5 districts in Kelantan to determine the concentration of ^{222}Rn and ^{226}Ra in groundwater from rural area for domestic usage. The method used for the evaluation of natural radionuclide in well water is LSC. The estimation of annual effective dose due to the intakes of groundwater was calculated for three different age groups. The dose estimation study is very useful by determining the amounts of radioactivity intakes and their effects to the human health and populations.

1.6 Thesis Outlines

This thesis consists of five chapters. The first chapter consists of introduction, research objectives, importance of this research, and scope of study and research methodology. Chapter two is the literature review. It covers the work of relevant studies carried out. Chapter three explains the methods and equipment used in the experimental work. Chapter four shows the data obtained from the field measurement, statistical presentation of data using histograms, analysis of groundwater using LSC, and estimate the annual effective dose due to the intake of natural radionuclide for three different age groups. Finally chapter five presents the conclusions of the project and the suggestions.

REFERENCES

- Agency for Toxic Substances and Disease Registry (ATSDR) (1990). *Toxicological Profile for Radium*. Public Health Service, U.S. Department of Health and Human Services, Atlanta, GA.
- Alabdula'aly, Abdulrahman I. (1996). Occurrence of Radon in Riyadh Groundwater Supplies. *Health Physics*. 70 (1), 103-108.
- Alam, M.N., Chowdhry, M.I, Kamal, M., Ghose, S, Islam, M.N and Anwaruddin, M. (1999). Radiological Assessment of Drinking Water of the Chittagong Region of Bangladesh. *Radiat. Prot. Dosim*. 82, 207-214.
- Andrzej Komosa, Ewa Madej, Magdalena Piekarz. (2008). Determination of a Supported Radon Activity Concentration in Bottled Mineral Waters. *Chem. Anal. (Warsaw)*. 53, 835.
- Baretka, J. and Mathew, P.J. (1985). Natural Radioactivity Activity of Australian Building Material, Industrial Wastes and By-products. *Health Physic*. 48(1), 87-95.
- Belloni, P., Cavaioli, M., Ingrao, G., Mancini, C., Notaro, M., Santaroni, P., Torri, G. and Vasselli, R. (1995). Optimization and Comparison of Three Different Methods for the Determination of Rn-222 in Water. *Science of the Total Environment*. 173/174, 61-67.

- Bem, H., and Bou-Rabee, F. (2004). Environmental and Health Consequences of Depleted Uranium Use in the 1991 Gulf War. *Environ. Int.* 30, 123-134.
- Bhasin, J.L. (2001). Impact of New Environmental and Safety Regulations on Uranium Exploration, Mining, Milling and Management of its Waste. *IAEA-TECDOC*. 1244, 189-200.
- Burnett, W.C., Cristoff, J., Stewart, B., Winters, T., and Wilbur, P. (2000). Reliable Gross Alpha/Beta-Particle Analysis of Environmental Samples via Liquid Scintillation Counting. *Radioactivity & Radiochemistry*. 11 (4), 26-44.
- Chalupnik, S. and Lebecka, Jolanta M. (1992). Determination of Ra-226, Ra-228 and Ra-224 in Water and Aqueous Solutions by Liquid Scintillation Counting. In ref. Liquid Scintillation Spectrometry 1992, *Proc. of the Int. Conf on Advances in LSC, LSC 92*. Sept 14-18. Vienna, Austria.
- Cothorn, C., and Rebers, P. A. (1990). *Radon, Radium and Uranium in Drinking Water*. Chelsea, MI: Lewis Publishers.
- Cothorn, C.R., Lappenbusch, W.L., Michel, L. (1986). Drinking Water Contribution to Natural Background. *Health Physics*. 50, 33–39.
- Dijkstra, A., Massart, D.L., Kaufman, L. (1993). *Evaluation and Optimization of Laboratory Methods and Analytical Procedures*. Netherlands. Elsevier Science Publishers.
- Escobar, V.G, Tomě, F.V., Lozano, J.C Sánchez, A.M. (1996). Determination of ^{222}Rn and ^{226}Ra in Aqueous Samples using a Low-level Liquid Scintillation Counter. *Applied Radiation and Isotopes*. 47 (9-10), 861-867.
- Fatima, I., Zaidi, J.H., Arif,M., and Tahir, S.N.A. (2005). Measurement of Natural Radioactivity in Bottled Drinking Water in Pakistan and Consequent Dose Estimates. *Radiation Protection Dosimetry*. 123 (2), 234-240.

- Godoy, J.M., Godoy, M.L. (2006). Natural radioactivity in Brazilian Groundwater. *J. Environ. Radioact.* 85, 71-83.
- Grasty, R.L., and La Marre, J.R. (2004). The Annual Effective Dose from Natural Sources of Ionizing Radiation in Canada. *Radiation Protection Dosimetry.* 108 (3), 215-226. Oxford University Press.
- Holbert, K.E., Stewart, B.D. and Eshraghi, P. (1995). Measurement of Radioactivity in Arizona Groundwater. *Health Physics.* 68 (2), 185-194.
- ICRP. (1991). ICRP Publication 60: 1990 Recommendations of the International Commission on Radiological Protection. J.Valentin (Ed). *International Commission of Radiological Protection, Annals of the ICRP.* 21, 1-3 .United Kingdom: Elsevier.
- Jack, K.H. (1985). *Natural Radioactivity in Water Supplies.* United State: Colorado.
- Juntunen, P., Ruutu, A. and Suksi, J. (2001). Determination of Ra-226 from Rock Samples Using LSC. Mobius.S., Noares.J., and Schönfer.F (Eds). *Advances in Liquid Scintillation Spectrometry* (pp.299-302). Arizona: the Arizona Board of Regent.
- Kaiholo, L., Oikari, T. and Suontausta, J. (1992). Direct Detection of Radon Gas in Air with a Liquid Scintillation Counter. *The 3rd International Conference on Nuclear and Radiochemistry.* September 7-11.Vienna, (Abstract Only).
- Karpińska.M., Mnich.Z.,Kapała.J., and Szpak.A. (2009). The Evaluation of Indoor Radon Exposure in Houses. *Polish J. of Environ. Stud.* 18(6), 1005-1012.
- Khalil, F.A., Rafat M.A., and Ei Fayoumi, M.A.K (2009). Natural Radioactive Nuclides and Chemical Components in the Groundwater of Beni Suef Governate, Middle Egypt. *J. Radiol. Prot.* 29, 105. Retrieved on October 13, 2008 from <http://iopscience.iop.org/0952-4746/29/1/N01>.

- Köhler, M., Preusse, W., Gleisberg, B., Schäfer, I., Heinrich, T., and Knobus, B. (2002). Comparison of Methods for the Analysis of Ra-226 in Water Samples. *Appl. Radiat. Isot.* 56, 387-392.
- Kozłowska, B., Hetman, A., Zipper, W. (1999). Determination of ^{222}Rn in Natural Water Samples from Health Resorts in the Sudety Mountains by the Liquid Scintillation Technique. *Applied Radiation and Isotopes*. 51, 475-480.
- Kozłowska, B., Morelli, D., Walencik, A., Dorda, J., Altamore, I., Chieffalo, V., Giammanco, S., Imme, G., Zipper, W. (2009). Radioactivity in Waters of Mt. Etna (Italy). *Radiation Measurements*. 44, 384-389.
- Kralik, C., Friedrich, M. and Vojir, F. (2003). Natural Radionuclides in Bottled Water in Austria. *J. Environmental Radioactivity*, 65(2), 233-241. Elsevier.
- Kuan, L.S. (2007). *Natural Background Radiation in the Kinta District, Perak, Malaysia*. Master Thesis. Universiti Teknologi Malaysia, Skudai.
- Lowry, J.D. and Lowry, S.B. (1988). Radionuclides in Drinking Water, *J. Am. Assoc. Water Works*, 80, 50-64.
- Malanca, A., Pessina, V. and Dallara, G. (1993). Assessment of the Natural Radioactivity in the Brazillian State of Rio Grande. *Health Physics*. 65, 298-302.
- Mays C.W., Rowland R.E., Stehney A.F. (1985). Canser Risk from the Lifetime Intake of Radium and Uranium Isotopes. *Health Physics*. 48, 635-647.
- National Research Council. (1999). *Risk Assessment of Radon in Drinking Water*. National Academy Press: Washington D.C.

- Radhakrishna, A.P., Somashekarappa, H.M., Narayana, Y. and Siddappa, K. (1993). A new Natural Background Radiation Area on the Southwest Coast of India. *Health Physics*. 65, 390-395.
- Salonen, L. (1988). Natural Radionuclides in Ground Water in Finland. *Radiat. Prot. Dosim.* 24, 163-166.
- Salonen, L. and Hukkanen, H. (1997). Advantages of Low-background Liquid Scintillation Alpha-spectrometry and Pulse Shape Analysis in Measuring Rn-222, Uranium and Ra-226 in Groundwater Samples. *J. Radionucl. Nucl. Chem.* 226 (1-2), 67-74.
- Sanchez, Martin, A., Rubio Montero, M.P., Gomez Escobar, V. and Jurado Vargas, M. (1999). Radioactivity of Bottled Mineral Waters. *Appl. Radiat. Isot.* 50, 1049-1055.
- Schönhofer, F. (1989). Determination of Radon-222 and Radium-226 in Mineral Water and Drinking Water. *A Survey in Austria. Analyst.* 114, 1345-7.
- Schönhofer, F. (1992). Measurement of Ra-226 in Water and Rn-222 in Water and Air by Liquid Scintillation Counting. *Radiation Protection Dosimetry.* 45 (1/5), 123-125.
- Schönhofer, F. (1995). Liquid Scintillation Spectrometry in Environmental Measurements. *The Science of the Total Environment.* 173/174, 29-40.
- Sherman, H.N., Fjeld, R.A., Brown, D.D., and Leybe, J. (1995). Evaluation of a Direct Extraction or Liquid Scintillation Counting Technique for the Measurement of Uranium in Water. *In 3rd Int. Conf on Methods and Applications of Radioanalytical Chemistry, MARC-III.* April 10-16, 1994. Kona, Hawaii. 193(1).

- Siegel, M.D., Bryan, C.R. (2005). Environmental Geochemistry of Radioactive Contamination. In: Sherwood Lollar, B. (Ed.), *Environmental Geochemistry*. (pp. 205—262). Treatise on Geochemistry: Elsevier.
- Soto, J., Fernández, P.L., Gómez, J., Ródenas, C. (1995). Study of the Occurrence of ^{222}Rn and ^{226}Ra in Drinking Water in Spain. *Health Physics*. 69 (6), 961-965.
- SSI. (1996). The Radium Distribution in Some Swedish Soils and Its Effect on Radon Emanation. Swedish Radiation Protection Institute (SSI P983.97; SSI P 1068.98; SSI P 1145.99; SSI P 1198.00; SSI 1228.00). *Swedish Radiation Protection Institute*. Stockholm, Sweden.
- Subba Ramu M.C., Ramachandran T.V., Muraleedhavan T.S., Shaikh A.N. (1990). Indoor Levels of Radon Daughters in some High Background Areas in India. *Radiat Prot Dosit*. 30. 41-44.
- Suomela, J. (1993). Method for Determination of Radon-222 in Water by Liquid Scintillation Counting. SSI-rapport 93-13, ISO Norm: ISO/TC147/SC3/WG6 Work. *Swedish Radiation Protection Institute*. Stockholm, Sweden.
- Tchokossa, P., Olomo, J.B., Osibote, O.A. (1999). Radioactivity in the Community Water Supplies of Ife-Central and Ife-East Local Government Areas of Osun State, Nigeria. *Nuc. Instr. and Meth in Phys. Res. A*. 422(1-3), 784-789.
- Tripathi, R.M, Sahoo, S.K., Jha, V.N., Khan, A.H., Puranik, V.D. (2008). Applied Radiation and Isotopes. Assessment of Environmental Radioactivity at Uranium Mining, Processing and Tailings Management Facility at Jaduguda, India. *Applied Radiation and Isotopes*. 66, 1666- 1670.
- UNSCEAR. (1993). Sources and Effects of Ionizing Radiation. UNSCEAR 1993 Report to the General Assembly. (Ed). *United Nations Scientific Committee on the Effects of Atomic Radiation*. New York.
- UNSCEAR. (2000). Volume I: Sources; Volume II: Effects.(Ed). *United Nations Scientific Committee on the Effects of Atomic Radiation, 2000 Report to the*

General Assembly, with scientific annexes. New York.

- U.S.EPA. (1993). Basic Information about the Radionuclides Rule. *The regulated radioactive drinking water contaminants and health effects.* Washington, D.C.
- U.S.EPA. (1999). Cancer Risk Coefficients for Environmental Exposures to Radionuclides Federal Guidance Report No. 13, EPA 402-R-99-001. *U.S. Environmental Protection Agency.* September 1999. Washington, D.C.
- U.S.EPA. (2000). National Primary Drinking Water Regulations; Radionuclides; Final Rule, 40 CFR Parts 9, 141, and 142 *Federal Register* 65. 236. 76708-76753. December 7, 2000. Washington, D.C.
- Vesterbacka, P. (2007). Natural Radioactivity in Drinking Water in Finland. *Boreal Environment Research.* 12, 11-16.
- Watson, J.E., Evans, J.P., and Mabry, A. M. (1993). Analysis of ^{222}Rn Concentrations in North Carolina Household Water Supplies Derived from Private Wells. *Health Physics.* 65, 156-160.
- WHO. (1993). Guidelines for Drinking Water Quality. *Recommendations.* (2nd Ed). Switzerland: WHO Press.
- WHO. (2004). Guidelines for Drinking Water Quality. *Recommendations.* (3rd Ed). Switzerland: WHO Press.
- WHO. (2009). WHO Handbook on Indoor Radon. *A Public Health Perspective.* Switzerland: WHO Press.
- Zhuo, W., Iida, T., and Yang, X. (2001). Occurrence of Rn-222, Ra-226, Ra-228 and U in Groundwater in Fujian Province, China. *Journal of Environmental Radioactivity.* 53. 111–120.