

**STUDY ON SOLUTIONS OF HEAT PROBLEMS USING FINITE  
DIFFERENCE METHODS AND METHOD OF LINES INCORPORATE  
WITH RK-LIKED METHODS**

**WAN RUKAIDA BINTI WAN ABDULLAH**

This dissertation is submitted as  
a partial fulfillment of the requirements for the award of  
the degree of Master of Science (Mathematics)

Faculty of Science  
Universiti Teknologi Malaysia

OCTOBER 2004

*To my beloved family and friends....*

## ACKNOWLEDGEMENT

First of all, thanks to Almighty Allah for graciously bestowing me the perseverance to undertake this study.

I would like to thank all of those who have supported me throughout the course of this dissertation. Without their time and effort, this endeavor would not have been possible. A special thanks and a deepest appreciation to my supervisor, Associate Prof. Dr. Nazeeruddin Yaacob for his kindness, guidance and valuable comments in completion of this work. A warmest gratitude dedicates to my family for their support and encouragement.

If anyone has had to be patient with me throughout this project, it is my special friend, Mohd Zul-Waqar Mohd Tohid. His love and companionship have helped me to put forth my full effort, and to maintain my sanity.

Finally, I wish to thank my friends for all the wonderful times we have had, it is without doubt that I would not have lasted through the dissertation without your encouragement and support.

## **ABSTRACT**

This dissertation reports a comparison of results from two classes of numerical methods for heat problems. The heat or diffusion equation, which is an example of parabolic equations are classified into the categories of the partial differential equations. Two classes of numerical methods, Method of Lines and Finite Difference Method are discussed. In Method of Lines, several Runge-Kutta methods were incorporated, including the third and fourth order. Finally, analysis on numerical results for the three heat problems is presented.

## ABSTRAK

Disertasi ini melaporkan dapatan kajian daripada dua kaedah berangka bagi menyelesaikan masalah haba. Masalah haba merupakan contoh persamaan parabolik dan dikategorikan dalam persamaan terbitan separa. Dua kaedah berangka iaitu Kaedah Garis dan Kaedah Pembeza Terhingga dibincangkan. Dalam Kaedah Garis, beberapa Kaedah Runge-Kutta digunakan termasuk Kaedah Runge-Kutta peringkat ketiga dan keempat. Akhir sekali, analisis ke atas dapatan berangka untuk tiga masalah haba dihuraikan.

## TABLE OF CONTENTS

CHAPTER	TITLE	PAGE
	<b>ACKNOWLEDGEMENT</b>	iv
	<b>ABSTRACT</b>	v
	<b>ABSTRAK</b>	vi
	<b>CONTENTS</b>	vii
	<b>LIST OF TABLES</b>	xi
	<b>LIST OF FIGURES</b>	xiii
	<b>LIST OF APPENDIXES</b>	xv
<b>CHAPTER I</b>	<b>RESEARCH FRAMEWORK</b>	
	1.1 Introduction	1
	1.2 Problem Statement	2
	1.3 Analysis Factors	6
	1.3.1 Accuracy of the Solution	6
	1.3.2 Computational Complexity	8
	1.3.3 Stability	8
	1.3.4 Simplicity of the Method	9
	1.4 Objectives	9
	1.5 Scope	9
	1.6 Thesis Organization	10

## **CHAPTER II      PARTIAL DIFFERENTIAL EQUATIONS**

2.1	Introduction	11
2.2	Partial Differential Equations	11
2.3	Examples of Partial Differential Equations	15
2.4	Solution of a Parabolic equation	17
	2.4.1 Separation of Variables	17
2.5	Stiff Equations	24

## **CHAPTER III      METHOD OF LINES AND RUNGE-KUTTA METHODS**

3.1	Introduction	25
3.2	The Method of Lines	25
	3.2.1 Example	28
3.3	Runge-Kutta Methods	30
	3.3.1 Error and Order	33
	3.3.2 Runge-Kutta-Liked Methods	35
	3.3.2.1 Heun's Third-Order Formula	35
	3.3.2.2 Kutta's Third Order Rule	37
	3.3.2.3 Stage-Arithmetic Mean	
	Runge-Kutta Method	38
	3.3.2.4 Classical Fourth Order Runge-Kutta Method	40

## **CHAPTER IV      FINITE DIFFERENCE METHODS**

4.1	Introduction	42
4.2	Finite Difference Method	42
4.3	Explicit Method and Implicit Method in a PDE of Parabolic Type	45
4.4	Explicit Methods for Heat Equation	48
4.5	Convergence and Stability	49

## **CHAPTER V      NUMERICAL EXPERIMENTS**

5.1	Introduction	51
5.2	Problem I	52
5.2.1	Solution of the Problem I Using Fourier Series	52
5.2.2	Solution of the Problem I by The Method of Lines	55
5.2.3	Solution of the Problem I Using Finite Difference Method	60
5.3	Problem II	64
5.3.1	Solution of the Problem II Using Separation of Variables	65
5.3.2	Solution of the Problem II by The Method of Lines	67
5.3.3	Solution of the Problem II Using Finite Difference Method	72
5.4	Problem III	76
5.4.1	Solution of the Problem III Using Separation of Variables	76
5.4.2	Solution of the Problem III by The Method of Lines	79
5.4.3	Solution of the Problem III Using Finite Difference Method	84
5.5	Summary of Solutions For The Problems	87

## **CHAPTER VI      COMPARISON AND CONCLUSIONS**

6.1	Comparison	94
6.1.1	accuracy	95
6.1.1.1	Problem I	96
6.1.1.2	Problem II	97



	6.1.1.3 Problem III	96
	6.1.2 Computational Complexity	98
	6.1.3 Stability	100
	6.1.4 Simplicity	110
6.2	Conclusions	111
6.3	Suggestions	114

## **REFERENCES**

### **APPENDIX**

	Appendix A	117
	Appendix B	160
	Appendix C	170

**LIST OF TABLES**

<b>TABLE NO.</b>	<b>TITLE</b>	<b>PAGE</b>
6.1	Analysis of the solutions for Problem I	95
6.2	Analysis of the solutions for Problem II	97
6.3	Analysis of the solutions for Problem III	98
6.4	Computational Complexity	99

## LIST OF FIGURES

FIGURE NO.	TITLE	PAGE
1.1	A thin circular rod	2
2.1	Time-Dependent Problem	15
3.1	Method of Lines	26
3.2	Subdivision of the domain to solve ( 3.7 )	28
4.1	A computational molecule for Finite Difference Method	43
4.2	Explicit Method	45
4.3	Implicit Method	46
4.4	Procedure of Finite Difference Method	47
4.5	Procedure of Explicit Method	47
4.6	Procedure of Implicit Method	48
5.1	Grid of Discretized solutions For Problem I	57
5.2 (a)	Solutions of Problem I at $x = 0.2$ using the Method of Lines	58
5.2 (b)	Solutions of Problem I at $x = 0.4$ using the Method of Lines	58
5.2 (c)	Solutions of Problem I at $x = 0.6$ using the Method of Lines	59
5.2 (d)	Solutions of Problem I at $x = 0.8$ using the Method of Lines	59
5.3 (a)	Solutions of Problem I at $x = 0.2$ using the Finite Difference Method	62
5.3 (b)	Solutions of Problem I at $x = 0.4$ using the Finite Difference Method	62

5.3 (c)	Solutions of Problem I at $x = 0.6$ using the Finite Difference Method	63
5.3 (d)	Solutions of Problem I at $x = 0.8$ using the Finite Difference Method	63
5.4	Grid of Discretized solutions For Problem II	69
5.5 (a)	Solutions of Problem II at $x = 0.2$ using the Method of Lines	70
5.5 (b)	Solutions of Problem II at $x = 0.4$ using the Method of Lines	70
5.5 (c)	Solutions of Problem II at $x = 0.6$ using the Method of Lines	71
5.5 (d)	Solutions of Problem II at $x = 0.8$ using the Method of Lines	71
5.6 (a)	Solutions of Problem II at $x = 0.2$ using the Finite Difference Method	73
5.6 (b)	Solutions of Problem II at $x = 0.4$ using the Finite Difference Method	74
5.6 (c)	Solutions of Problem II at $x = 0.6$ using the Finite Difference Method	74
5.6 (d)	Solutions of Problem II at $x = 0.8$ using the Finite Difference Method	75
5.7	Grid of Discretized solutions For Problem III	81
5.7 (a)	Solutions of Problem III at $x = 0.2$ using the Method of Lines	82
5.7 (b)	Solutions of Problem III at $x = 0.4$ using the Method of Lines	83
5.7 (c)	Solutions of Problem III at $x = 0.6$ using the Method of Lines	83
5.7 (d)	Solutions of Problem III at $x = 0.8$ using the Method of Lines	84
5.8 (a)	Solutions of Problem III at $x = 0.2$ using the Finite Difference Method	85
5.8 (b)	Solutions of Problem III at $x = 0.4$ using the Finite Difference Method	86

5.8 (c)	Solutions of Problem III at $x = 0.6$ using the Finite Difference Method	86
5.8 (d)	Solutions of Problem III at $x = 0.8$ using the Finite Difference Method	87
5.9 (a)	Solutions of Problem I at $x = 0.2$	88
5.9 (b)	Solutions of Problem I at $x = 0.4$	88
5.9 (c)	Solutions of Problem I at $x = 0.6$	89
5.10 (d)	Solutions of Problem I at $x = 0.8$	89
5.10 (a)	Solutions of Problem II at $x = 0.2$	90
5.10 (b)	Solutions of Problem II at $x = 0.4$	90
5.10 (c)	Solutions of Problem II at $x = 0.6$	91
5.10 (d)	Solutions of Problem II at $x = 0.8$	91
5.11 (a)	Solutions of Problem III at $x = 1.0$	92
5.11 (b)	Solutions of Problem III at $x = 2.0$	92
5.11 (c)	Solutions of Problem III at $x = 3.0$	93
5.11 (d)	Solutions of Problem III at $x = 4.0$	93
6.1	Relative errors at several locations ( Heun's Third-Order Formula ) for Problem I	100
6.2	Relative errors at several locations ( Kutta's Third Order Rule ) for Problem I	101
6.3	Relative errors at several locations ( Stage-Arithmetic Mean Runge-Kutta Method ) for Problem I	101
6.4	Relative errors at several locations ( Classical Fourth Order Runge-Kutta Method ) for Problem I	102
6.5	Relative errors at several locations ( Forward Finite Difference Method ) for Problem I	102
6.6	Relative errors at several locations ( Centered Finite Difference Method ) for Problem I	103
6.7	Relative errors at several locations ( Heun's Third-Order Formula ) for Problem II	104
6.8	Relative errors at several locations ( Kutta's Third Order Rule ) for Problem II	104
6.9	Relative errors at several locations ( Stage-Arithmetic Mean Runge-Kutta Method ) for Problem II	105

6.10	Relative errors at several locations ( Classical Fourth Order Runge-Kutta Method ) for Problem II	105
6.11	Relative errors at several locations ( Forward Finite Difference Method ) for Problem II	106
6.12	Relative errors at several locations ( Centered Finite Difference Method ) for Problem II	106
6.13	Relative errors at several locations ( Heun's Third-Order Formula ) for Problem III	107
6.14	Relative errors at several locations ( Kutta's Third Order Rule ) for Problem III	108
6.15	Relative errors at several locations ( Stage-Arithmetic Mean Runge-Kutta Method ) for Problem III	108
6.16	Relative errors at several locations ( Classical Fourth Order Runge-Kutta Method ) for Problem III	109
6.17	Relative errors at several locations ( Forward Finite Difference Method ) for Problem III	109
6.18	Relative errors at several locations ( Centered Finite Difference Method ) for Problem III	110

## LIST OF APPENDIXES

<b>APPENDIXES</b>	<b>TITLE</b>	<b>PAGE</b>
A-1	Heun's Thord Order Formula for Problem I	117
A-2	Kutta's Third Order Rule for Problem I	128
A-3	Stage-Arithmetic Mean Runge-Kutta Method for Problem I	144
A-4	Forward Finite Difference Method for Problem I	156
B-1	Solutions of The Method of Lines for Problem I	160
B-2	Approximation Solutions and Relative Errors For The Finite Difference Method For Problem I	163
B-3	Solutions of The Method of Lines for Problem II	164
B-4	Approximation Solutions and Relative Errors For The Finite Difference Method For Problem II	166
B-5	Solutions of The Method of Lines for Problem III	167
B-6	Approximation Solutions and Relative Errors For The Finite Difference Method For Problem III	169
C-1	Summary of Approximation Solutions Using The Methods For Problem I	170
C-2	Summary of Approximation Solutions Using The Methods For Problem II	172
C-1	Summary of Approximation Solutions Using The Methods For Problem III	174

## CHAPTER I

### RESEARCH FRAMEWORK

#### 1.1 Introduction

The study of heat problems started since in the 18<sup>th</sup> century. A large number of papers have been published on numerical methods for heat problems. However, papers comparing these methods are usually restricted to an examination of the stability of the finite difference schemes used by the methods [Price, 1974].

This study reports a comparison of solutions from two classes of the more important numerical methods for heat problem which is The Method of Lines and Finite Difference Method. We will compare the accuracy, computational complexity, stability and simplicity of the methods for the solutions of heat problems.

Forward Finite Difference Method ( FFDM ) and Centered Finite Difference Method ( CFDM ), which are the two approximations in the Finite Difference Method will be considered. Using The Method of Lines, will we considered the Runge-Kutta methods to find the solutions, which are Heun's Third-Order Formula ( HTOF ), Kutta's Third Order Rule ( KTOR ), Stage-Arithmetic Mean Runge-Kutta Method ( SAMRKM ) and Classical Fourth-Order Runge-Kutta Method ( CFORKM ). We have considered the Runge-Kutta methods because the methods are the most popular choice for numerical solution of Ordinary Differential Equations ( ODE ) in the context of Method of Lines.




## 1.2 Problem Statement

The problem can be stated as follows :

$$\frac{\partial u}{\partial t} = k \frac{\partial^2 u}{\partial x^2}, \quad k > 0. \quad (1.1)$$

The partial differential equation (1.1) is known as the one-dimensional heat equation. Equation ( 1.1 ) occurs in the theory of heat flow – that is, heat transferred by conduction in a rod or in a thin wire. The function  $u(x,t)$  represents temperature at a point  $x$  along the rod at some time  $t$ .



**Figure 1.1 :** A thin circular rod

We have to make many simplifying assumptions; it is worthwhile to see how equation ( 1.1 ) arises. Suppose a thin circular rod of length  $L$  has a cross-sectional area  $A$  and coincide with the  $x$ -axis on the interval  $[0, L]$ . See Figure 1.1. Let us suppose [Zill and Cullen, 2001] :

- The flow of heat within the rod takes place only in the  $x$ -direction.
- The lateral, or curved, surface of the rod is insulated; that is, no heat escapes from this surface.
- No heat is being generated within the rod.
- The rod is homogeneous: that is, its mass per unit volume  $\rho$  is a constant.

- The specific heat  $\gamma$  and thermal conductivity  $K$  of the material of the rod are constant.

To derive the partial differential equation satisfied by the temperature  $u(x,t)$  we need two empirical laws of heat conduction [Zill and Cullen, 2001] :

- The quantity of heat  $Q$  in an element of mass  $m$  is

$$Q = \gamma mu, \quad (1.2)$$

where  $u$  is the temperature of the element.

- The Rate of heat flow  $Q_t$  through the cross-section indicated in Figure 1.1 is proportional to the area  $A$  of the cross-section and the partial derivative with respect to  $x$  of the temperature :

$$Q_t = -KAu_x. \quad (1.3)$$

Since heat flows in the direction of decreasing temperature, the minus sign in ( 1.3 ) is used to ensure that  $Q_t$  is positive for  $u_x < 0$  ( heat flow to the right ) and negative for  $u_x > 0$  ( heat flow to the left ). If the circular slice of the rod shown in Figure 1.1 between  $x$  and  $x + \Delta x$  is very thin, then  $u(x,t)$  can be taken as the approximate temperature at each point in the interval. Now the mass of the slices is  $m = \rho(A\Delta x)$ , and so it follows from ( 1.2 ) that the quantity of heat in it is

$$Q = \gamma \rho A \Delta x u. \quad (1.4)$$

Furthermore, when heat flows in the positive  $x$ -direction, we see from ( 1.3 ) that heat builds up in the slice at the net rate

$$-KAu_x(x,t) - [-KAu_x(x + \Delta x,t)] = KA[u_x(x + \Delta x,t) - u_x(x,t)]. \quad (1.5)$$

By differentiating ( 1.6 ) with respect to  $t$  we see that the net rate is also given by

$$Q_t = \gamma\rho A\Delta x u_t. \quad (1.6)$$

Equating ( 1.5 ) and ( 1.6 ) gives

$$\frac{K}{\gamma\rho} \frac{u_x(x + \Delta x, t) - u_x(x, t)}{\Delta x} = u_t. \quad (1.7)$$

Taking the limit of ( 1.7 ) as  $\Delta x \rightarrow 0$  finally yields ( 1.1 ) in the form

$$\frac{K}{\gamma\rho} u_{xx} = u_t. \quad (1.8)$$

It is customary to let  $k = \frac{K}{\gamma\rho}$  and call this positive constant the thermal diffusivity

[Zill and Cullen, 2001].

In this study, we shall be concerned with finding solutions of the three problems of the one dimensional heat equation with two classes of numerical methods, namely Method of Lines and Finite Difference Method. Our work involves the comparison of these solutions. The three problems are :

### 1. Problem I

The heat problem :

$$u_t = k u_{xx}; \quad 0 < x < 1$$

with

$$u(0, t) = 50^\circ C, \quad (1.9)$$

$$u(1, t) = 20^\circ C; \quad 0 < t < 0.3$$

$$u(x, 0) = 70^\circ C; \quad 0 \leq x \leq 1$$

and  $k = 0.1$  [Rokiah *et al*, 2002]. The problem is considered because the initial value condition consist a constant value.

## 2. Problem II

The boundary value problem :

$$\begin{aligned}
 &u_t = ku_{xx}; && 0 < x < 1 \\
 \text{with} & && \\
 &u(0, t) = 0, && (1.10) \\
 &u(1, t) = 0; && 0 < t < 0.4 \\
 &u(x, 0) = \sin(\pi x); && 0 \leq x \leq 1
 \end{aligned}$$

and  $k = 1$  [Zill and Cullen, 2001]. The problem is considered because the initial value condition consist an oscillation function.

## 3. Problem III

The heat problem :

$$\begin{aligned}
 &u_t = ku_{xx}; && 0 < x < 5 \\
 \text{with} & && (1.11) \\
 &u(0, t) = 10, \\
 &u(5, t) = 30; && 0 < t < 0.30 \\
 &u(x, 0) = 10 - 2x; && 0 \leq x \leq 5
 \end{aligned}$$

and  $k = 1.0$  [McOwen, 1996]. The problem is considered because the initial value condition consist a linear function.

Some problem statements will be considered. The study is focus on the questions as follows :

1. Which is the best numerical method to solve the heat problem between The Method of Lines and Finite Difference Method ?

2. How the procedures of The Method of Lines and Finite Difference Method to solve the heat problem ?
3. What the advantages and disadvantages of the methods ?
4. How to analysis the accuracy, computational complexity, simplicity and stability of the methods ?

### 1.3 Analysis Factors

The aim of the investigations presented was to study the solutions of the two classes of numerical methods, namely Method of Lines incorporate with RK-like methods and The Finite Difference Methods with respect to their ability to simulate the heat problems.

The following points should be looked at in order to judge the applicability of a numerical method:

1. Accuracy of the solution
2. Computational complexity
3. Stability
4. Simplicity of the methods

#### 1.3.1 Accuracy of the Solution

A common way of classifying and comparing methods is to give their order of accuracy [Smith, 1979]. For our purpose, the order of a method qualitatively describes the errors expression. The errors analysis of computations is characterized according to several different criteria. Let  $x$  denote an exact value and let variable  $x^*$  denote an approximation of  $x$ . One may think of  $x^*$  as the result of a computation. The errors which are consider in this study as follows :

1. Average mean square error ( *amse* ),

The *amse* is defined to be

$$amse = \frac{\sum_{i=0}^N |x - x^*|}{N}$$

where  $N$  is a number of the solutions. In the comparison of the numerical methods, the lowest value of *amse* is the best method.

2. Maximum absolute error ( *mae* ),

The absolute error ( *ae* ) is defined to be

$$ae = |x - x^*|,$$

and the maximum absolute error is a maximum or the biggest value of the solutions for a numerical method [ Zill and Cullen, 2001].

3. Root mean square error ( *rmse* ),

The *rmse* is defined to be

$$rmse = \left( \frac{\sum_{i=0}^N (x - x^*)}{N} \right)^{1/2}$$

where  $N$  is a number of the solutions.

#### 4. Maximum relative error ( *mre* ) .

The relative error ( *re* ) is defined to be

$$re = \frac{ae}{|x|} = \frac{|x - x^*|}{|x|}$$

where the maximum relative error is a maximum or the biggest value of the solutions for a numerical method [ Zill and Cullen, 2001]. The relative error is usually the more important error measure : one wants to know how many digits in a computed result are correct [Yakowitz and Szidarovszky, 1986].

### 1.3.2 Computational Complexity

Computational complexity is a total of arithmetic operation numbers for any two real numbers [Norma, 2004]. The arithmetic operation such as plus ( + ), minus ( - ), multiply ( x ) and divide ( ÷ ). Application of trigonometry functions such as sinus, cosinus and tangent, exponent function and absolute function also effect to the computational complexity.

### 1.3.3 Stability.

An important consideration in using numerical methods to approximate the solution of an initial-value problem is the stability of the method. Simply stated, a numerical method is stable if small changes in the initial condition result in only small changes in computed solution [Zill and Cullen, 2001]. A numerical method is said to be unstable if it is not stable.

A numerical method is unstable if round-off errors or any other errors grow too rapidly as the computations proceed [Zill and Cullen, 2001].

#### **1.3.4 Simplicity of the Method.**

Simplicity of the method is a degree of hardness in implements the methods. We can conclude about the simplicity of the method by considered the computational complexity and the time consuming.

#### **1.4 Objective**

The objectives of this study are as follows:

1. To study the partial differential equations and their examples.
2. To study the procedure of the Method of Lines and the Runge-Kutta Methods.
3. To study the procedure of the Finite Difference Methods.
4. To find the better numerical method.
5. To compare the advantages and disadvantages between the methods.

#### **1.5 Scope**

Our work is focusing in the comparison of the solution from the numerical experiments of the heat problems. The comparison is between the Finite Difference Methods and Methods of Lines incorporate with RK-Liked Methods.



## 1.6 Thesis Organization

Chapter II starts with the introduction of partial differential equations. This is followed by some examples of the partial differential equations. The procedure to find the exact value of partial differential equations using the separation of variables is discussed.

Chapter III begins with an introduction to the Methods of Lines and an example of the heat equation to illustrate the procedure of the Method of Lines in partial differential equations. It then touches briefly on the Runge-Kutta ( RK ) methods and the Runge-Kutta Methods which used in the numerical experiment for the heat problems.

Chapter IV is concerned with the Finite Difference Methods. It contains some fundamental in the Finite Difference Methods. Then, we describe the procedure of Explicit Methods and Implicit Methods. This is followed by the discussion of convergence and stability of the Finite Difference Methods.

In chapter V, we discuss briefly the numerical experiments for the three heat problems using Method of Lines and Finite Difference Methods. Method of Lines will incorporate with Heun's Third-Order Formula, Kutta's Third-Order Rule, Stage-Arithmetic Mean Runge-Kutta Method and Classical Fourth Order Runge-Kutta Method. The Finite Difference Methods will use the Forward Difference Formula and Centered Difference Formula.

Chapter VI discusses the comparison of the solution from the numerical experiment in Chapter V and summarizes the conclusions from the comparison. Finally, we summarized the work and give the conclusions. The future works were mentioned here. This is followed by references and appendices.

7. The Final conclusion is The Finite Difference Method is the best method for desk computation but The Method of Lines is the best method for computerize.

### 6.3 Suggestions

1. Application of 5-points finite difference approximation for the Method of Lines may produce more accurate result. This is because we can reduce the errors by using 5-points finite difference approximation. But the method will become more difficult and we need to use software.
2. Using parallel algorithm for Method of Lines with large size of matrix. For example, the 5-points finite difference approximation for the Method of Lines will produce a large size of matrix. We can produce a parallel algorithm using the Parallel Virtual Machine ( PVM ).

## REFERENCES

- Ames, W. F. (1992). *Numerical Methods for Partial Differential Equations*. 3<sup>rd</sup> ed.  
Boston: Academic Press.
- Brezis, H. and Browder, F. (1998). *Partial Differential Equations in the 20<sup>th</sup> Century*.  
London : Prentice Hall.
- Butcher, J. D. (1987). *The Numerical Analysis of Ordinary Differential Equations :  
Ruge-Kutta and General Linear Method*. Chichester : John Wiley & Sons.
- Carnahan, B. and Luther, H. A. (1969). *Applied Numerical Methods*. New York: John  
Wiley & Sons.
- Chapra, S. C. and Canale, R. P. (1988). *Numerical Methods for Engineers*. 2<sup>nd</sup> ed.  
New York: McGraw-Hill Book Company.
- Fröberg, C. E. (1985). *Numerical Mathematics: Theory and Computer Application*.  
California: The Benjamin/Cummings Publishing Company.
- Gautschi, Walter. (1997). *Numerical Analysis: An Introduction*. Boston: Birkhauser.
- Lambert, J. D. (1973). *Computational Methods in Ordinary Differential Equations*.  
London: John Wiley & Sons.
- MacDonald, C. B. (2001) *Constructing High-Order Runge-Kutta Methods with  
Embedded Strong-Stability-Preserving Pairs*. Acadia University : M. Sc.  
Thesis.
- McOwen, R. (1996). *Partial Differential Equations : Methods and Applications*. New  
Jersey : Prentice Hall.
- Nazeeruddin Yaacob (1996). *New Nonlinear Runge-Kutta Methods for Solving Initial  
Value Problems*. Universiti Teknologi Malaysia : Ph. D Thesis.

- Norma Alias (2004). *Pembinaan dan Pelaksanaan Algorithma Selari bagi Kaedah Kelas TTHS dan TTKS dalam Menyelesaikan Persamaan Parabolik pada Sistem Komputer Selari Ingatan Teragih*. Universiti Kebangsaan Malaysia : Ph. D. Thesis.
- Pinsky, M. A. (1998). *Partial Differential Equations and Boundary-Value Problems with Applications*. 3<sup>rd</sup> ed. Boston : McGraw-Hill.
- Price, R. K. (1974). Comparison of Four Numerical Methods for Flood Routing. *Journal of The Hydraulics Division*.
- Rokiah @ Rozita Ahmad and *et. al.* (2002). Solving Stiff Partial Differential Equations Using Explicit Third Order Mean Runge-Kutta Method. *Prosiding Simposium Kebangsaan Sains Matematik ke-XI*. Malaysia : Persatuan Matematik Malaysia.
- Rice, J. R. (1993). *Numerical Methods, Software and Analysis*. 2<sup>nd</sup> ed. Boston: Academic Press.
- Smith, W. A. (1979). *Elementary Numerical Analysis*. New York : Harper & Row, Publishers.
- Yakowitz, S. and Szidarovszky, F. (1986). *An Introduction to Numerical Computations*. New York : MacMillan Publishing Company.
- Zwillinger, D. (1989). *Handbook of Differential Equations*. London : Academic Press Inc.
- Zill, D. G. and Cullen, M. R. (2001). *Differential Equations With Bounday-Value Problems*. United States : Brooks / Cole.