

Draft Genome Sequence of *Jeotgalibacillus soli* DSM 23228, a Bacterium Isolated from Alkaline Sandy Soil

Kian Mau Goh,^a Kok-Gan Chan,^b Amira Suriaty Yaakop,^a Chia Sing Chan,^a Robson Ee,^b Wen-Si Tan,^b Han Ming Gan^c

Faculty of Biosciences and Medical Engineering, Universiti Teknologi Malaysia, Skudai, Johor, Malaysia^a; Division of Genetics and Molecular Biology, Institute of Biological Sciences, Faculty of Science, University of Malaya, Kuala Lumpur, Malaysia^b; Monash School of Science, Monash University Sunway Campus, Petaling Jaya, Selangor, Malaysia^c

***Jeotgalibacillus soli*, a bacterium capable of degrading *N*-acyl homoserine lactone, was isolated from a soil sample in Portugal. *J. soli* constitutes the only *Jeotgalibacillus* species isolated from a nonmarine source. Here, the draft genome, several interesting glycosyl hydrolases, and its putative *N*-acyl homoserine lactonases are presented.**

Received 15 April 2015 Accepted 23 April 2015 Published 21 May 2015

Citation Goh KM, Chan K-G, Yaakop AS, Chan CS, Ee R, Tan W-S, Gan HM. 2015. Draft genome sequence of *Jeotgalibacillus soli* DSM 23228, a bacterium isolated from alkaline sandy soil. *Genome Announc* 3(3):e00512-15. doi:10.1128/genomeA.00512-15.

Copyright © 2015 Goh et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/3.0/).

Address correspondence to Kian Mau Goh, gohkianmau@utm.my.

Jeotgalibacillus is an underexplored halophilic genus of family *Planococcaceae*. The cell wall peptidoglycan of members of this genus is of the A1 α type, linked directly through L-Lys. The major quinones of *Jeotgalibacillus* spp. are MK-7 and MK-8 (1). With the exception of *J. soli* DSM 23228 (also known as strain P9), isolated from alkaline sandy soil, representatives of this genus are associated with marine sources or fermented seafood. *J. soli* has been identified as being strictly aerobic, oxidase and catalase positive, and positive for H₂S production (2). Cells have single polar or subpolar flagella. *J. soli* is distinctive from other *Jeotgalibacillus* spp. in its limited tolerance to NaCl (maximum 9% [wt/vol]). Other species, such as *J. alimentarius*, *J. salarius*, *J. malaysiensis*, and *J. campisalis*, for instance, are able to tolerate concentrations of 15 to 30% (wt/vol) (1, 3, 4).

Strain DSM 23228 was obtained from the Deutsche Sammlung von Mikroorganismen und Zellkulturen, and its genome was sequenced using an Illumina MiSeq sequencer. An average coverage of 200-fold was obtained for the draft genome of 3,776,953 bp in 24 contigs with an *N*₅₀ of 525,494. The *de novo* assembly was performed using SPAdes (5). Gene prediction was carried out using Glimmer 3.02 (6), tRNA prediction with tRNAscan-SE (7), and rRNA prediction with HMMER (8), while BLAST searches were performed against several databases including CatFam, COG, NCBI RefSeq, and SEED.

The G+C content of the *J. soli* genome is 39.7%. The total number of predicted genes is 3,938, and 5 rRNA and 78 tRNA genes were identified. Protein coding genes with predicted functions number 3,040, equivalent to approximately 80% of the total number of predicted genes. Of these, 968 sequences putatively code for catalytic enzymes. The ability of *J. soli* to use starch as a carbon source may be explained by the presence of enzymes (3 α -amylases and 2 pullulanases) that act on α -1,4 and α -1,6 glycosidic bonds. In addition, our detection of the key glycogen-degrading enzyme oligo-1,4-1,6- α -glucosidase is consistent with the capacity of *J. soli* to grow on glycogen. Although β -glucosidase activity was not observed using a standard API 50 CHB/E test

(bioMérieux), a gene encoding this enzyme was identified in the *J. soli* genome. In addition, the *N*-acyl homoserine lactone (AHL) degradation capability of *J. soli* was validated using an AHL inactivation assay performed with *N*-hexanoyl-L-homoserine lactone and *N*-(3-oxohexanoyl)-L-homoserine lactone, and several putative *N*-acyl homoserine lactonases (9, 10) were identified in the genomic sequence. Based on the same assay, *J. alimentarius*, *J. salarius*, *J. malaysiensis*, and *J. campisalis* were found to be unable to degrade AHL.

Nucleotide sequence accession numbers. This whole-genome shotgun project has been deposited in DDBJ/EMBL/GenBank under the accession no. [JXRP00000000](https://www.ncbi.nlm.nih.gov/nuccore/JXRP00000000). The version described in this paper is the first version, [JXRP01000000](https://www.ncbi.nlm.nih.gov/nuccore/JXRP01000000).

ACKNOWLEDGMENTS

This work was supported by the University of Malaya via High Impact Research Grants (UM C/625/1/HIR/MOHE/CHAN/01, grant A-000001-50001 and UM-MOHE HIR grant UM C/625/1/HIR/MOHE/CHAN/14/1, H-50001-A000027), which were awarded to K.-G.C. K.M.G. thanks Universiti Teknologi Malaysia for the awarded GUP grant 06H31. UTM Zamalah also awarded a graduate scholarship to A.S.Y.

REFERENCES

1. Yoon J-H, Weiss N, Lee K-C, Lee I-S, Kang KH, Park Y-H. 2001. *Jeotgalibacillus alimentarius* gen. nov., sp. nov., a novel bacterium isolated from jeotgal with L-lysine in the cell wall, and reclassification of *Bacillus marinus* Rügner 1983 as *Marinibacillus marinus* gen. nov., comb. nov. *Int J Syst Evol Microbiol* 51:2087–2093. <http://dx.doi.org/10.1099/00207713-51.0.2087-0>.
2. Cunha S, Tiago I, Paiva G, Nobre F, Da Costa MS, Verissimo A. 2012. *Jeotgalibacillus soli* sp. nov., a gram-stain-positive bacterium isolated from soil. *Int J Syst Evol Microbiol* 62:608–612. <http://dx.doi.org/10.1099/ijs.0.028878-0>.
3. Yoon J-H, Kang S-J, Schumann P, Oh T-K. 2010. *Jeotgalibacillus salarius* sp. nov., isolated from a marine saltern, and reclassification of *Marinibacillus marinus* and *Marinibacillus campisalis* as *Jeotgalibacillus marinus* comb. nov. and *Jeotgalibacillus campisalis* comb. nov., respectively. *Int J Syst Evol Microbiol* 60:15–20. <http://dx.doi.org/10.1099/ijs.0.008318-0>.
4. Yaakop AS, Chan K-G, Ee R, Kahar UM, Kon WC, Goh KM. 10 April

2015. Isolation of *Jeotgalibacillus malaysiensis* sp. nov. from a sandy beach in Malaysia, with an emended description of the *Jeotgalibacillus* genus. *Int J Syst Evol Microbiol*. <http://dx.doi.org/10.1099/ijs.0.000242>.
5. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <http://dx.doi.org/10.1089/cmb.2012.0021>.
 6. Delcher AL, Harmon D, Kasif S, White O, Salzberg SL. 1999. Improved microbial gene identification with Glimmer. *Nucleic Acids Res* 27: 4636–4641. <http://dx.doi.org/10.1093/nar/27.23.4636>.
 7. Lowe TM, Eddy SR. 1997. tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Res* 25: 955–964. <http://dx.doi.org/10.1093/nar/25.5.0955>.
 8. Finn RD, Clements J, Eddy SR. 2011. HMMER Web server: interactive sequence similarity searching. *Nucleic Acids Res* 39:W29–W37. <http://dx.doi.org/10.1093/nar/gkr367>.
 9. Ghani NA, Norizan SN, Chan XY, Yin WF, Chan KG. 2014. *Labrenzia* sp. BM1: a quorum quenching bacterium that degrades N-acyl homoserine lactones via lactonase activity. *Sensors (Basel)* 14:11760–11769. <http://dx.doi.org/10.3390/s140711760>.
 10. Ghani NA, Sulaiman J, Ismail Z, Chan XY, Yin WF, Chan KG. 2014. *Rhodotorula mucilaginosa*, a quorum quenching yeast exhibiting lactonase activity isolated from a tropical shoreline. *Sensors (Basel)* 14:6463–6473. <http://dx.doi.org/10.3390/s140406463>.