

SEMI-CLOSED CIRCULATION INTEGRATED MULTI-TROPHIC AQUACULTURE  
TREATMENT SYSTEM USING AQUATIC ORGANISMS AS BIOFILTERS  
TO IMPROVE SHRIMP EFFLUENT QUALITY

LAVANIA A/P BALOO

A thesis submitted in fulfilment of the  
requirements for the award of the degree of  
Doctor of Philosophy (Civil Engineering)

Faculty of Civil Engineering  
Universiti Teknologi Malaysia

MARCH 2015

*Dedicated to my lovely*

*Parents, siblings and friends*

*thank you for your support, sacrifice and patience all this while*

## ACKNOWLEDGEMENT

I wish to express my sincere appreciation and deepest gratitude to my supervisor Dr. Shamila Azman, for her valuable guidance, patience and continuous moral support throughout the research period and in preparing this dissertation. My appreciation also extended to my co-supervisor, Assoc. Prof. Dr. Mohd Ismid Mohd Said, for his valuable advices, comments and ideas that he shares with me to endure my research work painstakingly. Without them, I wouldn't complete my research and this dissertation might be meaningless. Completely, they showed me the way for success and nurture myself with inner strength to complete my studies. I would like to convey my heartfelt appreciation to both of my mentors.

I would also like to grab this opportunity to thank all the staff at Environmental Laboratory, such as Mr. Razale, Mr. Ramli, Mr. Azreen, and Mrs. Rahimah for their kind assistance throughout my research period. The time and energy contributed during sampling activity and field work, besides helping me to perform heavy tasks which I can't do it alone as a female student. In addition, my deepest gratitude to Mr. Saberi Mawi, research officer of Fishery Department of Gelang Patah, for providing technical support and advices on the field that I am not familiar with in the beginning.

Not forgetting my lovely family, who have supported me all this while and being there for me whenever I am in a difficult situation. All of you are the reason to make me stronger and confront any obstacles thrown during this journey. You have faith in me that one day I will complete my research successfully, and most importantly for never pressuring me to hurry. You have been the pillar of my achievement.

Last but not least, I also owes special thanks to my friends, who have been there for me, to lend their ears listening to my grumbling, frustration and have assisted me in research work. I deeply appreciate your companions and motivations that sincerely shown to me. You all have made my 4 years the most enjoyable and memorable one.

In addition I wish to acknowledge and would like to thank Ministry of Science Technology and Innovation (MOSTI) for providing National Science Fund (NSF) scholarship, to Universiti Teknologi Malaysia (UTM), and also Ministry of Higher Education for providing Exploratory Research Grant Scheme( ERGS Vot No.: 4L060) for financial support throughout my study period. Finally, my greatest appreciation to the Almighty, to bless me with good health, happiness, contentment, positivity and inner strength during my journey in completing this dissertation.

## ABSTRACT

Rapid development of aquaculture industry has contributed to the degradation of the coastal environment, and the waterways receiving effluent discharged. The wastewater contains high amount of excess nutrients and total suspended solids (TSS). Shrimp wastewater is rich in excretory waste products because the cultured shrimps could assimilate only 23-31% nitrogen and 10-13% phosphorus of the total inputs. This significantly causes water quality deterioration, outbreaks of shrimp diseases and affect shrimp production. This study focused on biological treatment option using macroalgae to balance the negative impact to the ecosystem. Thereby, *Gracilaria edulis* and *Ulva lactuca* were selected and potential as biofilters to improve shrimp water quality has been tested and verified in several laboratory and outdoor tank-scale experiments. Biofiltration potential of macroalgae in outdoor tank shrimp wastewater recirculation system had demonstrated considerably high nutrient removal efficiencies for ammonium, nitrate and phosphate concentrations such as 86%, 53% and 78% for *G. edulis* and 70%, 42% and 90% for *U. lactuca*, respectively. Furthermore, semi-closed circulation integrated multi-trophic aquaculture treatment system fabricated at outdoor of laboratory with integration of tiger shrimp cultivation and treatment units such as sedimentation tank, green lipped mussel in a spray tank cultivation system and macroalgae tank. Sedimentation tank, reduced TSS, chlorophyll-a, and turbidity by 40%, 22% and 43%, respectively. Mussel system depicted reduction of 65%, 67% and 54.0%, respectively. Whereas, macroalgae had presented remarkable removal efficiencies for ammonium, nitrate and phosphate concentrations by 98%, 79% and 89% for *G. edulis* and 85%, 63% and 96% for *U. lactuca*. Besides that, *G. edulis* had exhibited removal for total nitrogen (TN) by 40%. On the other hand, *U. lactuca* had shown greater removal for total phosphorus (TP) with 80% removal efficiency, followed by *G. edulis* by 62.0%. Tissue analysis had demonstrated that the final nitrogen content almost doubled than the initial value in both macroalgae. The final phosphorus content of the tissue in *G. edulis* has doubled and *U. lactuca* has shown an increase of about 1.5 times. In addition, the mean growth rate for *G. edulis* and *U. lactuca* were about 4.4 % d<sup>-1</sup> and 3.7% d<sup>-1</sup> respectively. The mean growth rate of shrimp in treatment tank was 1.31 ± 0.76 % d<sup>-1</sup>, doubled compared to control tank with 100% survival rate. Furthermore, maximum sustainable yield approach revealed the optimum macroalgae biomass for harvest to improve the performance of biofiltration capacity. Thus, *G. edulis* and *U. lactuca* are suitable as biofilters and potential applications of these findings include improvement of shrimp water quality to an acceptable level that ultimately enhance shrimp and macroalgae productivity besides produces an ecologically sustainable treatment and integrated system.

## ABSTRAK

Pembangunan pesat industri akuakultur telah menyumbang kepada kemerosotan persekitaran pantai, dan laluan air penerimaan pelepasan. Air sisa ini mengandungi jumlah yang tinggi dalam lebihan nutrien dan pepejal terampai jumlah (TSS). Air sisa udang kaya dengan bahan buangan perkumuhan kerana udang yang diternak boleh mengasimilasikan hanya 23-31% nitrogen dan 10-13% fosforus daripada jumlah input. Oleh yang demikian, kualiti air udang bertambah teruk, penyebaran serangan penyakit dan menjejaskan penghasilan udang. Kajian ini memberi tumpuan kepada pilihan olahan biologi menggunakan makroalga untuk mengimbangi kesan negatif kepada ekosistem. Dengan itu, *Gracilaria edulis* dan *Ulva lactuca* dipilih dan potensi sebagai biofilter untuk meningkatkan kualiti air udang telah diuji dan disahkan melalui beberapa eksperimen di dalam dan di luar makmal berskala tangki. Potensi makroalga sebagai biofilter di dalam tangki air sisa udang sistem peredaran semula telah menunjukkan efisiensi penyingkiran nutrien yang tinggi untuk ammonium, nitrat dan fosfat seperti 86%, 53% dan 78% bagi *G. edulis* dan 70%, 42% dan 90% bagi *U. lactuca*, masing-masingnya. Tambahan pula, sistem olahan semi-peredaran tertutup 'integrated multi-trophic aquaculture' telah difabrikasi di luar makmal dengan integrasi penternakan udang harimau dan unit olahan seperti tangki pemendapan, kupang di dalam sistem 'spray tank cultivation' dan tangki makroalga. Tangki pemendapan, mengurangkan TSS, klorofil-a, dan kekeruhan sebanyak 40%, 22% dan 43%, masing-masing. Sistem kupang menggambarkan pengurangan 65%, 67% dan 54%, masing-masing. Manakala, makroalga telah menunjukkan potensi penyingkiran yang baik untuk ammonium, nitrat dan fosfat sebanyak 98%, 79% dan 89% bagi *G. edulis* dan 85%, 63% dan 96% untuk *U. lactuca*, masing-masingnya. Di samping itu, *G. edulis* telah mempamerkan penyingkiran untuk jumlah nitrogen (TN) sehingga 40%. Sebaliknya, *U. lactuca* telah menunjukkan penyingkiran lebih besar untuk jumlah fosforus (TP) dengan efisiensi penyingkiran sebanyak 80%, diikuti oleh *G. edulis* sebanyak 62%. Analisis tisu telah menunjukkan bahawa kandungan nitrogen akhir tisu hampir dua kali ganda daripada nilai awal dalam kedua-dua makroalga. Kandungan fosforus akhir tisu dalam *G. edulis* juga telah bertambah dua kali ganda, manakala *U. lactuca* telah menunjukkan peningkatan lebih kurang 1.5 kali dari nilai awal. Purata kadar pertumbuhan untuk *G. edulis* dan *U. lactuca*, masing-masing adalah  $4.4 \% d^{-1}$  dan  $3.7 \% d^{-1}$ . Sebaliknya, udang telah mencatatkan kadar pertumbuhan dengan nilai purata  $1.31 \pm 0.76 \% d^{-1}$ , lebih dua kali ganda berbanding udang di tangki kontrol dengan 100% kadar hidup. Tambahan pula, pendekatan 'maximum sustainable yield' mendedahkan nilai biojisim makroalga optimum untuk tuaian bagi meningkatkan prestasi keupayaan biofiltrasi. Maka, *G. edulis* dan *U. lactuca* adalah sesuai sebagai biofilter dan aplikasi penemuan ini termasuk peningkatan kualiti air udang kepada tahap sesuai yang akhirnya juga meningkatkan produktiviti udang dan makroalga, selain menghasilkan sistem olahan dan integrasi yang seimbang dari segi ekologi.

## TABLE OF CONTENT

CHAPTER	TITLE	PAGE
	DECLARATION	ii
	DEDICATION	iii
	ACKNOWLEDGEMENT	iv
	ABSTRACT	vi
	ABSTRAK	vii
	TABLE OF CONTENT	viii
	LIST OF TABLES	xv
	LIST OF FIGURES	xviii
	LIST OF ABBREVIATIONS	xxvii
	LIST OF SYMBOLS	xxix
	LIST OF APPENDICES	xxx
<b>1</b>	<b>INTRODUCTION</b>	<b>1</b>
	1.1 Background of the Study	1
	1.1.1 Preamble of Macroalgae	3
	1.2 Problem Statement	5
	1.3 Objectives of the Study	7
	1.4 Scope of the Study	8
	1.5 Significance of the Study	9

1.6	Thesis Organization	11
<b>2</b>	<b>LITERATURE REVIEW</b>	<b>12</b>
2.1	Global Macroalgae Production	12
2.2	Prospect of Macroalgae Cultivation in Malaysia	14
2.2.1	The Perspective and Challenges of Macroalgae Cultivation in Malaysia	15
2.3	<i>Gracilaria</i> and <i>Ulva</i> Species as Ecological Engineering Tools	18
2.3.1	<i>Gracilaria</i> sp.	18
2.3.1.1	Cultivation Methods of <i>Gracilaria</i> sp.	20
2.3.2	<i>Ulva</i> sp.	23
2.3.2.1	Cultivation Methods of <i>Ulva</i> sp.	25
2.4	Nitrogen Requirement by <i>Gracilaria</i> and <i>Ulva</i> Species for Growth	26
2.5	Shrimp Farming Industry	28
2.6	Black Tiger Shrimp ( <i>Penaeus monodon</i> )	29
2.7	An Overview of Malaysian Shrimp Production	31
2.8	Shrimp Cultivation Practices and Evolution of Cultivation Methods	32
2.8.1	Intensive Cultivation of Shrimp	34
2.9	Shrimp Wastewater Characterization	36
2.10	Shrimp Pond and Nutrient Budget	38
2.11	Potential Solutions to Improve Shrimp Water Quality	41



2.12	Integrated Culture of Macroalgae with Aquaculture Wastewater	48
2.13	Potential of <i>Gracilaria</i> sp. as Biofilter and its Application in Integrated System	49
2.14	Potential of <i>Ulva</i> sp. as Biofilter and its Application in Integrated System	53
2.15	Integrated Multi-trophic Aquaculture (IMTA) System	61
2.16	Potential and Combination of Varying Species in an IMTA System	63
2.17	Transformation of IMTA System	65
2.18	Maximum Sustainable Yield (MSY) Concept for Macroalgae Biomass	69
<b>3</b>	<b>RESEARCH METHODOLOGY</b>	<b>71</b>
3.1	Introduction	71
3.2	Sampling Location	73
3.3	Specimens Collection and Acclimatization	75
3.4	Materials	76
3.5	Preliminary Study on Effect of Carbon Dioxide Concentration on <i>G. edulis</i> Growth	77
3.5.1	Batch Scale Laboratory Experiment	77
3.5.2	Tank Scale Experiment for CO <sub>2</sub> Utilisation by <i>G.edulis</i>	79
3.6	Experimental Design for Biofiltration Studies	80
3.6.1	Preliminary Study of <i>G. edulis</i> Growth in PES medium	82

3.6.1.1	Preparation of Pre-culture for Experimental Purposes	82
3.6.1.2	Macroalgae Culture in Provasoli Enriched Seawater (PES) Medium	83
3.6.1.3	Growth of <i>G. edulis</i> in Varying Volume of PES Medium	85
3.6.1.4	Growth of <i>G. edulis</i> in Varying Components of PES Medium	86
3.6.2	Biofiltration Capacity and Kinetics of Nutrient Uptake Rate for $\text{NH}_4^+$ , $\text{NO}_3^-$ , and $\text{PO}_4^{3-}$ using Synthetic Condition	87
3.6.3	Biofiltration Potential of Macroalgae for Nutrient Removal in Outdoor Tank Shrimp Wastewater Recirculation System	89
3.6.4	Biofiltration Potential of Macroalgae for Ammonium Removal in Outdoor Tank Shrimp Wastewater Recirculation System	91
3.6.4.1	Experimental Design for Outdoor Shrimp Recirculation System	92
3.6.4.2	Laboratory Experiment for Ammonium Removal	93
3.7	Semi-closed Circulation Integrated Multi-Trophic Aquaculture (IMTA) System	93
3.8	Kinetic of Nutrient Uptake Rate using Michealis- Menten Approach	98
3.9	Characterization Studies	98
3.9.1	Physical Analysis Procedures	100
3.9.2	Chemical Analysis of the Water Sample	100
3.10	Proximate Analysis of Macroalgae Tissues	101
3.11	Tissue Content Analysis for Metals	102

	3.12	Maximum Sustainable Yield (MSY) of Macroalgae Production	103
	3.13	Statistical Analysis	106
<b>4</b>		<b>FACTORS AFFECTING MACROALGAE GROWTH: LABORATORY SCALE EXPERIMENTS</b>	<b>108</b>
	4.1	Introduction	108
	4.2	Macroalgae Characterization	109
	4.2.1	Physical Characteristics of Macroalgae	110
	4.2.2	Physicochemical Characterization of Macroalgae Environment	112
	4.3	Proximate Analysis of Macroalgae Tissue	114
	4.3.1	Nitrogen and Phosphorus Profile of Macroalgae Tissue	117
	4.4	Carbon Dioxide Acquisition by <i>G. edulis</i> and Its Effect on Growth	121
	4.4.1	Batch Scale Laboratory Experiment for CO <sub>2</sub> Utilization	121
	4.4.2	Tank Scale Laboratory Experiment for CO <sub>2</sub> Utilization	128
	4.5	<i>G. edulis</i> Cultivation Techniques under Controlled Environment	130
	4.5.1	Effect of PES Volume Medium on <i>G. edulis</i> Growth	130
	4.5.2	Effect of Different PES Medium Components on <i>G. edulis</i> Growth	132
	4.6	Biofiltration Capacity and Kinetics of Nutrient Uptake Rate by <i>G. edulis</i> and <i>U. lactuca</i>	135
	4.7	Conclusions	146

<b>5</b>	<b>BIOFILTRATION OF NUTRIENTS IN OUTDOOR TANK SHRIMP WASTEWATER RECIRCULATION SYSTEM</b>	<b>148</b>
5.1	Introduction	148
5.2	Biofiltration Capacity for Ammonium, Nitrate and Phosphate by <i>G. edulis</i> and <i>U. lactuca</i>	149
5.2.1	Physical Parameters and Nutrient Removal Efficiency	150
5.2.2	Specific Growth Rate of Macroalgae and Shrimp	155
5.2.3	Tissue Content of Macroalgae in Treatment Tanks	157
5.3	Ammonium Removal in Outdoor Tank Shrimp Wastewater Recirculation System by Macroalgae	163
5.3.1	Ammonium Removal for Two Shrimp Growth Period	164
5.3.2	Laboratory Batch Experiment	166
5.3.3	Specific Growth Rate of Macroalgae and Shrimp	167
5.4	Conclusions	170
<b>6</b>	<b>SEMI-CLOSED CIRCULATION INTEGRATED MULTI-TROPHIC AQUACULTURE (IMTA) SYSTEM</b>	<b>172</b>
6.1	Introduction	172
6.2	Water Quality Parameters for Shrimp Cultivation	173
6.3	Nitrogen Cycle and Profile in Shrimp Wastewater	181
6.4	Sedimentation Tank Performance in a Semi-closed Circulation IMTA system	183

6.5	Mussel Filtration of Shrimp Wastewater and its Effect on Water Quality	188
6.5.1	Mussel Cultivation Water Characteristics in a Semi-closed Circulation System	189
6.5.2	Determination of Effect of Mussel Density and Survival Rate	193
6.5.3	Mussel Spray Tank Cultivation System Performance in a Semi-closed IMTA System	196
6.6	Macroalgae as Biofilters in a Semi-closed Circulation IMTA System	202
6.6.1	Physical Parameters of Water Quality in Macroalgae Treatment Tank	202
6.6.2	Nutrient Removal by <i>G. edulis</i> and <i>U. lactuca</i> in a Semi-closed Circulation IMTA System	208
6.6.3	Macroalgae Tissue Analysis of <i>G. edulis</i> and <i>U. lactuca</i>	214
6.6.4	Macroalgae Specific Growth Rate in a semi-closed Circulation IMTA System	218
6.6.5	Shrimp Growth Rate and Productivity in a Semi-closed Circulation IMTA System	221
6.6.6	Maximum Sustainable Yield (MSY) of Macroalgae	224
6.7	Conclusions of Treatment Efficiency in a Semi-closed Circulation IMTA System	229
<b>7</b>	<b>CONCLUSIONS AND RECOMMENDATIONS</b>	<b>232</b>
7.1	Conclusions	233
7.2	Recommendations	235
	<b>REFERENCES</b>	<b>237</b>
	Appendices A-E	270-304

**LIST OF TABLES**

<b>TABLE NO.</b>	<b>TITLE</b>	<b>PAGE</b>
2.1	Macroalgae production in Malaysia.	15
2.2	Taxonomic classification of <i>Gracilaria edulis</i> (Guiry and Guiry, 2010)	19
2.3	Taxonomic classification of <i>Ulva lactuca</i> (Guiry and Guiry, 2010)	25
2.4	Common shrimp species utilized in shrimp farming (FAO, 1989).	28
2.5	Life cycle of giant tiger prawn, <i>Penaeus monodon</i> (Motoh, 1985).	30
2.6	Comparison of brackish water shrimp farming systems (Anh <i>et al.</i> , 2010)	35
2.7	Recommended water quality criteria for shrimp cultivation (Boyd and Gautier, 2000)	36
2.8	Shrimp wastewater characteristics based on intensive cultivation and data collected by various researchers	37
2.9	Summary of treatment options for shrimp wastewater	45

2.10	Existing integrated system of varying species, aquaculture wastewaters and cultivation methods	58
2.11	Tank cultivation of varying species of <i>Ulva</i> sp. and <i>Gracilaria</i> sp. shows nutrient removal efficiency	60
2.12	Conceptual flow of ecologically sustainable aquaculture-macroalgae based integrated system	70
3.1	List of equipments used throughout the study	76
3.2	Phases of the research for nutrient biofiltration from lab-scale study to application on fabricated outdoor tank-scale study	81
3.3	Provasoli enriched natural seawater (PES) medium components (Andersen, 2005).	84
3.4	PES medium formulated with different components for <i>G. edulis</i> growth	87
3.5	Conditions of synthetic medium for nutrient uptake rate	88
4.1	The taxonomic classification of red and green macroalgae species	110
4.2	Physicochemical characteristics of macroalgae environment	113
4.3	Proximate analysis for macroalgae tissue content in dry weight basis. Data represent mean, range, and T-test values	115
5.1	Means of water physical characteristics of recirculation system.	151

5.2	Specific growth rate measurement of both macroalgae and also shrimp in each treatment aquarium	157
5.3	Tissue C, N, P contents and ratios of C:N and N:P in <i>G. edulis</i> and <i>U. lactuca</i> , with significant t-test values.	159
5.4	Specific growth rates of macroalgae and shrimp at two different shrimp growout periods in a shrimp wastewater outdoor recirculation system.	168
6.1	Characteristics of shrimp water quality cultivated in a polyethylene tank throughout shrimp growout period	175
6.2	Correlation coefficients (n = 6) between environmental parameters during the study period	178
6.3	Chemical and physical characteristics of mussel spray cultivation tank and on-site mussel cultivation	190
6.4	Analysis of mussel survival rate at four different stocking densities	193
6.5	Tissue C, N, P and ratios of C:N and N:P in <i>G. edulis</i> and <i>U. lactuca</i> , with significant t-test values	217



## LIST OF FIGURES

<b>FIGURES NO.</b>	<b>TITLE</b>	<b>PAGE</b>
1.1	Associated problems of shrimp pond effluent	5
2.1	<i>G. edulis</i> with irregular branching form (Guiry and Guiry, 2010)	20
2.2	<i>U. lactuca</i> in large sheet form (Guiry and Guiry, 2010)	23
2.3	Adult tiger shrimp during harvesting phase after 15 month	29
2.4	Total aquaculture production in Malaysia (FOA, 2012)	31
2.5	(a) Nitrogen sources and (b) fates in shrimp pond. Loss of nitrogen represents loss by denitrification and volatilization process (Jackson <i>et al.</i> , 2003)	39
2.6	Schematic diagram of a semi-recirculating integrated system (Neori <i>et al.</i> , 2003)	56
2.7	Schematic diagram of 25% recirculation unit and flow-through unit (Robertson-Andersson <i>et al.</i> , 2008)	57
2.8	Nitrogen and phosphorus budget in IMTA system. Reduction in the amount of nutrients is significant (Crab <i>et al.</i> , 2007)	62

2.9	An IMTA site in the Bay of Fundy, Canada with salmon cages on the left, a mussel raft on the right foreground and a macroalgae raft on the right background (Guerrero <i>et al.</i> , 2012)	63
2.10	Transformation of aquaculture wastewater treatment options and subsequent introduction of semi-closed recirculation IMTA system	66
2.11	Consideration criteria for selection of suitable macroalgae species	68
2.12	Advantages of IMTA system over monoculture	68
3.1	Research framework of the study	72
3.2	Map of sampling stations which is located nearby Johor Straits	73
3.3	Specimens (a), (b) and (c) were collected from site and (d) reared for experimental purposes	74
3.4	a) <i>G edulis</i> grown in brackish water and b) in shrimp effluent, with the presence of carbon dioxide gas	79
3.5	a) <i>G edulis</i> grown in brackish water and b) in shrimp effluent, without supply of carbon dioxide gas (control)	80
3.6	A series of conical flasks that contains varying volume of PES medium for <i>G. edulis</i> culture	86
3.7	Different concentrations of synthetic medium to determine nutrient uptake rate	89

3.8	Experimental set-up of outdoor tank shrimp wastewater recirculation system where a) stocked with <i>G. edulis</i> , b) stocked with <i>U. lactuca</i> and c) control	91
3.9	Schematic diagram of fabricated semi-closed circulation integrated multi-trophic aquaculture (IMTA) system, where (1) sedimentation tank, (2) mussel spray culture tank, (3) <i>G. edulis</i> and <i>U. lactuca</i> tanks and (4) collection tank	94
3.10	Shrimp cultivation tank and (b) gravitational flow of wastewater through pipe to sedimentation tank	95
3.11	Wastewater flows to the sedimentation, mussel and macroalgae tanks, respectively	96
3.12	Characterization studies and parameters that have been tested for analysis	99
3.13	Logistic growth curve as function of time	104
3.14	Maximum sustain harvest model and the value represented by the peak	106
4.1	Herbarium specimen of <i>Gracilaria edulis</i>	111
4.2	Herbarium specimen of <i>Ulva lactuca</i>	112
4.3	Correlation of tissue nitrogen content of <i>U. lactuca</i> and dissolved inorganic nitrogen content in the water sample	118
4.4	Correlation of tissue nitrogen content of <i>G. edulis</i> and dissolved inorganic nitrogen content in water sample	118
4.5	Correlation of tissue phosphorus content of <i>U. lactuca</i> and dissolved phosphorus content in the water sample	119

4.6	Correlation of tissue phosphorus content of <i>G. edulis</i> and dissolved phosphorus content in the water sample	120
4.7	Macroalgae growth in culture media aerated with normal air, and enriched carbon dioxide air over a 14 days period	122
4.8	Diurnal variations of pH during light and dark period at normal atmospheric carbon dioxide air, and enriched carbon dioxide air at 700 ppm and 1400 ppm, respectively	123
4.9	Response of photosynthetic O <sub>2</sub> evolution to inorganic carbon concentration. Circle marker represents experimental data and the line graph is an exponential curve obtained using curve expert software	125
4.10	<i>G. edulis</i> growth rate in the presence of carbon dioxide and control, in brackish water and shrimp effluent	128
4.11	Growth of <i>G. edulis</i> fresh weight at different volume of PES medium.	131
4.12	Mean daily growth rate of <i>G. edulis</i> in varying PES medium volume. Bar marked with same letters were not significantly different according to DMRT (P=0.05)	131
4.13	Treatment of different composition of PES medium and its response to <i>G. edulis</i> daily growth rate. Each data point is the mean of five replicates (means $\pm$ standard deviation). Bar marked with same letters were not significantly different according to DMRT (P=0.05)	133

4.14	Depletion of ammonium concentrations during 8 hours experiment. U and G represents <i>U. lactuca</i> and <i>G. edulis</i> respectively at different concentrations of U1& G1-25 $\mu$ M; U2&G2-50 $\mu$ M and U3&G3-100 $\mu$ M	137
4.15	Ammonium uptake rate of <i>G. edulis</i> and <i>U. lactuca</i> as a function of substrate concentration with data fitted to Michealis- Menten equation	138
4.16	Depletion of nitrate concentrations during 8 hours experiment. U and G represents <i>U. lactuca</i> and <i>G. edulis</i> respectively at different concentrations of U1& G1-25 $\mu$ M; U2&G2-50 $\mu$ M and U3&G3-100 $\mu$ M	139
4.17	Nitrate uptake rate of <i>G. edulis</i> and <i>U. lactuca</i> as a function of substrate concentration with data fitted to Michealis-Menten equation	140
4.18	Depletion of phosphate concentrations during 8 hours experiment. U and G represents <i>U. lactuca</i> and <i>G. edulis</i> respectively at different concentrations of U1& G1-25 $\mu$ M; U2&G2-50 $\mu$ M and U3&G3-100 $\mu$ M	141
4.19	Phosphate uptake rate of <i>G. edulis</i> and <i>U. lactuca</i> as a function of substrate concentration with data fitted to Michealis-Menten equation	142
5.1	Ammonium removal efficiency by <i>G. edulis</i> and <i>U. lactuca</i> in a shrimp integrated recirculation system, where A1-G: aquarium with <i>G. edulis</i> ; A2-U: aquarium with <i>U. lactuca</i> ; A3-C: aquarium without macroalgae	152

5.2	Nitrate removal efficiency by <i>G. edulis</i> and <i>U. lactuca</i> in a shrimp integrated recirculation system, where A1-G: aquarium with <i>G. edulis</i> ; A2-U: aquarium with <i>U. lactuca</i> ; A3-C: aquarium without macroalgae	153
5.3	Phosphate removal efficiency by <i>G. edulis</i> and <i>U. lactuca</i> in a shrimp integrated recirculation system, where A1-G: aquarium with <i>G. edulis</i> ; A2-U: aquarium with <i>U. lactuca</i> ; A3-C: aquarium without macroalgae	154
5.4	Specific growth rate of both macroalgae during the cultivation period integrated with shrimp cultivation. Each data point is the mean of five replicates (means $\pm$ SD). Bar marked with the same letters are not significantly different according to DMRT analysis (P=0.05)	156
5.5	Water physico-chemical parameters of two different shrimp growout periods in the integrated outdoor recirculation system	164
5.6	The initial and final concentration of ammonium after 14 days are shown as bars. The ‘*’ mark indicates the removal rates. G1, U1, C1 and G2, U2, C2 are <i>G. edulis</i> , <i>U. lactuca</i> and control during 60 days and 120 days of shrimp growout periods, respectively	165
5.7	The initial and final ammonium concentration after 14 days are shown as bars. The ‘*’ mark indicates removal rates of ammonium in a laboratory batch culture experiment	167
6.1	Sedimentation tank performance at different retention time period	184

6.2	Turbidity and chlorophyll-a reduction efficiency in sedimentation tank	185
6.3	Dissolved inorganic nutrients reduction efficiency in sedimentation tank	186
6.4	Effect of mussel stocking density on its survival rate in shrimp wastewater. Standard deviation bar is included, however the value is too small	194
6.5	Removal efficiency of TSS, chl-a, turbidity, TN and TP at low, medium and high mussel stocking density, respectively. Positive change represents an increase and negative change represents a decrease. Standard deviation bar is included	195
6.6	Reduction of total suspended solid, chlorophyll-a, and turbidity in mussel spray cultivation tank within two weeks of study period	197
6.7	Mean ammonium and nitrate profiles in mussel spray tank cultivation system during two weeks of study period	198
6.8	Mean phosphate profile in mussel spray tank cultivation system during two weeks of study period	199
6.9	TN and TP nutrients profile in mussel spray tank cultivation system during two weeks of study period	200
6.10	Physical parameters in <i>G. edulis</i> , <i>U. lactuca</i> and control tanks during the study period	204
6.11	TSS concentrations in macroalgae and control tank during two weeks of treatment period	206

6.12	Chlorophyll-a ( $\mu\text{g/L}$ ) and turbidity (NTU) concentrations in macroalgae and control tank during two weeks of treatment period	207
6.13	Ammonium removal in macroalgae treatment and control tank	208
6.14	Nitrate removal in macroalgae treatment and control tank	209
6.15	Phosphate removal in macroalgae treatment and control tank	210
6.16	TN and TP removal efficiency by (a) <i>G. edulis</i> and (b) <i>U. lactuca</i>	211
6.17	Specific growth rate of both macroalgae during the cultivation period semi-closed IMTA system. Each data point is the mean of three replicates (means $\pm$ SD). Bar marked with same letters are not significantly different according to DMRT test ( $P=0.05$ )	219
6.18	Shrimp body weight gain in treatment and control tank throughout the shrimp rearing period	222
6.19	Shrimp body length in treatment and control tank throughout the shrimp rearing period	222
6.20	Harvested shrimp from the treatment tank at the completion of the experiment	224
6.21	Biomass growth of <i>G. edulis</i> ( $\circ$ ) and <i>U. lactuca</i> ( $\bullet$ ) throughout the treatment where the circle marks represents experimental data and line graph is simulated data from the model. $K_G$ and $K_U$ are the carrying capacity for <i>G. edulis</i> and <i>U. lactuca</i> , respectively	225



6.22	MSY prediction for <i>G. edulis</i> and maximum harvest value	226
6.23	MSY prediction for <i>U. lactuca</i> and maximum harvest value	227
6.24	Conclusion of the findings obtained for semi-closed recirculation IMTA system	231

**LIST OF ABBREVIATIONS**

BOD	-	Biochemical oxygen demand
C	-	Carbon
Chl-a	-	Chlorophyll-a
C:N	-	Carbon to nitrogen
CO <sub>2</sub>	-	Carbon dioxide
COD	-	Chemical oxygen demand
DMRT	-	Duncan multiple range test
DO	-	Dissolved oxygen
D.wt	-	Dry weight
EPA	-	Environmental Protection Act
FAO	-	Food and Agriculture Organization
FCR	-	Food conversion ratio
<i>G. edulis</i>	-	<i>Gracilaria edulis</i>
HDPE	-	high density polyethylene
IMTA	-	Integrated Multi-trophic Aquaculture
L:D	-	Light and dark
MSY	-	Maximum sustainable yield
N	-	Nitrogen
NRE	-	Nutrient removal efficiency

NUR	-	Nutrient uptake rate
P	-	Phosphate
PES	-	Provasoli Enriched Seawater (PES)
RAS	-	Recirculation aquaculture system
S	-	Substrate concentration
SGR	-	specific growth rate
SRP	-	Soluble reactive phosphorus
sp.	-	species
SSW	-	Sterilized seawater
TAN	-	Total ammoniacal nitrogen
TN	-	Total nitrogen
TP	-	Total phosphorus
TSS	-	Total suspended solid
<i>U. lactuca</i>	-	<i>Ulva lactuca</i>
UTM	-	Universiti Teknologi Malaysia
WQI	-	Water quality index
Wt.	-	Weight

**LIST OF SYMBOLS**

$C_0$	-	Substrate concentration at initial period
$C_t$	-	Substrate concentration at final period
$K_s$	-	Half saturation constant
$^{\circ}\text{C}$	-	degree celcius
$t$	-	time
$V$	-	Nutrient uptake rate
$W_0$	-	Initial weight of macroalgae
$W_t$	-	Final weight of macroalgae
SA	-	surface area
Y	-	yield

**LIST OF APPENDICES**

<b>APPENDIX</b>	<b>TITLE</b>	<b>PAGE</b>
A	Details of calculations	270
B	AAS, pH meter and multi-parameter water quality meter operational guidelines	273
C	Proximate analysis method (AOAC Manual)	283
D	Statistical analysis with Minitab	290
E	Collection of publications	300

## **CHAPTER 1**

### **INTRODUCTION**

#### **1.1 Background of the Study**

Intensive shrimp production expands rapidly in most south-east Asian countries as it serves major protein source for human consumption (Naylor *et al.*, 2000; Anh *et al.*, 2010). In Malaysia, shrimp cultivation is supported actively due to its relatively clean and abundant supply of water with extensive coastline. Malaysia is one of the top ten shrimp producers in the world with production capacity of 2.8% in total world production of shrimp. In 2003, 70% of total aquaculture production in the country was generated by brackish water aquaculture species (FAO, 2012a) where shrimp production alone stood at 30,000 tonnes (Hashim and Kathamuthu, 2005) and in 2012 this figure has increased to 65,000 tonnes for meeting shrimp demand for local consumption and export targets (DoFM, 2012a). Furthermore, Asian countries are well known as shrimp producers and exporters since temperate regions have unfavourable environmental condition to culture similar shrimp species.

The rapid expansion of aquaculture industry has contributed to the degradation of coastal environment, due to the discharged of aquaculture wastewater directly to the sea or waterways prior to treatment. Thakur and Lin (2003) had revealed on nutrient budget in a closed shrimp culture system that shrimp could assimilate only 23-31% nitrogen and 10-13% phosphorus of the total inputs. Shrimp wastewater has high load of excess nutrient and organic matter, in the form of excretory waste products and

excess leftover feed materials. Consequently, the high amount of nutrient and organic loads will cause eutrophication and occurrence of red tide which affects marine organism and degrades the sustainability of coastal environment (Buschmann *et al.*, 1996; Thakur and Lin, 2003; Fei, 2004). Large amount of nitrogen discharge into rivers are notable in Cambodia, Malaysia, Thailand, and Vietnam with a value of 300,000 tonnes of nitrogen per year from various source (Todd *et al.*, 2010). This figure shows a large amount of nitrogen discharged without concern on the impact to the environment.

Integration of macroalgae cultivation in aquaculture wastewater is considered an ecologically sound aquaculture practice since macroalgae are known as potential nutrient strippers to assimilate excess nutrient from wastewater (Abreu *et al.*, 2011a; Kang *et al.*, 2011). Macroalgae biomass also contributes to the economic development with valuable by-products such as phycocolloid agar (Lüning and Pang, 2003; Neori *et al.*, 2003), human food (Norziah and Ching, 2000; Besada *et al.*, 2009), animal feed (Fleurence, 1999) and biofuel (Bastianoni *et al.*, 2008; Migliore *et al.*, 2012). The macroalgae genera most common in mariculture biofiltration are *Ulva* and *Gracilaria* which have been utilised in a few studies integrated with aquaculture animals in open sea, in ponds and as well as in tank culture systems (Buschmann *et al.*, 1996; Troell *et al.*, 1997; Fei, 2004; Hernandez *et al.*, 2005; Msuya *et al.*, 2006; Marinho-Soriano *et al.*, 2011).

Buschmann *et al.*, (1996) found that in an integrated salmon-*G. Chilensis* tank cultivation system, the macroalgae was capable of removing 95% of ammonium and 32% of orthophosphate. In addition, Troell *et al.*, (1997) stated that a considerable reduction in dissolved ammonium and phosphate can be achieved in an integrated culture of the *Gracilaria chilensis*-salmon, which ultimately reduce the risk of eutrophication. Fei (2004) reported that the *Gracilaria* can solve the eutrophication problem with 6.6% of growth rate. Tank cultivation of *G. caudata* and the filter feeder *Artemia franciscana* had biofiltration capacity for ammonium of 29.4% (Marinho-Soriano *et al.*, 2011). The *U. rotundata*, a type of *Ulva* genera, removes 54% of the total dissolved inorganic nitrogen in 600 L effluents of gilthead seabream in 70 days

(Hernandez *et al.*, 2005). Msuya *et al.*, (2006) found that *U. reticulata* can remove total ammonia nitrogen (TAN) of  $6.5 \text{ g m}^{-2} \text{ day}^{-1}$  with 65% removal efficiency, which is a significant figure.

Based on the findings, this study proposed the integration of macroalgae for treatment of shrimp wastewater as a viable treatment approach to balance the negative impacts to the ecosystem. The hypothesis of this study is selected macroalgae, *U. lactuca* and *G. edulis* are potential biofilters in removing the excess nutrients from shrimp wastewater effectively and can result in higher biomass production of the shrimp and macroalgae. *G. edulis* and *U. lactuca* were selected as biofilters which have been found to grow naturally in canals around the shrimp cultivation pond and at Sg. Melayu estuaries of Johor Bahru, Malaysia. In order to assess biofiltration capacity of both macroalgae, several laboratory and outdoor experiments were conducted to confirm the hypothesis.

### **1.1.1 Preamble of Macroalgae**

Macroalgae or also known as seaweed, are the most suitable candidate for biofiltration because of their high productivity of all plants and economically feasible to be cultured in large scale (Chopin *et al.*, 2001a; Neori *et al.*, 2004). Macroalgae biomass resource has contributed to large industrial products development such as food, soil conditioner, chemical and pharmaceutical products (Troell *et al.*, 1997; Capo *et al.*, 1999). Besides that, the major industrial use for macroalgae is as a source of phycocolloids, such as agar that mostly used as tissue culture media. The increasing demand for hydrocolloid extracts during the past 40 years has led to dramatic growth of the macroalgae industry in few countries (Briggs and Smith, 1993; Dawes, 1995b; Alveal *et al.*, 1997). In Eastern Asian countries, particularly in Japan, China and Korea, macroalgae cultivation has become an integral part of daily diet as protein source for human and animal feed (Bezerra and Marinho-Soriano, 2010). The society




is becoming aware of the benefits and potential of macroalgae and new algal products. An imminent in novel uses of macroalgae are stimulant to encourage more research and development in macroalgae cultivation (Lüning and Pang, 2003). Thus, cultivation of macroalgae has become a prominent industry where it generates income for the country and improved the socio-economic condition.

Besides that, this naturally growing species, has the potential to act as a biofiltration agent where it considerably reduces pollution load of aquaculture effluent and subsequently improve water quality (Buschmann *et al.*, 1996; Troell *et al.*, 1999; Martínez-Aragón *et al.*, 2002; Neori *et al.*, 2004). As a case study, Ann and his colleagues (2010) had discussed on water pollution occurred rapidly from intensive shrimp farming in south-east Vietnam, to meet the demand of food production for human consumption and impact to the environment. In this perspective, integration of macroalgae cultivation in aquaculture wastewater is a biological approach for mitigation of this problem and has been practiced by few researchers either in integrated manner or in an integrated multi-trophic aquaculture (IMTA) system (Jones *et al.*, 2001; Zhou *et al.*, 2006; Xu *et al.*, 2008a).

Macroalgae are important elements in many proposed integrated or multi trophic level mariculture systems as biofilters. Yet, there are only few successfully tested species integrated with shrimp cultivation out there and little information on economic viability as biofilters is available (Neori *et al.*, 2004; Troell *et al.*, 2003) and it should be economically interesting species to be cultivated. Macroalgae biomass relative growth rate increases by incorporating the excess nutrients of shrimp wastewater (particularly C, N and P) with the presence of solar energy. In integrated aquaculture, waste nutrients are considered not a burden but a resource that auxiliary macroalgae growth (Neori *et al.*, 2004).

## 1.2 Problem Statement

The study focused on the production of shrimp wastewater and the effect of its effluent to the environment prior to discharge without any proper treatment. The increased number of farms with high shrimp production rates and high effluent discharge has significantly causes environmental degradation in the shrimp ponds and the adjacent waterways. The typical duration for shrimp cultivation cycle is 100 to 120 days and harvested after this period. After harvesting, shrimp effluent in ponds will be discharged to the adjacent canals, or waterways without any treatment. The effluent is highly polluted and causes deleterious effect. Shrimp wastewater produces large amount of excess nutrient, such as ammonia that is harmful for shrimps and organic matter during the cultivation cycle. Figure 1.1 exhibits the problems associated with shrimp effluents.



**Figure 1.1:** Associated problems of shrimp pond effluent.

Consequently, the amount of nutrients and organic load from the wastewater has to be reduced to ensure better survival and productivity of the shrimp. Typically, shrimp wastewater will be discharged and replenished with fresh seawater as a method of treatment. However, this method of treatment is not environmentally friendly. Thus, the present study proposed on integration of macroalgae, where it could function as biofilters by removing excess nutrients load in the wastewater. This method is known as a more sustainable and ecologically sound aquaculture practices with biological treatment options by using macroalgae to improve the shrimp water quality.

However, another problem encountered when developing the integrated treatment system such as the choice of macroalgae used as biofilters that will significantly affect the shrimp production. The selection of the potential macroalgae species must take into consideration few aspects such as its tolerance to the hypereutrophic condition of the wastewater, high yield and growth rate, has economic values, abundantly found and thrives naturally, and safe for integration with shrimp. Thus, the hypothesis of this study suggests that *U. lactuca* and *G. edulis* are assume to meet these requirements. In addition, this study has also solved the bottleneck problem of design consideration for a sustainable treatment system that is affordable by farmers. Typically, the integration involved between, waste produced species and macroalgae only and very little study is dedicated on combination of few species. Thereby, a second hypothesis was pointed out to develop integrated multi-trophic aquaculture (IMTA) treatment system that involved combination of organic extractive species such as green lipped mussel with the inorganic extractive, macroalgae species. Interestingly, the choices of species selected, the design consideration and semi-closed recirculation IMTA system is the breakthrough for this kind of integrated system. Furthermore, the macroalgae biomass improved by the approach of maximum sustainable yield (MSY) that suggests the optimum biomass for harvest to maintain the biofiltration efficiency of the treatment system.

### 1.3 Objectives of the Study

The aim of this study is to perform modification of an integrated macroalgae and aquaculture treatment system and assess potential of macroalgae as biofilters. Special attention is dedicated on the physicochemical effect and removal performances specifically on the nutrient removal. In addition, organic and nutrient loads were also observed for the enhancement of the treatment system. This can be achieved by the following specific objectives:-

1. To investigate the dynamics of nutrients and organic matter that is produced during shrimp cultivation and characterization of its wastewater constituent for understanding the pollution load that is generated by shrimp cultivation.
2. To assess biofiltration capacity and kinetics of nutrient uptake by *G. edulis* and *U. lactuca* for nutrient removal from shrimp wastewater and the effect on macroalgae and shrimp biomass production.
3. To perform modification of a semi-closed circulation integrated multi-trophic aquaculture (IMTA) system that efficiently removes excess organic matter and nutrients from shrimp wastewater and reuse of the treated water for shrimp cultivation.
4. To perform process optimization of nutrient removal by a maximum sustainable yield (MSY) model approach that predicts the maximum biomass for harvest to maintain the optimum function of the biofiltration system and also biomass production of macroalgae.

#### 1.4 Scope of the Study

This study was conducted using macroalgae species, *Gracilaria edulis* from the brackish water canal located at Fisheries Department of Brackish Water Culture of Gelang Patah, Johor, (PPTAP) and *Ulva lactuca* from the estuaries of Sg. Melayu, Johor, Malaysia. Both macroalgae species possesses different morphology, characteristics and physical properties that follows its taxonomic classification. However, both could act as potential biofilters in treatment of shrimp wastewater and significantly assimilates the dissolved inorganic nutrients.

The eco-physiological properties of the selected macroalgae could adapt and survive to brackish shrimp wastewater condition. In addition, the macroalgae species and green lipped mussels selected were found thrives naturally, for easy accessibility of culture materials. In addition, tiger shrimp, *Penaeus monodon*, of post larvae size was selected to observe its growth throughout shrimp production cycle and physico-chemical characteristics of the wastewater. The biofiltration capacity for nutrient removal and uptake rate investigated by using both macroalgae in several laboratories and outdoor tank-scale experiments. For outdoor tank-scale experiments, two treatment systems were designed- shrimp wastewater recirculation system and semi-closed circulation integrated multi trophic aquaculture (IMTA) system that was built outdoor of Environmental Engineering Laboratory, Universiti Teknologi Malaysia (UTM).

## 1.5 Significance of the Study

Aquaculture wastewater treatment have been extensively reported, particularly, for wastewater derived from varieties of fish cultivation integrated with macroalgae cultivation (Hernandez *et al.*, 2005; Yang *et al.*, 2005b; Zhou *et al.*, 2006; Hayashi *et al.*, 2008; Abreu *et al.*, 2011b; Huo *et al.*, 2012). However, very few research have been conducted on treatment of shrimp wastewater and improvement of its water quality. Application of ecological engineering tools such as IMTA for water pollution control by macroalgae integration is a viable approach for reducing the pollution load caused by shrimp cultivation. Thus, in this study water quality deterioration of shrimp wastewater has been addressed and as a mitigation, application of macroalgae biomass which act as a biofilter has been proposed as a biological treatment options. Besides its potential as biofilters, these macroalgae have considerable economic value and advantageous to be used in IMTA system. In addition, once the biofilter is selected, harvesting rate could be the most important controlling factor in design of a biofiltration system. To date, research on MSY approach to develop a biofiltration system is not given much attention. Thus, the application of MSY concepts was highlighted in this study and contributes to the design of a high efficiency sustainable biofiltration system.

Typically, IMTA system is developed and practiced by a few countries with the aim to produce a more productive, economically profitable and environmentally sound aquaculture system. However, major modification was made where the combination of aquaculture species used were tiger shrimp, green lipped mussel and macroalgae species i.e. *G. edulis* and *U. lactuca*. This is a new configuration which have not been used by others and known as semi-closed circulation integrated multi-trophic aquaculture (IMTA) system. In addition, the design of this system and execution in a semi-closed circulation approach, significantly improves the water quality and the combination of these species are safe for shrimp cultivation. Besides that, frequent water exchange that associated with the possibility of diseases outbreak during shrimp cultivation also improved in this closed circulation IMTA system.

Thereby, this study claims novelty for the design of this integrated treatment system and has been patented. The importance of this study are as follows:-

- I. This study has revealed the amount of wastewater discharged that is produced from shrimp cultivation throughout its life cycle. The fate of nutrients that enter the shrimp tank in the form of feed were also identified.
- II. The study exhibited that the selected macroalgae species can be cultured in shrimp wastewater and produces extra biomass while utilizing nitrogen and phosphorus from the wastewater.
- III. This study provides a systematic treatment design which could be applicable for massive scale of shrimp cultivation and reuse of treated wastewater via the fabrication of a semi-closed circulation integrated multi-trophic aquaculture (IMTA) system.
- IV. This study also provides information to design an integrated treatment system with combination of several treatment units and species successfully with considerable reduction of nutrients and organic load, that results in better yield of macroalgae and shrimp production.
- V. The study provides information regarding MSY approach that can be applied for maximum harvest of macroalgae biomass for optimum biofiltration capacity.
- VI. This study also offers multi-product harvesting at the end of the treatment stage and conversion into useful products where macroalgae biomass could be processed into fertilizer or animal feed, with shrimp and mussel for seafood commercialization purposes.

## 1.6 Thesis Organization

The thesis consist of seven chapters. Chapter 1, introduction of the study was explained, which contains the underlying considerations made in embarking this study. Problem statement of the selected issues that concern of this study were highlighted. Followed by the objectives or motivation for the study to be carried-out, which reflects the solution proposed for the problems proclaimed previously. The scope of the study aimed to confine the study to meet the objectives and satisfies the significance of the study. Lastly, the output of the study would be required to conform the significance of study which is known as the driving force for the overall research that has been reported in this thesis.

Collection of literature reviews presented in Chapter 2. Many books, journals, literature collections, government reports, and proceedings were sought to form the foundation of this chapter. The prospect and challenges of shrimp and macroalgae industry, internationally and nationally were discussed. The significant integration of macroalgae with aquaculture wastewater was described and the efficiency of the existing IMTA system was discussed. The approach of maximum sustainable yield (MSY) is also pointed out.

Chapter 3 recorded the research materials and methodology applied to assist the research work to meet the objectives. Data analysis performed using Minitab software. Chapter 4 presents few laboratory experiments to assess on factors that affects macroalgae growth. Chapter 5 exhibits the potential of macroalgae as biofilters in an outdoor tank shrimp wastewater recirculation system integrated with macroalgae. Eventually, Chapter 6 represents the combined treatment efficiencies in a semi-closed circulation integrated multi-trophic aquaculture (IMTA) system. The conclusions were highlighted in Chapter 7. Moreover, this chapter also covers future research recommendation and exploration extended from this study.



## REFERENCES

- Abreu, M. H. Pereira, R. Buschmann, A. H. Sousa-Pinto, I. and Yarish, C. (2011a). Nitrogen uptake responses of *Gracilaria vermiculophylla* (Ohmi) Papenfuss under combined and single addition of nitrate and ammonium. *Journal of Experimental Marine Biology and Ecology*. 407(2), 190-199.
- Abreu, M. H. Pereira, R. Yarish, C. Buschmann, A. H. and Sousa-Pinto, I. (2011b). IMTA with *Gracilaria vermiculophylla*: Productivity and nutrient removal performance of the seaweed in a land-based pilot scale system. *Aquaculture*. 312(1-4), 77-87.
- Ahemad, S. Ismail, A. and Mohammad, R. (2006). The seaweed industry in Sabah, East Malaysia. *Jati*. 11, 97-107.
- Ale, M. T. Mikkelsen, J. D. and Meyer, A. S. (2011). Differential growth response of *Ulva lactuca* to ammonium and nitrate assimilation. *Journal of Applied Phycology*. 23, 345-351.
- Allan, G. L. and Maguire, G. B. (1992). Effects of pH and salinity on survival, growth and osmoregulation in *Penaeus monodon* Fabricius. *Aquaculture*. 107(1), 33-47.
- Alveal, K. Romo, H. Werlinger, C. and Oliveira, E. C. (1997). Mass cultivation of the agar-producing alga *Gracilaria chilensis* (Rhodophyta) from spores. *Aquaculture*. 148(2-3), 77-83.
- Andersen, R. A. (2005). *Algal Culturing Techniques*. (pp. USA: Elsevier Academic Press.

- Andreas Schuenhoff Leonardo Mata and Santos, R. (2006). The tetrasporophyte of *Asparagopsis armata* as a novel seaweed biofilter. *Aquaculture*. 252, 3–11.
- Andria, J. R. Vergara, J. J. and Perez-Llorens, J. L. (1999a). Biochemical responses and photosynthetic performance of *Gracilaria* sp. (Rhodophyta) from Cadiz, Spain, cultured under different inorganic carbon and nitrogen levels. *European Journal of Phycology*. 34, 497-504.
- Andria, J. R. Vergara, J. J. and Perez-Llorens, J. L. (1999b). Biochemical responses and photosynthetic performance of *Gracilaria* sp. (Rhodophyta) from Cádiz, Spain, cultured under different inorganic carbon and nitrogen levels. *European Journal of Phycology*. 34(05), 497-504.
- Anh, P. T. Kroeze, C. Bush, S. R. and Moi, A. P. J. (2010). Water Pollution Control by Intensive Brackish Shrimp Farming in South-east Vietnam: Causes and Options for Control. *Agricultural Water Management*. 97, 872-882.
- Anon (2003). *Annual Fisheries Statistics Volume 1, 2003*. Malaysia: Department of Fisheries Malaysia. Ministry of Agriculture and Agro-Based Industry.
- AOAC (1990). *Official methods of analysis* 16th ed. Washington, DC: Association of Official Analytical Chemist.
- APHA/ AWWA (2005). Standard Methods for the Examination of Water and Wastewater. In: Eaton, A. D., Clesceri, L. S., Rice, E. W. & Greenberg, A. E. (ed.) (eds.) 21<sup>st</sup> ed. Washington, D.C: American Public Health Association / American Water Works Association / Water Environment Federation.
- Armisen, R. (1995). World-wide use and importance of *Gracilaria*. *Journal of Applied Phycology*. 7(3), 231-243.
- Axelsson, L. M, M. J. and L, F. F. (2000). Utilization of  $\text{HCO}_3^-$  at high pH by the brown macroalga *Laminaria saccharina*. *European Journal of Phycology*. 35, 53-59.

- Bastianoni, S. Coppola, F. Tiezzi, E. Colacevich, A. Borghini, F. and Focardi, S. (2008). Biofuel potential production from the Orbetello lagoon macroalgae: A comparison with sunflower feedstock. *Biomass and Bioenergy*. 32(7), 619-628.
- Berges, J. A. and Franklin, D. J. (2001). Evolution of an artificial seawater medium: Improvements in enriched seawater, artificial water over the last two decades. *Journal Phycology*. 37, 1138-1145.
- Besada, V. Andrade, J. M. Schultze, F. and González, J. J. (2009). Heavy metals in edible seaweeds commercialised for human consumption. *Journal of Marine Systems*. 75(1–2), 305-313.
- Bezerra, A. F. and Marinho-Soriano, E. (2010). Cultivation of the red seaweed *Gracilaria birdiae* (Gracilariales, Rhodophyta) in tropical waters of northeast Brazil. *Biomass and Bioenergy*. 34(12), 1813-1817.
- Bidwell, R. G. S. and McLachlan, J. (1985). Carbon nutrition of seaweeds: photosynthesis, photorespiration and respiration. *Journal of Experimental Marine Biology and Ecology*. 86, 15-46.
- Bjork, M. Haglund, K. Ramazanov, Z. Gareia-Reina, G. and Pederson, M. (1992). Inorganic-carbon assimilation in the green seaweed *Ulva rigida* C.Ag. (Chlorophyta). *Planta*. 187, 152-156.
- Björnsäter, B. R. and Wheeler, P. A. (1990). Effect of nitrogen and phosphorus supply on growth and tissue composition of *Ulva fenestrata* and *Enteromorpha intestinalis* (ulvales, chlorophyta). *Journal of Phycology*. 26(4), 603-611.
- Bower, C. E. and Bidwell, J. P. (1978). Ionization of ammonia in seawater: effects of temperature, pH, and salinity. *Journal of the Fisheries Board of Canada*. 35(7), 1012-1016.
- Boyd, C. and Gautier, D. (2000). Effluent composition and water quality standards. *Global Aquaculture Advocate*. 3(5), 61-66.

- Boyd, C. E. (2003). Guidelines for aquaculture effluent management at the farm-level. *Aquaculture*. 226, 101-112.
- Boyd, C. E. and Musig, Y. (1992). Shrimp pond effluents: observations of the nature of the problem on commercial farms. *Proceedings of the Special Session on Shrimp Farming*. World Aquaculture Society, Baton Rouge, Louisiana. 195-197.
- Braga, A. D. C. and Yoneshigue-Valentin, Y. (1996). Nitrogen and phosphorus uptake by the Brazilian kelp *Laminaria abyssalis* (Phaeophyta) in culture. *Hydrobiologia*. 326/327, 445-450.
- Briggs, M. R. P. and Smith, S. J. F.-. 1993. Macroalgae in Aquaculture: An Overview and their Possible Role in Shrimp Culture. *Marine Biotechnology*. Bangkok, Thailand.
- Bruhn, A. Dahl, J. Nielsen, H. B. Nikolaisen, L. Rasmussen, M. B. Markager, S. Olesen, B. Arias, C. and Jensen, P. D. (2011). Bioenergy potential of *Ulva lactuca*: Biomass yield, methane production and combustion. *Bioresource Technology*. 102(3), 2595-2604.
- Bui, T. D. Luong-Van, J. and Austin, C. M. (2012). Impact of Shrimp Farm Effluent on Water Quality in Coastal Areas of the World Heritage-Listed Ha Long Bay. *American Journal of Environmental Sciences*. 8(2), 104-116.
- Burford, M. A. and Williams, K. C. (2001). The fate of nitrogenous waste from shrimp feeding. *Aquaculture*. 198(1), 79-93.
- Buschmann, A. H. Correa, J. A. Westermeier, R. Hernández-González, M. a. D. C. and Norambuena, R. (2001). Red algal farming in Chile: a review. *Aquaculture*. 194(3), 203-220.

- Buschmann, A. H. Troell, M. Kautsky, N. and Kautsky, L. (1996). Integrated tank cultivation of salmonids and *Gracilaria chilensis* (Gracilariales, Rhodophyta). *Hydrobiologia*. 326-327(1), 75-82.
- Byrne, K. Zuccarello, G. C. West, J. Liao, M.-L. and Kraft, G. T. (2002). Gracilaria species (Gracilariaceae, Rhodophyta) from southeastern Australia, including a new species, *Gracilaria perplexa* sp. nov.: Morphology, molecular relationships and agar content. *Phycological Research*. 50(4), 295-311.
- Cahill, P. L. Hurd, C. L. and Lokman, M. (2010a). Keeping the water clean — Seaweed biofiltration outperforms traditional bacterial biofilms in recirculating aquaculture. *Aquaculture*. 306(1–4), 153-159.
- Cahill, P. L. Lhurd, C. and Lokman, M. (2010b). Keeping the water clean- Seaweed biofiltration outperforms traditional bacterial biofilms in recirculating aquaculture. *Aquaculture*. 306, 153-159.
- Capo, T. Jaramillo, J. Boyd, A. Lapointe, B. and Serafy, J. (1999). Sustained high yields of *Gracilaria* (Rhodophyta) grown in intensive large-scale culture. *Journal of Applied Phycology*. 11(2), 143-147.
- Chen, F. and Johns, M. R. (1991). Effect of C/N ratio and aeration on the fatty acid composition of heterotrophic *Chlorella sorokiniana*. *Journal of Applied Phycology*. 3(3), 203-209.
- Chen, J.-C. and Lei, S.-C. (1990). Toxicity of Ammonia and Nitrite to *Penaeus monodon* Juveniles. *Journal of the World Aquaculture Society*. 21(4), 300-306.
- Chen, J. C. Cheng, S. Y. and Chen, C. T. (1994). Nitrogen excretion and changes of hemocyanin, protein and free amino acid levels in the hemolymph of *Penaeus monodon* exposed to different concentrations of ambient ammonia-N at different salinity levels. *Marine Ecology*. 110, 85-94.

- Chopin, T. Buschmann, A. H. Halling, C. Troell, M. Kautsky, N. Neori, A. Kraemer, G. P. Zertuche-González, J. A. Yarish, C. and Neefus, C. (2001a). Integrating Seaweeds into Marine Aquaculture System: A Key Toward Sustainability. *Journal of Phycology*. 37(6), 975-986.
- Chopin, T. Buschmann, A. H. Halling, C. Troell, M. Kautsky, N. Neori, A. Kraemer, G. P. Zertuche González, J. A. Yarish, C. and Neefus, C. (2001b). Integrating seaweeds into marine aquaculture systems: a key toward sustainability. *Journal of Phycology*. 37(6), 975-986.
- Chopin, T. Buschmann, A. H. Hulling, C. Troell, M. Kautsky, N. Neon, A. Kraemer, G. P. Zertuche-Gonzala, J. A. Yarish, C. and Neefus, C. (2001c). Mini review- Integrating seaweeds into marine aquaculture systems: A key toward sustainability. *Journal of Phycology*. 37, 975-986.
- Chuntapa, B. Powtongsook, S. and Menasveta, P. (2003). Water quality control using *Spirulina platensis* in shrimp culture tanks. *Aquaculture*. 220(1-4), 355-366.
- Ciba (2003). *Shrimp culture: feeds and feed management*. India: Indian Council of Agricultural Research.
- Copertino, M. D. Tormena, T. and Seeliger, U. (2009). Biofiltering efficiency, uptake and assimilation rates of *Ulva clathrata* (Roth) J. Agardh (Clorophyceae) cultivated in shrimp aquaculture waste water. *Journal of Applied Phycology*. 21(1), 31-45.
- Costanzo, S. D. O'donohue, M. J. and Dennison, W. C. (2000a). *Gracilaria edulis* (Rhodophyta) as a biological indicator of pulsed nutrients in oligotrophic waters. *Journal of Phycology*. 36(4), 680-685.
- Costanzo, S. D. O'donohue, M. J. and Dennison, W. C. (2000b). *Gracilaria edulis* (rhodophyta) as a biological indicator of pulsed nutrients in oligotrophic waters. *Journal of Phycology*. 36, 680-685.

- Crab, R. Avnimelech, Y. Defoirdt, T. Bossier, P. and Verstraete, W. (2007). Nitrogen removal techniques in aquaculture for a sustainable production. *Aquaculture*. 270(1–4), 1-14.
- Critchley, A. T. and Ohno, M. (1998). *Seaweed Resources of the World*. Yokosuka, Japan: Japan Internatioanl Cooperation Agency.
- Cruz-Suárez, L. E. León, A. Peña-Rodríguez, A. Gustavo Rodríguez-Peña C Moll, B. and Ricque-Marie, D. (2010). Shrimp/*Ulva* co-culture: A sustainable alternative to diminish the need for artificial feed and improve shrimp quality. *Aquaculture*. 301, 64-68.
- Cuomo, V. Merrill, J. Palomba, I. and Perretti, A. (1993). Systematic collection of *Ulva* and mariculture of *Porphyra*: biotechnology against eutrophication in the Venice Lagoon. *International journal of environmental studies*. 43(2-3), 141-149.
- D'elia, C. F. and Deboer, J. A. (1978). Nutritional studies of two red algae. II. Kinetics of ammonium and nitrate uptake. *Journal of Phycology*. 14(3), 266-272.
- Dawes, C. (1995a). Suspended cultivation of *Gracilaria* in the sea. *Journal of Applied Phycology*. 7(3), 303-313.
- Dawes, C. J. Orduña-Rojas, J. and Robledo, D. (1999). Response of the tropical red seaweed *Gracilaria cornea* to temperature, salinity and irradiance. *Journal of Applied Phycology*. 10(5), 419-425.
- Dawes, C. P. (1995b). Suspended cultivation of *Gracilaria* in the sea. *Journal of Applied Phycology*. 7(3), 303-313.
- De Graaf, G. and Xuan, T. (1998). Extensive shrimp farming, mangrove clearance and marine fisheries in the southern provinces of Vietnam. *Mangroves and Salt Marshes*. 2(3), 159-166.

- Debusk, T. Blakeslee, M. and Ryther, J. (1986). Studies on the Outdoor Cultivation of *Ulva lactuca* L. *Botanica Marina*. 29(5), 381-386.
- Dhargalkar, V. and Pereira, N. (2005). Seaweed: promising plant of the millennium. *Science and culture*. 71(3-4), 60-66p.
- Dhargalkar, V. K. and Verlecar, X. N. (2009). Southern Ocean seaweeds: A resource for exploration in food and drugs. *Aquaculture*. 287(3-4), 229-242.
- DoFM (1996). *Annual Fisheries Statistics*. Kuala Lumpur: Ministry of Agriculture Malaysia.
- DoFM (2012a). *Annual Fisheries Statistics*. Kuala Lumpur, Malaysia: Department of Fisheries Malaysia, Ministry of Agriculture Malaysia.
- DoFM (2012b). *Perangkaan Perikanan Tahunan 2012-Awalan*. Putrajaya, Malaysia: Jabatan Perikanan Malaysia.
- Ebeling, J. M. Timmons, M. B. and Bisogni, J. (2006). Engineering analysis of the stoichiometry of photoautotrophic, autotrophic, and heterotrophic removal of ammonia–nitrogen in aquaculture systems. *Aquaculture*. 257(1), 346-358.
- El-Naggar, A. H. Osman, M. E. El-Sheekh, M. M. and Gheda, S. F. (2005). Influence of the aqueous extracts of *Ulva lactuca* and *Chlorella kessleri* on growth and yield of *Vicia faba*. *Algological Studies*. 116(1), 213-229.
- Emerson, K. Russo, R. C. Lund, R. E. and Thurston, R. V. (1975). Aqueous ammonia equilibrium calculations: effect of pH and temperature. *Journal of the Fisheries Board of Canada*. 32(12), 2379-2383.
- Erler, D. Pollard, P. Duncan, P. and Knibb, W. (2004). Treatment of shrimp farm effluent with omnivorous finfish and artificial substrates. *Aquaculture Research*. 35(9), 816-827.



- Faazaz, A. L. (1986). *Preliminary results on the experimental culture of the red seaweed Gracilaria spp. in Malaysia*. Malaysia: Department of Fisheries, Ministry of Agriculture.
- Fábregas, J. Vázquez, V. Cabezas, B. and Otero, A. (1993). Tris not only controls the pH in microalgal cultures, but also feeds bacteria. *Journal of Applied Phycology*. 5(5), 543-545.
- FAO (1989). *Some advances attained in shrimp farming research and management practices: Insights to future prospects for expansion of production*. Manila, Philippines: Food and Agriculture Organization of the United Nations.
- FAO (1996). *Aquaculture Production Statistics 1985–1994*. Rome: Food and Agriculture Organization of the United Nations.
- FAO (2010). *National Aquaculture Sector Overview*. Malaysia: Fisheries and Aquaculture Department of Malaysia.
- FAO (2012a). *Fishstat Plus, Fisheries production database, software (in CD)*. Malaysia: Fisheries and Aquaculture Department of Malaysia.
- FAO (2012b). *The state of world fisheries and aquaculture*. Rome: Food and Agriculture Organization of the United Nations.
- Fei, X. (2004). Solving the coastal eutrophication problem by large scale seaweed cultivation. *Hydrobiologia*. 512(1), 145-151.
- Fleurence, J. (1999). Seaweed proteins: biochemical, nutritional aspects and potential uses. *Trends in Food Science & Technology*. 10(1), 25-28.
- Friedlander, M. and Levy, I. (1995). Cultivation of Gracilaria in outdoor tanks and ponds. *Journal of Applied Phycology*. 7(3), 315-324.

- Fujita, R. M. (1985). The role of nitrogen status in regulating transient ammonium uptake and nitrogen storage by macroalgae. *Journal of Experimental Marine Biology and Ecology*. 92(2-3), 283-301.
- Funge-Smith, S. J. and Briggs, M. R. P. (1998). Nutrient budgets in intensive shrimp ponds: implications for sustainability. *Aquaculture*. 164, 117-133.
- Gao, K. Aruga, Y. Asada, K. Ishihara, T. Akano, T. and Kiyohara, M. (1991). Enhanced growth of the red alga *Porphyra yezoensis* Ueda in high CO<sub>2</sub> concentrations. *Journal of Applied Phycology*. 3, 355-362.
- Gao, K. Aruga, Y. Asada, K. and Kiyohara, M. (1993). Influence of enhanced CO<sub>2</sub> on growth and photosynthesis of the red algae *Gracilaria sp.* and *G. chilensis*. *Journal of Applied Phycology*. 5, 563-571.
- Gao, K. and McKinley, K. R. (1994). Use of macroalgae for marine biomass production and CO<sub>2</sub> remediation: a review. *Journal of Applied Phycology*. 6, 45-60.
- Gao, Q.-F. Xu, W.-Z. Liu, X.-S. Cheung, S. G. and Shin, P. K. S. (2008). Seasonal changes in C, N and P budgets of green-lipped mussels *Perna viridis* and removal of nutrients from fish farming in Hong Kong. *Marine Ecology Progress Series*. 353, 137-146.
- Garcia, S. Sparre, P. and Csirke, J. (1989). Estimating surplus production and maximum sustainable yield from biomass data when catch and effort time series are not available. *Fisheries Research*. 8(1), 13-23.
- Gerung, S. G. Ohno, M. and Yamamoto, H. (1999). Growth rates and agar properties on some species of *Gracilaria Grev.* (Rhodophyta, Gigartinales) from Manado, Indonesia. *Buletin of marine sciences and fisheries, Kochi University*. 19, 9-14.
- Gil, M. N. Torres, A. I. and Esteves, J. L. (2005). Uptake of sewage derived nitrogen by *Ulva rigida* (Chlorophyceae) in Bahía Nueva (Golfo Nuevo, Patagonia, Argentina). *Hydrobiologia*. 532, 39-43.

- Gross, A. Boyd, C. E. and Wood, C. W. (2000). Nitrogen transformations and balance in channel catfish ponds. *Aquacultural Engineering*. 24, 1-14.
- Guiry, M. D. and Guiry, G. M. 2010. AlgaeBase. National University of Ireland, Galway: World-wide electronic publication.
- Haglund, K. and Pedersén, M. (1993). Outdoor pond cultivation of the subtropical marine red alga *Gracilaria tenuistipitata* in brackish water in Sweden. Growth, nutrient uptake, co-cultivation with rainbow trout and epiphyte control. *Journal of Applied Phycology*. 5(3), 271-284.
- Hanisak, M. D. 1983. The nitrogen relationships of marine macroalgae. . In Carpenter, E. J. & Capone, D. G. (Eds.), *Nitrogen in the marine environment* New York: Academic Press.
- Hanisak, M. D. (1990a). The use of *Gracilaria tikvahiae* (Gracilariales, Rhodophyta) as a model system to understand the nitrogen nutrition of cultured seaweeds. *Hydrobiologia*. 204(1), 79-87.
- Hanisak, M. D. (1990b). The use of *Gracilaria tikvahiae* (Gracilariales, Rhodophyta) as a model system to understand the nitrogen nutrition of cultured seaweeds. *Hydrobiologia*. 204-205(1), 79-87.
- Hashim, M. B. and Kathamuthu, S. (2005). *Shrimp farming in Malaysia: Regional Technical Consultation on the Aquaculture of P. vannamei and Other Exotic Shrimps in Southeast Asia*. Manila, Philippines: SEAFDEC Aquaculture Department.
- Hayashi, L. Yokoya, N. S. Ostini, S. Pereira, R. T. L. Braga, E. S. and C.Oliveira, E. (2008). Nutrients removed by *Kappaphycus alvarezii* (Rhodophyta, Solieriaceae) in integrated cultivation with fishes in re-circulating water. *Aquaculture*. 277, 185-191.

- He, P. Xu, S. Zhang, H. Wen, S. Dai, Y. Lin, S. and Yarish, C. (2008). Bioremediation efficiency in the removal of dissolved inorganic nutrients by the red seaweed, *Porphyra yezoensis*, cultivated in the open sea. *Water Research*. 42(4-5), 1281-1289.
- Healey, F. (1980). Slope of the Monod equation as an indicator of advantage in nutrient competition. *Microbial Ecology*. 5(4), 281-286.
- Hein, M. Pedersen, M. F. and Sandjensen, K. (1995). Size-dependent nitrogen uptake in micro- and macroalgae. *Marine Ecology Progress Series*. 118(1-3), 247-253.
- Hellblom, F. Beer, S. Björk, M. and Axelsson, L. (2001). A buffer sensitive inorganic carbon utilisation system in *Zostera marina*. *Aquatic Botany*. 69 55–62.
- Hernandez, I. Fernandez-Engo, M. A. Perez-Llorens, J. L. and Vergara, J. J. (2005). Integrated outdoor culture of two estuarine macroalgae as biofilters for dissolved nutrients from *Sparus aurata* waste waters. *Journal of Applied Phycology*. 17(6), 557-567.
- Hiraoka, M. and Oka, N. (2008). Tank cultivation of & *Ulva prolifera* in deep seawater using a new “germling cluster” method. *Journal of Applied Phycology*. 20(1), 97-102.
- Ho, Y. (1990a). *Ulva lactuca* as bioindicator of metal contamination in intertidal waters in Hong Kong. *Hydrobiologia*. 203(1-2), 73-81.
- Ho, Y. B. (1990b). *Ulva lactuca* as bioindicator of metal contamination in intertidal waters in Hong Kong. *Hydrobiologia*. 203(1-2), 73-81.
- Huo, Y. Wu, H. Chai, Z. Xu, S. Han, F. Dong, L. and He, P. (2012). Bioremediation efficiency of *Gracilaria verrucosa* for an integrated multi-trophic aquaculture system with *Pseudosciaena crocea* in Xiangshan harbor, China. *Aquaculture*. 326, 99–105.
- Huo, Y. Z. Xu, S. N. Wang, Y. Y. Zhang, J. H. Zhang, Y. J. Wu, W. N. Chen, Y. Q. and He, P. M. (2011). Bioremediation efficiencies of *Gracilaria verrucosa* cultivated in

an enclosed sea area of Hangzhou Bay, China. *Journal of Applied Phycology*. 23, 173-182.

Jackson, C. Presto, N. Thompson, P. J. and Burford, M. (2003a). Nitrogen budget and effluent nitrogen components at an intensive shrimp farm. *Aquaculture*. 218, 397-411.

Jackson, C. J. Preston, N. Burford, M. A. and Thompson, P. J. (2003b). Managing the development of sustainable shrimp farming in Australia: the role of sedimentation ponds in treatment of farm discharge water. *Aquaculture*. 226(1-4), 23-34.

Jackson, C. J. Preston, N. Burford, M. A. and Thompson, P. J. (2003c). Managing the development of sustainable shrimp farming in Australia: the role of sedimentation ponds in treatment of farm discharge water. *Aquaculture*. 226, 23-34.

James, C. S. (1996). *Analytical Chemistry of Foods*. (pp. New York: Chapman and Hall.

Jayasankar, R. and Ramamoorthy, N. (1997). Propagation of *Gracilaria edulis* (Gmelin) Silva by Reproductive Method. *Indian Journal of Fisheries*. 44(4), 353-360.

Jayasankar, R. and Varghese, S. (2002). Short communication: Cultivation of marine red alga *Gracilaria edulis* (Gigartinales, Rhodophyta) from spores *Indian Journal of Marine Sciences*. 31(1), 75-77.

Jiménez Del Rio, M. Ramazanov, Z. and Garcia-Reina, G. (1994). Optimization of yield and biofiltering efficiencies of *Ulva rigida* C. cultivated with *Sparus aurata* L. waste waters. *Scientia Marina*. 58(4), 329-335.

Jiménez Del Rio, M. Ramazanov, Z. and García Reina, G. (1995). Effect of nitrogen supply on photosynthesis and carbonic anhydrase activity in the green seaweed *Ulva rigida* (Chlorophyta). *Marine Biology*. 123(4), 687-691.

Jones, A. B. Dennison, W. C. and Preston, N. P. (2001). Integrated treatment of shrimp effluent by sedimentation, oyster filtration and macroalgal absorption: a laboratory scale study. *Aquaculture*. 193(1-2), 155-178.

- Jones, A. B. Dennison, W. C. and Stewart, G. R. (1996). Macroalgal responses to nitrogen source and availability: Amino acid metabolic profiling as a bioindicator using *Gracilaria edulis* (rhodophyta). *Journal of Phycology*. 32(5), 757-766.
- Jones, A. B. and Preston, N. P. (1999). Oyster filtration of shrimp farm effluent, the effects on water quality. *Aquaculture Research*. 30, 51-57.
- Jones, A. B. Preston, N. P. and Dennison, W. C. (2002). The efficiency and condition of oyster and macroalgae used as biological filters of shrimp pond effluent. *Aquaculture Research*. 33, 1-19.
- Kain, J. and Destombe, C. (1995). A review of the life history, reproduction and phenology of *Gracilaria*. *Journal of Applied Phycology*. 7(3), 269-281.
- Kakita, H. and Kamishima, H. (2006). Effects of environmental factors and metal ions on growth of the red alga *Gracilaria chorda* Holmes (Gracilariales, Rhodophyta). *Journal of Applied Phycology*. 18, 469-474.
- Kaliaperumal, N. and Ramalingam, J. (2005). Effect of different fertilizers on the growth of *Gracilaria edulis* (Gmelin) silva in onshore cultivation. *Indian Hydrobiology*. 7, 63-67.
- Kalita, T. L. and Titlyanov, E. A. (2011). The effect of temperature on infradian rhythms of reproduction in *Ulva fenestrata* postels et ruprecht, 1840 (Chlorophyta: Ulvales). *Russian Journal of Marine Biology*. 37(1), 52-61.
- Kalkman, L. Rajendran, I. and Angell, C. L. 1991. Bay of Bengal Programme: Small-Scale Fisherfolk Communities. *Seaweed (Gracilaria edulis) Farming in Vedalai and Chinnapalam, India*. Madras, India: Fisheries Department.
- Kang, Y. H. Park, S. R. and Chung, I. K. (2011). Biofiltration efficiency and biochemical composition of three seaweed species cultivated in a fish-seaweed integrated culture. *Algae*. 26(1), 97-108.

- Kaur, C. R. and Margaret, A. G. H. 2009. Seminar on Developing the Seaweed Aquaculture Sector in Malaysia (Full Report).
- Kautsky, N. Ro`Nnb`Ack, P. Tedengren, M. and Troell, M. (2000). Ecosystem perspectives on management of disease in shrimp pond farming. *Aquaculture*. 191, 145–161.
- Khan, S. I. and Satam, S. (2003). Seaweed mariculture: scope and potential in India. *Aquac Asia*. 8, 26-29.
- Kim, S. Park, S. Kang, Y. Kim, G.-Y. Lee, K.-S. Lee, H. Won, N.-I. and Kil, H.-J. (2013). Usefulness of tissue nitrogen content and macroalgal community structure as indicators of water eutrophication. *Journal of Applied Phycology*, 1-10.
- Kinne, P. N. Samocha, T. M. Jones, E. R. and Browdy, C. L. (2001). Characterization of Intensive Shrimp Pond Effluent and Preliminary Studies on Biofiltration. *North American Journal of Aquaculture*. 63, 25-33.
- Kongkeo, H. (1997). Comparison of intensive shrimp farming systems in Indonesia, Philippines, Taiwan and Thailand. *Aquaculture Research*. 28(10), 789-796.
- Körner, S. Das, S. K. Veenstra, S. and Vermaat, J. E. (2001). The effect of pH variation at the ammonium/ammonia equilibrium in wastewater and its toxicity to *Lemna gibba*. *Aquatic Botany*. 71, 71-78.
- Kuhn, D. D. Angier, M. W. Barbour, S. L. Smith, S. A. and Flick Jr, G. J. (2013). Culture feasibility of eastern oysters (*Crassostrea virginica*) in zero-water exchange recirculating aquaculture systems using synthetically derived seawater and live feeds. *Aquacultural Engineering*. 54(0), 45-48.
- Lahaye, M. (1991). Marine algae as sources of fibres: determination of soluble and insoluble dietary fibre contents in some ‘sea vegetables’. *Journal of the Science of Food and Agriculture*. 54(4), 587-594.

- Lahaye, M. and Jegou, D. (1993). Chemical and physical-chemical characteristics of dietary fibres from *Ulva lactuca* (L.) Thuret and *Enteromorpha compressa* (L.) Grev. *Journal of Applied Phycology*. 5(2), 195-200.
- Lapointe, B. Tenore, K. and Dawes, C. (1984). Interactions between light and temperature on the physiological ecology of *Gracilaria tikvahiae* (Gigartinales: Rhodophyta). *Marine Biology*. 80(2), 161-170.
- Lapointe, B. E. and Duke, C. S. (1984). Biochemical strategies for growth of *Gracilaria tikvahiae* (rhodophyta) in relation to light intensity and nitrogen availability. *Journal of Phycology*. 20(4), 488-495.
- Lapointe, B. E. and Tenore, K. R. (1981). Experimental outdoor studies with *Ulva fasciata* Delile. I. Interaction of light and nitrogen on nutrient uptake, growth, and biochemical composition. *Journal of Experimental Marine Biology and Ecology*. 53(2-3), 135-152.
- Lee, C. S. and Ang, P. (1991). A simple model for seaweed growth and optimal harvesting strategy. *Ecological Modelling*. 55(1-2), 67-74.
- Legović, T. Klanjšček, J. and Geček, S. (2010). Maximum sustainable yield and species extinction in ecosystems. *Ecological Modelling*. 221(12), 1569-1574.
- Lin, Y.-C. and Chen, J.-C. (2000). Acute toxicity of ammonia on *Litopenaeus vannamei* Boone juveniles at different salinity levels. *Journal of Experimental Marine Biology and Ecology*. 259, 109-119.
- Lin, Y.-F. Jing, S.-R. and Lee, D.-Y. (2003a). The potential use of constructed wetlands in a recirculating aquaculture system for shrimp culture. *Environmental Pollution*. 123(1), 107-113.



- Lin, Y.-F. Jing, S.-R. and Lee, D.-Y. (2003b). The potential use of constructed wetlands in a recirculating aquaculture system for shrimp culture. *Environmental Pollution*. 123, 107-113.
- Lindsey Zemke-White, W. and Ohno, M. (1999). World seaweed utilisation: An end-of-century summary. *Journal of Applied Phycology*. 11(4), 369-376.
- Lobban, C. S. and Harrison, P. J. (1994). *Seaweed Ecology and Physiology*. (pp. 37-39). Cambridge UK: Cambridge University Press.
- Lourenco, S. O. Barbarino, E. Nascimento, A. Freitas, J. N. P. and Diniz, G. S. (2006). Tissue nitrogen and phosphorus in seaweeds in a tropical eutrophic environment: What a long-term study tells us. *Journal of Applied Phycology*. 18, 389-398.
- Lüning, K. and Pang, S. (2003). Mass cultivation of seaweeds: current aspects and approaches. *Journal of Applied Phycology*. 15(2), 115-119.
- Luo, M. B. Liu, F. and Xu, Z. L. (2012). Growth and nutrient uptake capacity of two co-occurring species, *Ulva prolifera* and *Ulva linza*. *Aquatic Botany*. 100(0), 18-24.
- Macdonald, B. A. Robinson, S. M. C. and Barrington, K. A. (2011). Feeding activity of mussels (*Mytilus edulis*) held in the field at an integrated multi-trophic aquaculture (IMTA) site (*Salmo salar*) and exposed to fish food in the laboratory. *Aquaculture*. 314, 244-251.
- Mao, Y. Yang, H. Zhou, Y. Ye, N. and Fang, J. (2009). Potential of the seaweed *Gracilaria lemaneiformis* for integrated multi-trophic aquaculture with scallop *Chlamys farreri* in North China. *Journal of Applied Phycology*. 21, 649-656.
- Marinho-Soriano, E. Azevedo, C. a. A. Trigueiro, T. G. Pereira, D. C. Carneiro, M. a. A. and Camara, M. R. (2011). Bioremediation of aquaculture wastewater using macroalgae and *Artemia*. *International Biodeterioration & Biodegradation*. 65(1), 253-257.

- Marinho-Soriano, E. Fonseca, P. C. Carneiro, M. a. A. and Moreira, W. S. C. (2006). Seasonal variation in the chemical composition of two tropical seaweeds. *Bioresource Technology*. 97, 2402-2406.
- Marinho-Soriano, E. Morales, C. and Moreira, W. S. C. (2002). Cultivation of *Gracilaria* (Rhodophyta) in shrimp pond effluents in Brazil. *Aquaculture Research*. 33, 1081-1086.
- Marinho-Soriano, E. Nunes, S. O. Carneiro, M. a. A. and Pereira, D. C. (2009a). Nutrients' removal from aquaculture wastewater using the macroalgae *Gracilaria birdiae*. *Biomass and Bioenergy*. 33(2), 327-331.
- Marinho-Soriano, E. Panucci, R. A. Carneiro, M. a. A. and Pereira, D. C. (2009b). Evaluation of *Gracilaria caudata* J. Agardh for bioremediation of nutrients from shrimp farming wastewater. *Bioresource Technology*. 100(24), 6192-6198.
- Marinho-Soriano, E. Panucci, R. A. Carneiro, M. a. A. and Pereira, D. C. (2009c). Evaluation of *Gracilaria caudata* J. Agardh for bioremediation of nutrients from shrimp farming wastewater. *Bioresource Technology*. 100, 6192-6198.
- Martínez-Aragón, J. F. Hernández, I. Pérez-Lloréns, J. L. Vázquez, R. and Vergara, J. J. (2002). Biofiltering efficiency in removal of dissolved nutrients by three species of estuarine macroalgae cultivated with sea bass (*Dicentrarchus labrax*) waste waters 1. Phosphate. *Journal of Applied Phycology*. 14(5), 365-374.
- Mata, L. Schuenhoff, A. and Santos, R. (2010). A direct comparison of the performance of the seaweed biofilters, *Asparagopsis armata* and *Ulva rigida*. *Journal of Applied Phycology*. 22, 639-644.
- Matsunaga, K. Suzuki, Y. Kuma, K. and Kudo, I. (1994). Diffusion of Fe(II) from an iron propagation cage and its effect on tissue iron and pigments of macroalgae on the cage. *Journal of Applied Phycology*. 6(4), 397-403.

- Mcintosh, D. and Fitzsimmons, K. (2003). Characterization of effluent from an inland, low salinity shrimp farm: what contribution could this water make if used for irrigation. *Aquaculture Engineering*. 27 147-156.
- Menasveta, P. Panritdam, T. Sihanonth, P. Powtongsook, S. Chuntapa, B. and Lee, P. (2001). Design and function of a closed, recirculating seawater system with denitrification for the culture of black tiger shrimp broodstock. *Aquacultural Engineering*. 25, 35-49.
- Mercado, J. M. F. Javier, L. G. Niell, F. X. and Figueroa, F. L. (1999). Effects of different levels of CO<sub>2</sub> on photosynthesis and cell components of the red alga *Porphyra leucosticta*. *Journal of Applied Phycology*. 11, 455-461.
- Migliore, G. Alisi, C. Sprocati, A. R. Massi, E. Ciccoli, R. Lenzi, M. Wang, A. and Cremisini, C. (2012). Anaerobic digestion of macroalgal biomass and sediments sourced from the Orbetello lagoon, Italy. *Biomass and Bioenergy*. 42(0), 69-77.
- Miller, P. Flaherty, M. and Szuster, B. (1999). Inland shrimp farming in Thailand. *Aquaculture Asia. Bangkok*. 4(1), 27-32.
- Miranda, F. R. Lima, R. N. Crisostomo, L. A. and Santana, M. G. S. (2008). Reuse of inland low-salinity shrimp farm effluent for melon irrigation. *Aquaculture Engineering*. 39, 1-5.
- Mohamed, M. I. H. (1991). National management of Malaysian fisheries. *Marine Policy*. 15(1), 2-14.
- Motoh, H. (1985). Biology and ecology of *Penaeus monodon*. *Proceedings of the First International Conference on the Culture of Penaeid Prawns/Shrimps*. Iloilo City, Philippines. 27-36.
- Msuya, F. and Neori, A. (2008). Effect of water aeration and nutrient load level on biomass yield, N uptake and protein content of the seaweed *Ulva lactuca* cultured in seawater tanks. *Journal of Applied Phycology*. 20(6), 1021-1031.

- Msuya, F. E. Kyewalyanga, M. S. and Salum, D. (2006). The performance of the seaweed *Ulva reticulata* as a biofilter in a low-tech, low-cost, gravity generated water flow regime in Zanzibar, Tanzania. *Aquaculture*. 254(1-4), 284-292.
- Msuya, F. E. and Neori, A. (2002). *Ulva reticulata* and *Gracilaria crassa*: Macroalgae That Can Biofilter Effluent from Tidal Fishponds in Tanzania. *Western Indian Ocean Journal Marine Science*. 1(2), 117-126.
- Muangkeow, B. Ikejima, K. Powtongsook, S. and Yi, Y. (2007). Effects of white shrimp, *Litopenaeus vannamei* (Boone), and Nile tilapia, *Oreochromis niloticus* L., stocking density on growth, nutrient conversion rate and economic return in integrated closed recirculation system. *Aquaculture*. 269, 363-376.
- Mwandya, A. Mtolera, M. Pratap, H. and Jiddawi, N. (2001). Macroalgae as biofilters of dissolved inorganic nutrients in an integrated mariculture tank system in Zanzibar. *Marine Science Development in Tanzania and Eastern Africa. Proc. 20th Anniversary Conference on Advances in Marine Science in Tanzania*. 159-170.
- Naldi, M. and Viaroli, P. (2002a). Nitrate uptake and storage in the seaweed *Ulva rigida* C. Agardh in relation to nitrate availability and thallus nitrate content in a eutrophic coastal lagoon (Sacca di Goro, Po River Delta, Italy). *Journal of Experimental Marine Biology and Ecology*. 269, 65-83.
- Naldi, M. and Viaroli, P. (2002b). Nitrate uptake and storage in the seaweed *Ulva rigida* C. Agardh in relation to nitrate availability and thallus nitrate content in a eutrophic coastal lagoon (Sacca di Goro, Po River Delta, Italy). *Journal of Experimental Marine Biology and Ecology*. 269(1), 65-83.
- Naldi, M. and Wheeler, P. A. (2002). <sup>15</sup>N measurements of ammonium and nitrate uptake by *Ulva fenestrata* (chlorophyta) and *Gacilaria pacifica* (rhodophyta): Comparison of net nutrient disappearance, release of ammonium and nitrate, and <sup>15</sup>N accumulation in algal tissue. *Journal of Phycology*. 38(1), 135-144.

- Navarro-Angulo, L. and Robledo, D. (1999). Effects of nitrogen source, N:P ratio and N-pulse concentration and frequency on the growth of *Gracilaria cornea* (Gracilariales, Rhodophyta) in culture. *Hydrobiologia*. 398-399, 315–320.
- Naylor, R. L. Goldberg, R. J. Primavera, J. H. Kautsky, N. Beveridge, M. C. Clay, J. Folke, C. Lubchenco, J. Mooney, H. and Troell, M. (2000). Effect of Aquaculture on World Fish Supplies. *Nature*. 405(6790), 1017-1024.
- Nelson, S. G. Glenn, E. P. Conn, J. Moore, D. Walsh, T. and Akutagawa, M. (2001a). Cultivation of *Gracilaria parvispora* (Rhodophyta) in shrimp-farm effluent ditches and floating cages in Hawaii: a two-phase polyculture system. *Aquaculture*. 193(3), 239-248.
- Nelson, S. G. Glenn, E. P. Conn, J. Moore, D. Walsh, T. and Akutagawa, M. (2001b). Cultivation of *Gracilaria parvispora* (Rhodophyta) in shrimp-farm effluent ditches and floating cages in Hawaii: a two-phase polyculture system. *Aquaculture*. 193(3), 239-248.
- Neori, A. Chopin, T. Troell, M. Buschmann, A. H. Kraemer, G. P. Halling, C. Shpigel, M. and Yarish, C. (2004). Integrated aquaculture: rationale, evolution and state of the art emphasizing seaweed biofiltration in modern mariculture. *Aquaculture*. 231(1-4), 361-391.
- Neori, A. Cohen, I. and Gordin, H. (1991). *Ulva lactuca* Biofilters for Marine Fishpond Effluents. II. Growth Rate, Yield and C:N Ratio. *Botanica Marina*. 34(6), 483-489.
- Neori, A. Msuya, F. E. Shauli, L. Schuenhoff, A. Kopel, F. and Shpigel, M. (2003). A novel three-stage seaweed (*Ulva lactuca*) biofilter design for integrated mariculture. *Journal of Applied Phycology*. 15(6), 543-553.
- Neori, A. Shpigel, M. and Ben-Ezra, D. (2000). A sustainable integrated system for culture of fish, seaweed and abalone. *Aquaculture*. 186, 279-291.

- Nielsen, S. and Jensen, K. S. (1990). Allometric scaling of maximal photosynthetic growth rate to surface/volume ratio. *Limnology and Oceanography*. 35(1), 177-181.
- Norziah, M. H. and Ching, C. Y. (2000). Nutritional composition of edible seaweed *Gracilaria changgi*. *Food Chemistry*. 68(1), 69-76.
- Ortiz, J. Romero, N. Robert, P. Araya, J. Lopez-Hernandez, J. Bozzo, C. Navarrete, E. Osorio, A. and Rios, A. (2006). Dietary fiber, amino acid, fatty acid and tocopherol contents of the edible seaweeds *Ulva lactuca* and *Durvillaea antarctica*. *Food Chemistry*. 99(1), 98-104.
- Páez-Osuna, F. (2001). The Environmental Impact of Shrimp Aquaculture: Causes, Effects, and Mitigating Alternatives. *Environmental Management*. 28(1), 131-140.
- Páez-Osuna, F. Guerrero-Galván, S. R. Ruiz-Fernández, A. C. and Espinoza-Angulo, R. (1997). Fluxes and mass balances of nutrients in a semi-intensive shrimp farm in north-western Mexico. *Marine Pollution Bulletin*. 34(5), 290-297.
- Paniagua-Michel, J. and Garcia, O. (2003). Ex-situ bioremediation of shrimp culture effluent using constructed microbial mats. *Aquacultural Engineering*. 28, 131-139.
- Paula, E. J. D. Erbert, C. and Pereira, R. T. L. (2001). Growth rate of the carrageenophyte *Kappaphycus alvarezii* (Rhodophyta, Gigartinales) in vitro. *Phycological Research*. 49(3), 155-161.
- Peckol, P. and Rivers, J. S. (1995). Physiological responses of the opportunistic macroalgae *Cladophora vagabunda* (L.) van den Hoek and *Gracilaria tikvahiae* (McLachlan) to environmental disturbances associated with eutrophication. *Journal of Experimental Marine Biology and Ecology*. 190(1), 1-16.
- Pedersen, M. F. and Borum, J. (1992). Nitrogen dynamics of eelgrass *Zostera marina* during a late summer period of high growth and low nutrient availability. *Marine Ecology-Progress Series*(80), 65-73.

- Pedersen, M. F. and Borum, J. (1996). Nutrient control of algal growth in estuarine waters. Nutrient limitation and the importance of nitrogen requirements and nitrogen storage among phytoplankton and species of macroalgae. *Marine ecology progress series. Oldendorf*. 142(1), 261-272.
- Pérez-Lloréns, J. L. and Niell, F. X. (1995). Short-term phosphate uptake kinetics in *Zostera noltii* Hornem: a comparison between excised leaves and sediment-rooted plants. *Hydrobiologia*. 297(1), 17-27.
- Phang, S.-M. (1994). Some species of *Gracilaria* from peninsular Malaysia and Singapore. *Taxonomy of Economic Seaweeds*. 4, 125-134.
- Phang, S.-M. Shaharuddin, S. Noraishah, H. and Sasekumar, A. (1996). Studies on *Gracilaria changii* (Gracilariales, Rhodophyta) from Malaysian mangroves. *Hydrobiologia*. 326-327(1), 347-352.
- Phang, S. M. (2006). Seaweed resources in Malaysia: Current status and future prospects. *Aquatic Ecosystem Health & Management*. 9(2), 185-202.
- Pinchetti, J. L. G. Del Campo Fernández, E. Díez, P. M. and Reina, G. G. (1998a). Nitrogen availability influences the biochemical composition and photosynthesis of tank-cultivated *Ulva rigida* (Chlorophyta). *Journal of Applied Phycology*. 10(4), 383-389.
- Pinchetti, J. L. G. Fernandez, E. D. C. Diez, P. M. and Reina, G. G. (1998b). Nitrogen availability influences the biochemical composition and photosynthesis of tank-cultivated *Ulva rigida* (Chlorophyta). *Journal of Applied Phycology*. 10, 383-389.
- Plastino, E. and Oliveira, E. 1996. Approaches to the identification of terete Brazilian Gracilariaceae (Gracilariales, Rhodophyta). In: Lindstrom, S. & Chapman, D. (ed.) (eds.) *Fifteenth International Seaweed Symposium*. Springer Netherlands.

- Ponce-Palafox, J. Martinez-Palacios, C. A. and Ross, L. G. (1997). The effects of salinity and temperature on the growth and survival rates of juvenile white shrimp, *Penaeus vannamei*, Boone, 1931. *Aquaculture*. 157(1), 107-115.
- Qi, H. Zhang, Q. Zhao, T. Chen, R. Zhang, H. Niu, X. and Li, Z. (2005). Antioxidant activity of different sulfate content derivatives of polysaccharide extracted from *Ulva pertusa* (Chlorophyta) in vitro. *International Journal of Biological Macromolecules*. 37(4), 195-199.
- Raju, P. and Thomas, P. (1971). Experimental field cultivation of *Gracilaria edulis* (Gmel.) Silva. *Botanica Marina*. 14(2), 71-75.
- Ramus, J. and Rosenberg, G. (1980). Diurnal photosynthetic performance of seaweeds measured under natural conditions. *Marine Biology*. 56, 21-28.
- Raven, J. (1997). Inorganic carbon acquisition by marine autotrophs. *Advances in Botanical Research*. 27, 85-209.
- Ray, A. J. Dillon, K. S. and Lotz, J. M. (2011). Water quality dynamics and shrimp (*Litopenaeus vannamei*) production in intensive, mesohaline culture systems with two levels of biofloc management. *Aquacultural Engineering*. 45(3), 127-136.
- Ray, A. J. Lewis, B. L. Browdy, C. L. and Leffler, J. W. (2010a). Suspended solids removal to improve shrimp (*Litopenaeus vannamei*) production and an evaluation of a plant-based feed in minimal-exchange, superintensive culture systems. *Aquaculture*. 299, 89-98.
- Ray, A. J. Seaborn, G. Leffler, J. W. Wilde, S. B. Lawson, A. and Browdy, C. L. (2010b). Characterization of microbial communities in minimal-exchange, intensive aquaculture systems and the effects of suspended solids management. *Aquaculture*. 310(1-2), 130-138.



- Rice, D. L. and Lapointe, B. E. (1981). Experimental outdoor studies with *Ulva fasciata* delile. II. Trace metal chemistry. *Journal of Experimental Marine Biology and Ecology*. 54(1), 1-11.
- Rindi, F. Soler-Vila, A. and Guiry, M. 2012. Taxonomy of Marine Macroalgae Used as Sources of Bioactive Compounds. In: Hayes, M. (ed.) (eds.) *Marine Bioactive Compounds*. Springer US.
- Rivers, J. S. and Peckol, P. (1995). Summer decline of *Ulva lactuca* (chlorophyta) in a eutrophic embayment: Interactive effects of temperature and nitrogen availability. *Journal of Phycology*. 31(2), 223-228.
- Robertson-Andersson, D. Potgieter, M. Hansen, J. Bolton, J. Troell, M. Anderson, R. Halling, C. and Probyn, T. (2008). Integrated seaweed cultivation on an abalone farm in South Africa. *Journal of Applied Phycology*. 20(5), 579-595.
- Robertson-Andersson, J. J. B. D. V. and Kandjengo, D. S. L. (2008). Growing *Ulva* (Chlorophyta) in integrated systems as a commercial crop for abalone feed in South Africa: a SWOT analysis. *Journal of Applied Phycology*.
- Robertson, A. I. and Phillips, M. J. (1995). Mangroves as filters of shrimp pond effluent: predictions and biogeochemical research needs. *Hydrobiologia*. 295(1-3), 311-321.
- Rosas, C. Martinez, E. Gaxiola, G. Brito, R. Sánchez, A. and Soto, L. A. (1999). The effect of dissolved oxygen and salinity on oxygen consumption, ammonia excretion and osmotic pressure of *Penaeus setiferus* (Linnaeus) juveniles. *Journal of Experimental Marine Biology and Ecology*. 234(1), 41-57.
- Rosenberg, G. Littler, D. S. Littler, M. M. and Oliveira, E. C. (1995). Primary production and photosynthetic quotients of seaweeds from Sao Paulo State, Brazil. *Botanica Marina*. 38, 369-377.

- Roy, D. Hassan, K. and Boopathy, R. (2010). Effect of carbon to nitrogen ( C:N) ratio on nitrogen removal from shrimp production waste water using sequencing batch reactor. *Journal of Indian Microbiological Biotechnology*. 37, 1105-1110.
- Runcie, J. W. Ritchie, R. J. and Larkumb, A. W. D. (2003). Uptake kinetics and assimilation of inorganic nitrogen by *Catenella nipae* and *Ulva lactuca*. *Aquatic Botany*. 76, 155-174.
- Ruperez, P. (2002). Mineral content of edible marine seaweeds. *Food Chemistry*. 79(1), 23-26.
- Rupérez, P. (2002). Mineral content of edible marine seaweeds. *Food Chemistry*. 79(1), 23-26.
- Ryther, J. Corwin, N. Debusk, T. and Williams, L. (1981). Nitrogen uptake and storage by the red alga *Gracilaria tikvahiae* (McLachlan, 1979). *Aquaculture*. 26(1), 107-115.
- Sánchez-Machado, D. I. López-Cervantes, J. López-Hernández, J. and Paseiro-Losada, P. (2004). Fatty acids, total lipid, protein and ash contents of processed edible seaweeds. *Food Chemistry*. 85(3), 439-444.
- Sandifer, P. A. Hopkins, J. S. Stokes, A. D. and Browdy, C. L. (1993). Preliminary comparisons of the native *Penaeus setiferus* and Pacific *P. vannamei* white shrimp for pond culture in South Carolina, USA. *Journal of the World Aquaculture Society*. 24(3), 295-303.
- Santelices, B. and Doty, M. S. (1989). A review of *Gracilaria* farming. *Aquaculture*. 78(2), 95-133.
- Schuenhoff, A. Shpigel, M. Lupatsch, I. Ashkenazi, A. Msuya, F. E. and Neori, A. (2003). A semi-recirculating, integrated system for the culture of fish and seaweed. *Aquaculture*. 221(1-4), 167-181.

- Schuur, A. M. (2003). Evaluation of biosecurity applications for intensive shrimp farming. *Aquacultural Engineering*. 28(1), 3-20.
- Shang, Y. C. (1976). Economic aspects of *Gracilaria* culture in Taiwan. *Aquaculture*. 8(1), 1-7.
- Sheehy, J. E. Dionora, M. J. A. Mitchell, P. L. Peng, S. Cassman, K. G. Lemaire, G. and Williams, R. L. (1998). Critical nitrogen concentrations: implications for high-yielding rice (*Oryza sativa* L.) cultivars in the tropics. *Field Crops Research*. 59(1), 31-41.
- Sheehya, J. E. Dionoraa, M. J. A. Mitchell, P. L. Penga, S. Cassman, K. G. Lemaire, G. and Williams, R. L. (1998). Critical nitrogen concentrations: implications for high-yielding rice (*Oryza sativa* L.) cultivars in the tropics. *Field Crops Research*. 59, 31-41.
- Shi, H. Zheng, W. Zhang, X. Zhu, M. and Ding, D. (2013). Ecological–economic assessment of monoculture and integrated multi-trophic aquaculture in Sanggou Bay of China. *Aquaculture*. 410–411(0), 172-178.
- Shin, P. K. S. Yau, F. N. Chow, S. H. K.K. Tai and Cheung, S. G. (2002). Responses of the green-lipped mussel *Perna viridis* (L.) to suspended solids. *Marine Pollution Bulletin*. 45, 157-162.
- Sivapalan, A. and Theivendirarajah, K. (1985). Studies on the agarophyte, *Gracilaria edulis* - Experimental field cultivation and methods of improving yield and quality of agar. *Journal Natural Sciences Country Sri lanka*. 13(2), 197-212.
- SJER 2010. Natural and green environment-Iskandar Malaysia. In: Malaysia, I. (ed.) *Part 3: Physical Planning Initiatives*. Johor: South Johor Economic Region.
- Skriptsova, A. V. and Miroshnikova, N. V. (2011). Laboratory experiment to determine the potential of two macroalgae from the Russian Far-East as biofilters for integrated multi-trophic aquaculture (IMTA). *Bioresource Technology*. 102(3), 3149-3154.

- Smith, D. M. Burford, M. A. Tabrett, S. J. Irvin, S. J. and Ward, L. (2002). The effect of feeding frequency on water quality and growth of the black tiger shrimp (*Penaeus monodon*). *Aquaculture*. 207(1–2), 125-136.
- Smith, S. L. Yamanaka, Y. Pahlow, M. and Oschlies, A. (2009). Optimal uptake kinetics: physiological acclimation explains the pattern of nitrate uptake by phytoplankton in the ocean. *Marine Ecology Progress Series*. 384(1-12).
- Strickland, J. D. H. and Parsons, T. R. (1972). *A Practical Handbook of Seawater Analysis*. (pp. Canada: Fisheries Research Board of Canada.
- Stumm, W. and Morgan, J. (1981). *Aquatic Chemistry*. (pp. New York: Wiley.
- Subbaramaiah, K. and Thomas, P. (1990). Raft cultivation of *Gracilaria edulis* (Gmel.) Silva. *Proceedings: Plant Sciences*. 100(2), 123-127.
- Taboada, C. Millán, R. and Míguez, I. (2010). Composition, nutritional aspects and effect on serum parameters of marine algae *Ulva rigida*. *Journal of the Science of Food and Agriculture*. 90(3), 445-449.
- Taylor, R. B. Peek, J. T. A. and Rees, T. a. V. (1998). Scaling of ammonium uptake by seaweeds to surface area:volume ratio : geographical variation and the role of uptake by passive diffusion. *Marine Ecology Progress Series*. 169, 143-148.
- Teichert-Coddington, D. Rodriguez, R. and Toyofuku, W. (1994). Cause of cyclic variation in Honduran shrimp production. *World Aquaculture*. 25, 57-57.
- Teichert-Coddington, D. R. Rouse, D. B. Potts, A. and Boyd, C. E. (1999). Treatment of harvest discharge from intensive shrimp ponds by settling. *Aquacultural Engineering*. 19, 147-161.
- Tenore, K. R. and Dunstan, W. M. (1973). Comparison of feeding and biodeposition of three bivalves at different food levels. *Marine Biology*. 21(3), 190-195.

- Thakur, D. P. and Lin, C. K. (2003). Water quality and nutrient budget in closed shrimp (*Penaeus monodon*) culture systems. *Aquacultural Engineering*. 27(3), 159-176.
- Thirumaran, G. and Anantharaman, P. (2009). Daily growth rate of field farming seaweed *Kappaphycus alvarezii* (Doty) Doty ex P. Silva in Vellar Estuary. *World J. Fish. Mar. Sci.* 1(3), 144-153.
- Thuy, H. T. T. Nga, L. P. and Loan, T. T. C. (2011). Antibiotic contaminants in coastal wetlands from Vietnamese shrimp farming. *Environmental Science Pollution Research*. 18, 835-841.
- Tian, X. Li, D. Dong, S. Yan, X. Qi, Z. Liu, G. and Lu, J. (2001). An experimental study on closed-polyculture of penaeid shrimp with tilapia and constricted tagelus. *Aquaculture*. 202(1-2), 57-71.
- Tilley, D. R. Badrinarayanan, H. Rosati, R. and Son, J. (2002). Constructed wetlands as recirculation filters in large-scale shrimp aquaculture. *Aquaculture Engineering*. 26, 81-109.
- Todd, P. A. Ong, X. and Chou, L. M. (2010). Impacts of pollution on marine life in Southeast Asia. *Biodiversity and Conservation*. 19(4), 1063-1082.
- Troell, M. Halling, C. Neori, A. Chopin, T. Buschmann, A. H. Kautsky, N. and Yarish, C. (2003a). Integrated mariculture: asking the right questions. *Aquaculture*. 226(1-4), 69-90.
- Troell, M. Halling, C. Neori, A. Chopin, T. Buschmann, A. H. Kautsky, N. and Yarish, C. (2003b). Integrated mariculture: asking the right questions. *Aquaculture* 226, 69-90.
- Troell, M. Halling, C. Nilsson, A. Buschmann, A. H. Kautsky, N. and Kautsky, L. (1997). Integrated marine cultivation of *Gracilaria chilensis* (Gracilariales,

- Rhodophyta) and salmon cages for reduced environmental impact and increased economic output. *Aquaculture*. 156(1-2), 45-61.
- Troell, M. Joyce, A. Chopin, T. Neori, A. Buschmann, A. H. and Fang, J.-G. (2009). Ecological engineering in aquaculture — Potential for integrated multi-trophic aquaculture (IMTA) in marine offshore systems. *Aquaculture*. 297(1–4), 1-9.
- Troell, M. Rönnbäck, P. Halling, C. Kautsky, N. and Buschmann, A. (1999). Ecological engineering in aquaculture: use of seaweeds for removing nutrients from intensive mariculture. *Journal of Applied Phycology*. 11, 89-97.
- Trott, L. A. and Alongi, D. M. (2000). The Impact of Shrimp Pond Effluent on Water Quality and Phytoplankton Biomass in a Tropical Mangrove Estuary. *Marine Pollution Bulletin*. 40, 947-951.
- Tsagkamilis, P. Danielidis, D. Dring, M. J. and Katsaros, C. (2010). Removal of phosphate by the green seaweed *Ulva lactuca* in a small-scale sewage treatment plant (Ios Island, Aegean Sea, Greece). *Journal of Applied Phycology*. 22(3), 331-339.
- Tseng, K.-F. Su, H.-M. and Su, M.-S. (1998). Culture of *Penaeus monodon* in a recirculating system. *Aquacultural Engineering*. 17(2), 138-147.
- Turpin, D. H. (1991). Effects of inorganic N availability on algal photosynthesis and carbon metabolism. *Journal of Phycology*. 27(1), 14-20.
- Ugarte, R. and Santelices, B. (1992). Experimental tank cultivation of *Gracilaria chilensis* in central Chile. *Aquaculture*. 101(1), 7-16.
- Ursi, S. Guimaraes, M. and Plastino, E. M. (2008). Deleterious effect of TRIS buffer on growth rates and pigment content of *Gracilaria birdiae* Plastino & E.C. Oliveira (Gracilariales, Rhodophyta). *Acta Botanica Brasilica*. 22(3), 891-896.
- Valente, L. M. P. Gouveia, A. Rema, P. Matos, J. Gomes, E. F. and Pinto, I. S. (2006). Evaluation of three seaweeds *Gracilaria bursa-pastoris*, *Ulva rigida* and *Gracilaria*

cornea as dietary ingredients in European sea bass (*Dicentrarchus labrax*) juveniles. *Aquaculture*. 252(1), 85-91.

Vandermeulen, H. and Gordin, H. (1990). Ammonium uptake using *Ulva* (chlorophyta) in intensive fishpond systems - mass-culture and treatment of effluent. *Journal of Applied Phycology*. 2(4), 363-374.

Vijayaraghavan, K. Ahmad, D. and Fadzin, T. S. B. (2008). In situ hypochlorous acid generation for the treatment of brackish shrimp aquaculture wastewater. *Aquaculture Research*. 39, 449-456.

Wahab, M. A. Bergheim, A. and Braaten, B. (2003). Water quality and partial mass budget in extensive shrimp ponds in Bangladesh. *Aquaculture*. 218(1), 413-423.

Walne, P. R. (1972). The Influence of Current Speed, Body Size and Water Temperature On the Filtration Rate of Five Species of Bivalves. *Journal of the Marine Biological Association of the United Kingdom*. 52(02), 345-374.

Wang, R. Xiao, H. Wang, Y. Zhou, W. and Tang, X. (2007). Effects of three macroalgae, *Ulva linza* (Chlorophyta), *Corallina pilulifera* (Rhodophyta) and *Sargassum thunbergii* (Phaeophyta) on the growth of the red tide microalga *Prorocentrum donghaiense* under laboratory conditions. *Journal of Sea Research*. 58(3), 189-197.

Wyban, J. Walsh, W. A. and Godin, D. M. (1995). Temperature effects on growth, feeding rate and feed conversion of the Pacific white shrimp (*Penaeus vannamei*). *Aquaculture*. 138, 267-279.

Xu, Y. Fang, J. and Wei, W. (2008a). Application of *Gracilaria lichenoides* (Rhodophyta) for alleviating excess nutrients in aquaculture. *Journal of Applied Phycology*. 20, 199-203.

- Xu, Y. Fang, J. and Wei, W. (2008b). Application of *Gracilaria lichenoides* (Rhodophyta) for alleviating excess nutrients in aquaculture. *Journal of Applied Phycology*. 20(2), 199-203.
- Xu, Z. Zou, D. and Gao, K. (2010). Effects of elevated CO<sub>2</sub> and phosphorus supply on growth, photosynthesis and nutrient uptake in the marine macroalgae *Gracilaria lemaneiformis* (Rhodophyta). *Botanica Marina*. 53, 123-129.
- Yang, H. Zhou, Y. Mao, Y. Li, X. Liu, Y. and Zhang, F. (2005a). Growth characters and photosynthetic capacity of *Gracilaria lemaneiformis* as a biofilter in a shellfish farming area in Sanggou Bay, China. *Journal of Applied Phycology*. 17, 199-206.
- Yang, H. Zhou, Y. Mao, Y. Li, X. Liu, Y. and Zhang, F. (2005b). Growth characters and photosynthetic capacity of *Gracilaria lemaneiformis* as a biofilter in a shellfish farming area in Sanggou Bay, China. *Journal of Applied Phycology*. 17, 199-206.
- Yang, Y.-F. Fei, X.-G. Song, J.-M. Hu, H.-Y. Wang, G.-C. and Chung, I. K. (2006a). Growth of *Gracilaria lemaneiformis* under different cultivation conditions and its effects on nutrient removal in Chinese coastal waters. *Aquaculture*. 254(1-4), 248-255.
- Yang, Y.-F. Fei, X.-G. Song, J.-M. Hu, H.-Y. Wang, G.-C. and Chung, I. K. (2006b). Growth of *Gracilaria lemaneiformis* under different cultivation conditions and its effects on nutrient removal in Chinese coastal waters. *Aquaculture*. 254, 248-255.
- Yap, C. K. Ismail, A. Tan, S. G. and Omar, H. (2003). Accumulation, depuration and distribution of cadmium and zinc in the green-lipped mussel *Perna viridis* (Linnaeus) under laboratory conditions. *Hydrobiologia*. 498(1-3), 151-160.
- Ye, N. Wang, H. and Wang, G. (2006). Formation and early development of tetraspores of *Gracilaria lemaneiformis* (Gracilaria, Gracilariaceae) under laboratory conditions. *Aquaculture*. 254(1), 219-226.



- Yokoyama, H. and Ishihi, Y. (2010). Bioindicator and biofilter function of *Ulva* spp. (Chlorophyta) for dissolved inorganic nitrogen discharged from a coastal fish farm - potential role in integrated multi-trophic aquaculture. *Aquaculture*. 310(1-2), 74-83.
- Yow, Y.-Y. Lim, P.-E. and Phang, S.-M. (2011). Genetic diversity of *Gracilaria changii* (Gracilariaceae, Rhodophyta) from west coast, Peninsular Malaysia based on mitochondrial *cox1* gene analysis. *Journal of Applied Phycology*. 23(2), 219-226.
- Yu, C.-H. Lim, P.-E. and Phang, S.-M. (2013). Effects of irradiance and salinity on the growth of carpospore-derived tetrasporophytes of *Gracilaria edulis* and *Gracilaria tenuistipitata var liui* (Rhodophyta). *Journal of Applied Phycology*. 25(3), 787-794.
- Zang, C. Huang, S. Wu, M. Du, S. Scholz, M. Gao, F. Lin, C. and Dong, Y. G. Y. (2011). Comparison of Relationships Between pH, Dissolved Oxygen and Chlorophyll a for Aquaculture and Non-aquaculture Waters. *Water Air Soil Pollution*. 219, 157-174.
- Zhou, Y. Yang, H. Hu, H. Liu, Y. Mao, Y. Zhou, H. Xu, X. and Zhang, F. (2006). Bioremediation potential of the macroalgae *Gracilaria lemaneiformis* (Rhodophyta) integrated into fed fish culture in coastal waters of north China. *Aquaculture*. 252(2-4), 264-276.
- Ziemann, D. A. Walsh, W. A. Saphore, E. G. and Fulton-Bennett, K. (1992). A Survey of Water Quality Characteristics of Effluent from Hawaiian Aquaculture Facilities. *Journal of the World Aquaculture Society*. 23(3), 180-191.
- Zou, D. (2005). Effects of elevated atmospheric CO<sub>2</sub> on growth, photosynthesis and nitrogen metabolism in the economic brown seaweed, *Hizikia fusiforme* (Sargassaceae, Phaeophyta). *Aquaculture*. 250, 726-735.
- Zou, D. and Gao, K. (2009). Effects of elevated CO<sub>2</sub> on the red seaweed *Gracilaria lemaneiformis* (Gigartinales, Rhodophyta) grown at different irradiance levels. 48(6), 510-517.