

**SYNTHESIS AND ANTIOXIDANT ACTIVITIES OF  
HYDROXYLATED COUMARINYL CHALCONES**

**MOHAMAD SHAFIQ BIN SULEIMAN**

**UNIVERSITI TEKNOLOGI MALAYSIA**

SYNTHESIS AND ANTIOXIDANT ACTIVITIES OF  
HYDROXYLATED COUMARINYL CHALCONES

MOHAMAD SHAFIQ BIN SULEIMAN

A dissertation submitted in partial fulfillment of the  
requirements for the award of the degree of  
Master of Chemistry

Faculty of Science  
Universiti Teknologi Malaysia

FEBRUARY 2016

## ACKNOWLEDGEMENT

In the name of Allah, The Most Gracious and The Most Merciful, peace and blessings of Allah Almighty to our beloved Prophet Muhammad S.A.W and his relatives, all his companions and those who have followed. Alhamdulillah, with His willing has allowed me to complete my research project.

I would like to express my most sincere gratitude to my beloved supervisor Dr. Shajarahtunnur binti Jamil for her trust, encouragement, kindness, guidance and supervision from the beginning up to the completion point of this project which has enabled me to gain so much experience and knowledge in this field.

I would also like to thank the Faculty of Science, Universiti Teknologi Malaysia (UTM) for providing me the facilities and equipments to complete my project. It was wonderful to have such great cooperation from all laboratory staffs especially En. Azmi bin Md Rais, Pn. Suhani binti Md Tah, En. Mohd Faizz bin Mohamad Fuad, En. Rasyidi bin Abd Mubin and all other postgraduate students. Without their help, I would not think this project would be a success.

Thanks also to the members of my research group for the guidance and discussions necessary for the project that I am conducting. My special thanks also go to my beloved parents for their encouragement as well as moral and financial support and emphasis on the value of education. Not forgetting, my classmates and my best friends for all their support and thoughts, thank you.

Lastly, my special gratitude goes to the people that had contributed either directly or indirectly for the completion of this study.

## ABSTRACT

Hydroxylated coumarins and chalcones are known to possess potent antioxidant activities. The present study highlights the synthesis of two hydroxycoumarins namely 3-acetyl-7-hydroxycoumarin and 3-acetyl-6-hydroxycoumarin. Both compounds were synthesized using the Knoevenagel condensation method with respective hydroxybenzaldehydes under basic conditions. The synthesized hydroxycoumarins served as an intermediate for the subsequent coupling reaction to produce three new coumarinyl chalcones known as 7-hydroxy-3-[3-(4'-hydroxyphenyl)prop-2-enoyl]-2*H*-1-benzopyran-2-one, 6-hydroxy-3-[3-(4'-hydroxyphenyl)prop-2-enoyl]-2*H*-1-benzopyran-2-one and 7-hydroxy-3-[3-(3',4'-dihydroxyphenyl)prop-2-enoyl]-2*H*-1-benzopyran-2-one respectively. The target products were synthesized *via* the Claisen-Schmidt condensation reaction utilizing the Lewis acid, boron trifluoride-etherate (BF<sub>3</sub>-Et<sub>2</sub>O) in 1,4-dioxane. The structures of the synthetic compounds were confirmed by spectroscopic techniques which includes the Infrared (IR) and Nuclear Magnetic Resonance (NMR) (<sup>1</sup>H, <sup>13</sup>C and DEPT) Spectroscopies. The antioxidant activities of all synthesized compounds were evaluated using three antioxidant assays known as the 2,2-diphenyl-1-picrylhydrazyl (DPPH), 2,2'-azinobis-(3-ethylbenzothiazoline-6-sulfonic acid) (ABTS) and Ferric Reducing Ability of Plasma (FRAP) assays using the Bio-Tek Epoch microplate reader. Among the compounds tested, the coumarinyl chalcones exhibited a better antioxidant activity compared to the hydroxycoumarins. 7-Hydroxy-3-[3-(3',4'-dihydroxyphenyl)prop-2-enoyl]-2*H*-1-benzopyran-2-one proved to be the best antioxidant in all assays with SC<sub>50</sub> values of 0.57 mM and 0.036 mM in the ABTS and DPPH assays respectively. The FRAP value of this compound falls between the range of 0.09 mM to 1.67 mM. The values were comparable with the positive control used in these assays known as butylated hydroxyanisole (BHA).

## ABSTRAK

Hidroksi kumarin dan kalkon terkenal dengan aktiviti antioksidan yang kuat. Kajian ini menonjolkan hasil sintesis bagi dua hidroksikumarin yang bernama 3-asetil-7-hidroksikumarin dan 3-asetil-6-hidroksikumarin. Kedua-dua sebatian ini disintesis menggunakan kaedah kondensasi Knoevenagel dengan hidroksibenzaldehid masing-masing dalam keadaan bes. Dua hidroksikumarin yang disintesis ini digunakan sebagai sebatian perantara untuk tindak balas penggandingan yang dilakukan berikutnya untuk menghasilkan tiga sebatian kumarinil kalkon yang masing-masing dikenali sebagai 7-hidroksi-3-[3-(4'-hidroksifenil)prop-2-enoil]-2*H*-1-benzopiran-2-on, 6-hidroksi-3-[3-(4'-hidroksifenil)prop-2-enoil]-2*H*-1-benzopiran-2-on dan 7-hidroksi-3-[3-(3',4'-dihidroksifenil)prop-2-enoil]-2*H*-1-benzopiran-2-on. Ketiga-tiga sebatian sasaran disintesis melalui reaksi kondensasi Claisen-Schmidt dengan menggunakan sejenis asid Lewis, boron-trifluorida eterat (BF<sub>3</sub>-Et<sub>2</sub>O) dalam 1,4-dioksan. Struktur sebatian sintetik ini disahkan menggunakan teknik spektroskopi termasuk Spektroskopi Inframerah (IR) dan Resonans Magnet Nukleus (NMR) (<sup>1</sup>H, <sup>13</sup>C dan DEPT). Penilaian antioksidan ke atas sebatian yang disintesis juga dijalankan menerusi tiga ujian antioksidan iaitu ujian 2,2-difenil-1-pikrilhidrazil (DPPH), ujian 2,2'-azinobis-(3-etilbenzotiazolin-6-asid sulfonik) (ABTS) dan ujian Kemampuan Plasma Menurunkan Ferik (FRAP) menggunakan Bio-Tek pembaca mikroplat Epoch. Antara semua sebatian yang diuji, sebatian kumarinil kalkon menunjukkan aktiviti antioksidan yang lebih tinggi berbanding sebatian hidroksikumarin. 7-Hidroksi-3-[3-(3',4'-dihidroksifenil)prop-2-enoil]-2*H*-1-benzopiran-2-on terbukti menjadi antioksidan terbaik dalam semua ujian antioksidan yang dijalankan dengan bacaan SC<sub>50</sub> 0.57 mM dan 0.036 mM masing-masing dalam ujian ABTS dan ujian DPPH. Nilai FRAP sebatian ini ialah antara 0.09 mM hingga 1.67 mM. Nilai-nilai ini dapat dibandingkan dengan kawalan positif yang digunakan dalam ujian-ujian ini yang dikenali sebagai butil hidroksianisol (BHA).

**TABLE OF CONTENT**

<b>CHAPTER</b>	<b>TITLE</b>	<b>PAGE</b>
	<b>DECLARATION</b>	ii
	<b>ACKNOWLEDGEMENT</b>	iii
	<b>ABSTRACT</b>	iv
	<b>ABSTRAK</b>	v
	<b>TABLE OF CONTENTS</b>	vi
	<b>LIST OF TABLES</b>	ix
	<b>LIST OF SCHEMES</b>	x
	<b>LIST OF FIGURES</b>	xii
	<b>LIST OF ABBREVIATIONS</b>	xiii
	<b>LIST OF APPENDICES</b>	xiv
<b>1</b>	<b>INTRODUCTION</b>	
	1.1 Background of Study	1
	1.2 Problem Statement	3
	1.3 Objectives of Study	4
	1.4 Scope of Study	4
	1.5 Significance of Study	5
<b>2</b>	<b>LITERATURE REVIEW</b>	
	2.1 Chalcones	6
	2.1.1 Naturally Occurring Chalcones	7
	2.1.2 Biosynthesis of Chalcones	11
	2.1.3 Biological Activities of Chalcones	13
	2.1.4 Synthesis of Chalcones	15
	2.2 Coumarins	19

2.2.1	Naturally Occurring Coumarins	21
2.2.2	Biosynthesis of Coumarins	23
2.2.3	Biological Activities of Coumarins	25
2.2.4	Synthesis of Coumarins	26
2.3	Coumarinyl Chalcones	28
2.3.1	Biological Activities of Coumarinyl Chalcones	29
2.3.2	Synthesis of Coumarinyl Chalcones	30
2.4	Antioxidant Activity	32
2.4.1	Ferric Reducing Ability of Plasma (FRAP)	33
2.4.2	2,2'-Azinobis-(3-ethylbenzothiazoline-6- sulfonic acid) (ABTS)	34
2.4.3	2,2-Diphenyl-1-picrylhydrazyl (DPPH) Free Radicals	34
<b>3</b>	<b>RESULTS AND DISCUSSION</b>	
3.1	Synthesis of Hydroxycoumarins	36
3.1.1	Synthesis of 3-acetyl-7-hydroxycoumarin ( <b>97</b> ) and 3-acetyl-6-hydroxycoumarin ( <b>98</b> )	36
3.2	The Synthesis of Hydroxylated Coumarinyl Chalcones	40
3.2.1	Synthesis of 7-Hydroxy-3-[3-(4'- hydroxyphenyl)prop-2-enoyl]-2 <i>H</i> -1- benzopyran-2-one ( <b>105</b> ), 6-Hydroxy-3- [3-(4'-hydroxyphenyl)prop-2-enoyl]-2 <i>H</i> - 1-benzopyran-2-one ( <b>106</b> ) and 7- Hydroxy-3-[3-(3',4'-dihydroxyphenyl) prop-2-enoyl]-2 <i>H</i> -benzopyran-2-one ( <b>112</b> )	40
3.3	Antioxidant Properties of the Synthesized Compounds	47

3.3.1	FRAP Assay	48
3.3.2	ABTS Assay	50
3.3.3	DPPH Assay	51
3.3.4	Structural Features related to Antioxidant Properties	53
<b>4</b>	<b>EXPERIMENTAL</b>	
4.1	General Procedures	55
4.2	Instrumentations	55
4.3	Solvents and Chemicals	56
4.4	Synthesis of 3-Acetyl-7-hydroxycoumarin ( <b>97</b> )	56
4.5	Synthesis of 3-Acetyl-6-hydroxycoumarin ( <b>98</b> )	57
4.6	Synthesis of 7-Hydroxy-3-[3-(4'-hydroxyphenyl)prop-2-enoyl]-2 <i>H</i> -1-benzopyran-2-one ( <b>105</b> )	57
4.7	Synthesis of 6-Hydroxy-3-[3-(4'-hydroxyphenyl)prop-2-enoyl]-2 <i>H</i> -1-benzopyran-2-one ( <b>106</b> )	58
4.8	Synthesis of 7-Hydroxy-3-[3-(3',4'-dihydroxyphenyl)prop-2-enoyl]-2 <i>H</i> -1-benzopyran-2-one ( <b>112</b> )	59
4.9	Antioxidant Assays	59
4.9.1	FRAP Assay	60
4.9.2	ABTS Assay	60
4.9.3	DPPH Assay	61
4.9.4	Statistical Analysis	61
<b>5</b>	<b>CONCLUSION AND RECOMMENDATIONS</b>	
5.1	Conclusion	62
5.2	Recommendations	63
	<b>REFERENCES</b>	64
	Appendices 1 - 27	76-102



**LIST OF TABLES**

<b>TABLE NO.</b>	<b>TITLE</b>	<b>PAGE</b>
2.1	Classification of Coumarins and Their Structures	20
3.1	Ferric Reducing Antioxidant Power (FRAP) Results	49
3.2	ABTS Radical Scavenging Results	51
3.3	DPPH Radical Scavenging Results	52

## LIST OF SCHEMES

SCHEME NO.	TITLE	PAGE
2.1	Isomerization of the Chalcone-Flavanone System	7
2.2	The Biosynthetic Pathway of 2',4',4,6'-Tetrahydroxychalcone	12
2.3	Synthesis of Chalcone <i>via</i> Claisen-Schmidt condensation	15
2.4	Solvent-free reaction to produce Substituted Chalcones	16
2.5	Synthesis of Chalcones using SOCl <sub>2</sub> and EtOH as catalysts	16
2.6	Synthesis of 4,4'-Dihydroxychalcone using BF <sub>3</sub> -Et <sub>2</sub> O	17
2.7	The Algar-Flynn-Oyamada reaction to produce Aurones ( <b>51</b> ) and Flavones ( <b>52</b> )	18
2.8	Biosynthesis of Coumarin	24
2.9	Synthesis of Coumarins <i>via</i> Perkin reaction	26
2.10	Pechmann condensation	26
2.11	Pechmann Synthesis carried out by Maheshwara <i>et al.</i> (2006) [81]	27
2.12	Knoevenagel condensation in the presence of an Ionic Liquid	27
2.13	Classic Knoevenagel condensation	28
2.14	Synthesis method of Coumarinyl Chalcones	31
2.15	Claisen-Schmidt condensation to produce Coumarinyl Chalcones	32
3.1	Reaction to synthesize hydroxycoumarins ( <b>97</b> ) and ( <b>98</b> )	37

3.2	Reaction mechanism to produce hydroxycoumarins <b>(97)</b> and <b>(98)</b>	39
3.3	Synthesis of Coumarinyl Chalcones <b>(105)</b> , <b>(106)</b> and <b>(112)</b>	41
3.4	Proposed dissociation process of BF <sub>3</sub> complex	42
3.5	Reaction mechanism to produce Coumarinyl Chalcones <b>(105)</b> and <b>(106)</b>	45
3.6	Reaction mechanism to generate ABTS radicals and its Electron Transfer process	50
3.8	DPPH radical scavenging action	51
3.9	Proposed mechanism of Antioxidant Capacity of Compound <b>(112)</b>	54

**LIST OF FIGURES**

<b>FIGURE NO.</b>	<b>TITLE</b>	<b>PAGE</b>
3.1	FeSO <sub>4</sub> Calibration Curve	49

**LIST OF ABBREVIATIONS**

$\lambda$	-	Wavelength
ABTS	-	2,2'-Azinobis-(3-ethyl-benzothiazoline-6-sulfonic acid)
Ac <sub>2</sub> O	-	Acetic Anhydride
BHA	-	Butylated Hydroxy Anisole
CC	-	Column Chromatography
COSY	-	Correlation Spectroscopy
<sup>13</sup> C-DEPT	-	Carbon-13 Distortionless Enhancement by Polarization Transfer
<sup>13</sup> C-NMR	-	Carbon-13 Nuclear Magnetic Resonance Spectroscopy
DPPH	-	2,2-Diphenyl-1-picrylhydrazyl
ESR	-	Electron Spin Resonance
EtOAc	-	Ethyl Acetate
EtOH	-	Ethanol
FRAP	-	Ferric Reducing Ability of Plasma
<sup>1</sup> H-NMR	-	Proton Nuclear Magnetic Resonance Spectroscopy
IR	-	Infrared Spectroscopy
<i>J</i>	-	Coupling Constant
EtOK	-	Potassium Acetate
EtONa	-	Sodium Acetate
ORAC	-	Oxygen Radical Absorbance Capacity
PAL	-	Phenylalanine Ammonia Lyase
PMN	-	Polymorphonucleate
TLC	-	Thin Layer Chromatography
UDP	-	Uridine Diphosphate
UV	-	Ultraviolet

## LIST OF APPENDICES

APPENDIX	TITLE	PAGE
1	IR spectrum of 3-acetyl-7-hydroxycoumarin ( <b>97</b> )	76
2	<sup>1</sup> H NMR spectrum of 3-acetyl-7-hydroxycoumarin ( <b>97</b> )	77
3	<sup>1</sup> H NMR spectrum of 3-acetyl-7-hydroxycoumarin ( <b>97</b> ) (Expansion)	78
4	IR spectrum of 3-acetyl-6-hydroxycoumarin ( <b>98</b> )	79
5	<sup>1</sup> H NMR spectrum of 3-acetyl-6-hydroxycoumarin ( <b>98</b> )	80
6	<sup>1</sup> H NMR spectrum of 3-acetyl-6-hydroxycoumarin ( <b>98</b> ) (Expansion)	81
7	IR spectrum of 7-Hydroxy-3-[3-(4'-hydroxyphenyl) prop-2-enoyl]-2 <i>H</i> - benzopyran-2-one ( <b>105</b> )	82
8	<sup>1</sup> H NMR spectrum of 7-Hydroxy-3-[3-(4'-hydroxyphenyl) prop-2-enoyl]-2 <i>H</i> -1-benzopyran-2-one ( <b>105</b> ) (Expansion)	83
9	<sup>1</sup> H NMR spectrum of 7-Hydroxy-3-[3-(4'-hydroxyphenyl) prop-2-enoyl]-2 <i>H</i> -1-benzopyran-2-one ( <b>105</b> )	84
10	COSY spectrum of 7-Hydroxy-3-[3-(4'-hydroxyphenyl) prop-2-enoyl]-2 <i>H</i> -1-benzopyran-2-one ( <b>105</b> ) (Expansion)	85
11	<sup>13</sup> C NMR spectrum of 7-Hydroxy-3-[3-(4'-hydroxyphenyl) prop-2-enoyl]-2 <i>H</i> -1-benzopyran-2-one ( <b>105</b> )	86

12	<sup>13</sup> C NMR spectrum of 7-Hydroxy-3-[3-(4'-hydroxyphenyl)prop-2-enoyl]-2 <i>H</i> -1-benzopyran-2-one ( <b>105</b> ) (Expansion)	87
13	<sup>13</sup> C NMR and DEPT spectra of 7-Hydroxy-3-[3-(4'-hydroxyphenyl)prop-2-enoyl]-2 <i>H</i> -1-benzopyran-2-one ( <b>105</b> )	88
14	IR spectrum of 6-Hydroxy-3-[3-(4'-hydroxyphenyl)prop-2-enoyl]-2 <i>H</i> - benzopyran-2-one ( <b>106</b> )	89
15	<sup>1</sup> H NMR spectrum of 6-Hydroxy-3-[3-(4'-hydroxyphenyl)prop-2-enoyl]-2 <i>H</i> -1-benzopyran-2-one ( <b>106</b> )	90
16	<sup>1</sup> H NMR spectrum of 6-Hydroxy-3-[3-(4'-hydroxyphenyl)prop-2-enoyl]-2 <i>H</i> -1-benzopyran-2-one ( <b>106</b> ) (Expansion)	91
17	COSY spectrum of 6-Hydroxy-3-[3-(4'-hydroxyphenyl)prop-2-enoyl]-2 <i>H</i> -1-benzopyran-2-one ( <b>106</b> ) (Expansion)	92
18	<sup>13</sup> C NMR spectrum of 6-hydroxy-3-[3-(4'-hydroxyphenyl)prop-2-enoyl]-2 <i>H</i> -1-benzopyran-2-one ( <b>106</b> )	93
19	<sup>13</sup> C NMR spectrum of 6-hydroxy-3-[3-(4'-hydroxyphenyl)prop-2-enoyl]-2 <i>H</i> -1-benzopyran-2-one ( <b>106</b> ) (Expansion)	94
20	<sup>13</sup> C NMR and DEPT spectrum of 6-hydroxy-3-[3-(4'-hydroxyphenyl)prop-2-enoyl]-2 <i>H</i> -1-benzopyran-2-one ( <b>106</b> )	95
21	IR spectrum of 7-Hydroxy-3-[3-(3',4'-dihydroxyphenyl)prop-2-enoyl]-2 <i>H</i> - benzopyran-2-one ( <b>112</b> )	96
22	<sup>1</sup> H NMR spectrum of 7-Hydroxy-3-[3-(3',4'-dihydroxyphenyl)prop-2-enoyl]-2 <i>H</i> -1-benzopyran-2-one ( <b>112</b> )	97
23	<sup>1</sup> H NMR spectrum of 7-Hydroxy-3-[3-(3',4'-dihydroxyphenyl)prop-2-enoyl]-2 <i>H</i> -1-benzopyran-2-one ( <b>112</b> ) (Expansion)	98

24	COSY spectrum of 7-Hydroxy-3-[3-(3',4'-dihydroxyphenyl)prop-2-enoyl]-2 <i>H</i> -1-benzopyran-2-one ( <b>112</b> ) (Expansion)	99
25	<sup>13</sup> C NMR spectrum of 7-Hydroxy-3-[3-(3',4'-dihydroxyphenyl)prop-2-enoyl]-2 <i>H</i> -1-benzopyran-2-one ( <b>112</b> )	100
26	<sup>13</sup> C NMR spectrum of 7-Hydroxy-3-[3-(3',4'-dihydroxyphenyl)prop-2-enoyl]-2 <i>H</i> -1-benzopyran-2-one ( <b>112</b> ) (Expansion)	101
27	<sup>13</sup> C NMR and DEPT spectra of 7-Hydroxy-3-[3-(3',4'-dihydroxyphenyl)prop-2-enoyl]-2 <i>H</i> -1-benzopyran-2-one ( <b>112</b> )	102




## CHAPTER 1

### INTRODUCTION


#### 1.1 Background of Study

Naturally occurring compounds are produced as a result of biodiversity in which organisms interact with their surroundings to yield various chemical entities necessary for the organisms to survive [1]. Chalcone is a class of natural compounds that can be isolated from various plant species and their existence in the plant kingdom is vast. Chalcones are also known by the name of 1,3-diaryl-2-propen-1-one (**1**) [2]. There are quite a number of natural and synthetic chalcones that display useful bioactivities which includes cytotoxicity, chemoprotective, antibacterial, antifungal, antiviral and anti-inflammatory properties [3]. Based on the biosynthesis of flavonoids in plants, chalcones exist as an intermediate as well as the final product [2].



Another class of naturally occurring compounds is known as coumarin (**2**) or 1,2-benzopyrone. Naturally occurring coumarins are some of the most abundant chemicals in natural products [4]. It is the parent molecule of the compound dicoumarol. Compound (**2**) has the simplest structure within a huge class of phenolic compounds and they consist of a benzene ring fused to an  $\alpha$ -pyrone ring [5].

Previous reports have showed that compound (2) and its several simple derivatives have antitumor properties. The biological and pharmacological effects of these compounds largely depends on the position and the type of substituents attached to the compound [6]. Some reports have shown that coumarins possessed antimicrobial, inhibited lipooxygenase and cyclooxygenase metabolic pathways, antioxidant, inflammatory and antitumoral activities [7, 8].



(2)

A wide range of natural products such as the ones mentioned above can be taken as chemical scaffolds as they are able to provide templates with high potential for combinatorial chemistry since they have the ability to display chemical information in a three-dimensional space. These countless drug classes aids therapeutic areas of infectious diseases and oncology as they are able to interact with numerous specific targets within the cell. Also, for many years, they have been deliberated as the base molecules in the process of drug discovery and development. Libraries are constructed with a basis of those scaffolds thus having the potential for both lead discovery and lead optimization. In lead discovery the compound is hoped to have an effect against targets unrelated to the original activity of the natural product and in lead optimization the compound is derivatized with the hope of improving its properties over the natural product [9].

Nowadays, hybrid molecules are the current trend in drug development. Several biological characteristics are able to be observed with these hybrid molecules. Recently, several reports proposed that coumarins having coupled with other molecules possessing a different biological activity will exhibit dual bioactivities. The coupling process will also enhance the properties of compounds to exhibit activities such as antiplatelet, antioxidant and anti-inflammatory activities [10 - 12].

In recent years, the demands for natural product inspired drug-like molecules or their libraries have hiked and therefore, it imposes the necessity for the development of reaction sequences and linking strategies that allow complex and assorted target molecules to be constructed in a more facile and reliable manner [13]. Therefore, the overall mission or target in any organic synthesis is to build or construct any organic molecules that are desired. Synthesizing bioactive natural compounds are among the goal in doing organic synthesis. Presently, organic synthesis is considered to be very important since natural products synthesis is not the only pathway to synthesis compounds having useful properties anymore. Organic synthesis has also evolved that some useful properties are able to be discovered due to the synthetic studies conducted [2].

## **1.2 Problem Statement**

Recent developments in pharmacology are looking into the coupling or combination of two pharmacophores within a molecule. These combinations allow the availability of active sites that are able to accommodate two different targets within the same molecule. The coupling of these pharmacophores provides a way to surpass drug resistance [14] and lowering the emergence of new resistant strains [15]. Cellular oxidative stress are created due to a rise in free radicals, it plays an important role in the aging process through pathogenesis apart from other diseases which includes cancer, atherosclerosis, diabetes and Alzheimer's disease [16, 17]. Hence, it would be intriguing to synthesize hybrid molecules and investigate whether these molecules are able to relieve oxidative stress as the research and development of antioxidants have drawn a great deal of attention in recent years [18].






### 1.3 Objectives of Study

This study was designed based on the following objectives:

- To synthesize hydroxycoumarins and hydroxylated coumarinyl chalcones.
- To characterize the structure of synthetic compounds using Nuclear Magnetic Resonance (NMR) and Infrared Spectroscopies (IR).
- To evaluate the antioxidant properties of all synthesized compounds.

### 1.4 Scope of Study

The synthesis of hydroxycoumarins involved the usage of 2,4-dihydroxybenzaldehyde (**3**), 2,5-dihydroxybenzaldehyde (**4**) and ethyl acetoacetate (**5**) as the starting materials. Piperidine and glacial acetic acid acted as the catalyst and co-catalyst respectively. The subsequent reaction to produce the desired coumarinyl chalcones utilized the synthetic hydroxycoumarins with either 4-hydroxybenzaldehyde (**6**) or 3,4-dihydroxybenzaldehyde (**7**) as the starting materials. The catalyst used was the Lewis acid, boron-trifluoride etherate ( $\text{BF}_3\text{-Et}_2\text{O}$ ) and dioxane acted as the solvent. The reactions were monitored by means of the thin layer chromatographic (TLC) technique and column chromatography (CC) was adapted for the purification process.



The synthetic compounds were analysed using several spectroscopic methods which consisted of infrared (IR) spectroscopy, 1D NMR ( $^1\text{H}$ ,  $^{13}\text{C}$  and DEPT) and 2D NMR (COSY). All products are tested for their antioxidant activities using the Ferric Reducing Ability of Plasma (FRAP), the 2,2'-azinobis-(3-ethylbenzothiazoline-6-sulfonic acid) (ABTS) and 2,2 diphenyl-1-picrylhydrazyl (DPPH) assays. All of these assays were monitored using the Bio-Tek Epoch microplate reader.

## 1.5 Significance of Study

Coumarins are a type of heterocyclic molecules that positively impact the human health. The effects imposed by these compounds are due to the radical scavenging properties which are mostly connected with their antioxidant activities [19]. A number of publications reported the antioxidant activity of chalcones, particularly hydroxylated chalcones [20-24]. These studies indicate that hydroxychalcones are a potent radical scavenger [18]. The antioxidant properties are related closely to the radical scavenging potentials of these compounds and they are affected greatly by the substituents attached to the compounds. Much of this significant property is observed in compounds having hydroxyl groups or oxygenated substituents [25]. Therefore, this research is dedicated to synthesize several derivatives of natural hydroxycoumarins and also hybrid molecules known as coumarinyl chalcone which will have hydroxyl moieties. These compounds are expected to possess potent antioxidant properties. Through the development of these antioxidants, a cure for all free radical related diseases could be found or at least the risks of inflicting them could be minimized.

## REFERENCES

1. Mishra, B.B., and Tiwari, V.K. (2011). Natural Products: an evolving role in future drug discovery. *Eur. J. of Med. Chem.* **46(10)**, 4769-4807.
2. Suwito, H., Jumina, Mustoffa, Kristanti, A.N., and Puspitaningsih, N.N.T. (2014). Chalcones: Synthesis, structure diversity and pharmacological aspects. *J. Chem. Pharm. Res.* **6(5)**, 1076-1088
3. Dimmock, J.R., Elias, D.W., Beazely, M.A., and Kandepu, N.M. (1999). Bioactivity of Chalcones. In Atta-ur-Rahman (Ed.), *Curr. Med. Chem.* **6**, 1125-1149.
4. Prince, M., Campbell, C.T., Robertson, T.A., Wells, A.J., and Kleiner, H.E. (2006). Naturally occurring coumarins inhibit 7,12-dimethylbenz[*a*]anthracene DNA adduct formation in mouse mammary gland. *Carcinogenesis.* **27(6)**, 1204-1213.
5. Keating, G.J., and O'Kennedy, R. (1997). The chemistry and occurrence of coumarins. In O'Kennedy, R. and Thornes, R.D. (Eds.), *Coumarin: Biology, Application and Mode of Action*. West Sussex, England: John Wiley & Sons. 23-64.
6. Lopez-Gonzalez, J.S., Prado-Garcia, H., Molina-Guarneros, J.A., Mendoza-Patino, N., Rosas-Medina, P., and Mandoki, J.J. (2008). Mechanisms of Antitumor Effects of Simple Coumarins in Murine Melanoma and Human Non-small Cell Lung Carcinoma Cell Lines. In Edardes, J.P. (Ed.), *Coumarin Anticoagulant Research Progress*. New York, USA: Nova Science Publishers, Inc. 99-112.
7. Marshall, M.E., Mohler, J.L., Edmonds, K., Williams, B., Butler, K., Ryles, M., Weiss, L., Urban, D., Bueschen, A., Markiewicz, M., and Cloud, G. (1994). An updated review of the clinical development of coumarin (1,2-benzopyrone) and 7-hydroxycoumarin. *J. Cancer Res. Clin. Oncol.* **120**, Suppl: S39-S42.
8. Hoult, J.R., and Payá, M. (1996). Pharmacological and biochemical actions of simple coumarins: natural products with therapeutic potential. *Gen. Pharmacol.* **27(4)**, 713-722.
9. Hajduk, P.J., Galloway, W.R.J.D., and Spring, D.R. (2011). Drug discovery: A question of library design. *Nature.* **470(7332)**, 42-43.

10. Vilar, S., Quezada, E., Santana, L., Uriarte, E., Yáñez, M., Fraiz, N., Alcaide, C., Cano, E., & Orallo, F. (2006). Design, synthesis, and vasorelaxant and platelet antiaggregatory activities of coumarin-resveratrol hybrids. *Bioorg. Med. Chem. Lett.* **16**(2), 257-261.
11. Melagraki, G., Afantitis, A., Igglessi, M.O., Detsi, A., Koufaki, M., Kontogiorgis, C., and Hadjipavlou, L.D. (2009). Synthesis and evaluation of the antioxidant and anti-inflammatory activity of novel coumarin-3-aminoamides and their alpha-lipoic acid adducts. *Eur. J. of Med. Chem.* **44**(7), 3020-3026.
12. Song, H.Y., Ngai, M.H., Song, Z.Y., MacAry, P.A., Hobley, J., and Lear, M. J. (2009). Practical synthesis of maleimides and coumarin-linked probes for protein and antibody labelling via reduction of native disulfides. *Org. Biomol. Chem.* **7**(17), 3400-3406.
13. Samiulla, D.S., Vaidyanathan, V.V., Arun, P.C., Balan, G., Blaze, M., Bondre, S., Chandrasekhar, G., Gadakh, A., Kumar, R., Kharvi, G., Kim, H. O., Kumar, S., Malikayil, J.A., Moger, M., Mone, M.K., Nagarjuna, P., Ogbu, C., Pendhalkar, D., Rao, A.V., Rao, G.V., Sarma, V.K., Shaik, S., Sharma, G.V., Singh, S., Sreedhar, C., Sonawane, R., Timmanna, U., and Hardy, L.W. (2005). Rational selection of structurally diverse natural product scaffolds with favourable ADME properties for drug discovery. *Molec. Divers.* **9**(1-3), 131-139.
14. Hubschwerlen, C., Specklin, J.L., Sigwalt, C., Schroeder, S., and Locher, H. H. (2003). Design, Synthesis and Biological Evaluation of Oxazolidinone–Quinolone Hybrids. *Bioorg. Med. Chem.* **11**(10), 2313-2319.
15. Pingaew, R., Prachayasittikul, S., Ruchirawat, S., and Prachayasittikul, V. (2014). Synthesis and cytotoxicity of novel 4-(4-(substituted)-1H-1,2,3-triazol-1-yl)-N-phenethylbenzenesulfonamides. *Med. Chem. Res.* **23**(4), 1768-1780.
16. Pieta, P.G. (2000). Flavonoids as antioxidants. *J. Nat. Prod.* **63**(7), 1035-1042.
17. Maritim, A.C., Sanders, R.A., & Watkins III, J.B. (2003). Diabetes, oxidative stress, and antioxidants: A review. *J. Biochem. Mol. Toxicol.* **17**(1), 24-37.
18. Wang, G., Xue, Y., An, L., Zheng, Y., Dou, Y., Zhang, L., and Liu, Y. (2015). Theoretical study on the structural and antioxidant properties of some recently synthesized 2,4,5-trimethoxy chalcones. *Food Chem.* **171**, 89-97.

19. Bubols, G.B., Vianna, D.R., Medina-Remon, A., von Poser, G., Lamuela-Raventos, R.M., Eifler-Lima, V.L., and Garcia, S.C. (2013). The antioxidant activity of coumarins and flavonoids. *Mini. Rev. Med. Chem.* **13(3)**, 318-334.
20. Calliste, C.A., Le Bail, J.C., Trouillas, P., Pouget, C., Havrioux, G., Chulia, A.J., and Duroux, J.L. (2001). Chalcones: Structural requirements for antioxidant, estrogenic and antiproliferative activities. *Anticancer Res.* **21(6A)**, 3949-3956.
21. Rezk, B.M., Haenen, G.R., van der Vijgh, W.J., and Bast, A. (2002). The antioxidant activity of phloretin: The disclosure of a new antioxidant pharmacophore in flavonoids. *Biochem. Biophys. Res. Commun.* **295(1)**, 9-13.
22. Chen, W.J., Song, J.R., Guo, P., and Wen, Z.Y. (2006). Butein, a more effective antioxidant than  $\alpha$ -tocopherol. *Comp. Theor. Chem.* **763(1-3)**, 161-164..
23. Kozłowski, D., Trouillas, P., Calliste, C.A., Marsal, P., Lazzaroni, R., and Duroux, J.L. (2007). Density functional theory study of the conformational, electronic, and antioxidant properties of natural chalcones. *J. Phys. Chem. A*, **111(6)**, 1138-1145.
24. Todovora, I.T., Batovska, D.I., Stamboliyska, B.A., and Parushev, S.P. (2011). Evaluation of the radical scavenging activity of a series of synthetic hydroxychalcones towards the DPPH radical. *J. Serb. Chem. Soc.* **76(4)**, 491-497.
25. Vazquez-Rodriguez, S., João-Matos, M., Figueroa-Guñez, R., Maya, J.D., Lapiere, M., Olea-Azar, C., Pérez-Cruz, F., Uriarte, E., and Santana, L. (2012). Coumarin-chalcone derivatives as potential antitrypanosomal and antioxidant compounds. In *Proceedings of the 16<sup>th</sup> Int. Electron. Conf. Synth. Org. Chem.* 1-30 November 2012. Sciforum Electronic Conference Series. 1-14.
26. Sashidhara, K.V., Kumar, A., Kumar, M., Sarkar, J., and Sudhir, S. (2010). Synthesis and in vitro evaluation of novel coumarin–chalcone hybrids as potential anticancer agents. *Bioorg. Med. Chem. Lett.* **20(24)**, 7205-7211.
27. Daniel, M. (2006). *Medicinal Plants: Chemistry and Properties*. Enfield, NH, USA: Science Publishers. 157.
28. Sieglar, D.S. (1998b). Flavonoids. *Plant Secondary Metabolism*. Massachusetts, USA: Kluwer Academic Publishers. 151-192



29. Phrutivorapongkul, A., Lipipun, V., Ruangrungsi, N., Kirtikara, K., Nishikawa, K., Maruyama, S., Watanabe, T., and Ishikawa, T. (2003). Studies on the Chemical Constituents of Stem Bark of *Millettia leucantha*: Isolation of New Chalcones with Cytotoxic, Anti-herpes Simplex Virus and Anti-inflammatory Activities. *Chem. Pharm. Bull. (Tokyo)*. **51(2)**, 187-190.
30. Tanaka, T., Iinuma, M., Yuki, K., Fujii, Y., and Muzuno, M. (1992). Flavonoids in root bark of *Pongamia pinnata*. *Phytochemistry*. **31(3)**, 993-998.
31. Salem, M.A.I., Madkour, H.M.F., Soliman, E.S.A., and Mahmoud, N.F.H. (2000). Synthesis of Bactericides via Carbon Nucleophilic Addition on 1,3-Diaryprop-2-enones as Michael Acceptors. *Heterocycles*. **53(5)**, 1129-1143.
32. Tsopmo, A., Tene, M., Kamnaing, P., Ayafor, J.F., and Sterner, O. (1999). A new Diels–Alder-type adduct flavonoid from *Dorstenia barteri*. *J. Nat. Prod.* **62(10)**, 1432- 1434.
33. Ngadjui, B.T., Kouam, S.F., Dongo, E., Kapche, G.W.F., and Abegaz, B.M. (2000). Prenylated flavonoids from the aerial parts of *Dorstenia mannii*. *Phytochemistry*. **55(8)**, 915-919.
34. Fukai, T., Nishizawa, J., and Nomura, T. (1994). Five isoprenoid-substituted flavonoids from *Glycyrrhiza eurycarpa*. *Phytochem.* **35(2)**, 515-519.
35. Hano, Y., Itoh, N., Hanaoka, A., and Nomura, T. (1995). Paratocarpins F-L, seven new isoprenoid-substituted flavonoids from *Paratocarpus venenosa* Zoll. *Heterocycles*. **41(10)**, 2313-2326.
36. Abegaz, B.M., Ngadjui, B.T., Dongo, E., and Tamboue, H. (1998). Prenylated chalcones and flavones from the leaves of *Dorstenia kameruniana*. *Phytochem.* **49(4)**, 1147-1150.
37. Ngameni, B., Touaibia, M., Belkaid, A., Ambassa, P., Watchueng, J., Patnam, R., Ngadjui, B.T., Annabi, B., and Roy, R. (2007). Inhibition of matrix metalloproteinase-2 secretion by chalcones from the twigs of *Dorstenia barteri*. *ARKIVOC*. **2007(9)**, 91-103.
38. Jamil, S., Sirat, H.M., Jantan, I., Aimi, N., and Kitajima, M. (2008). A new prenylated dihydrochalcone from the leaves of *Artocarpus lowii*. *J. Nat. Med.* **62(3)**, 321-324.
39. Shajarahtunnur Jamil (2006). Phytochemical and bioactivities of Malaysia *Artocarpus lowii* King, *A. scortechinii* King, *A. teysmanii* MIQ. Universiti Teknologi Malaysia: Doctor of Philosophy Thesis.

40. Jayasinghe, L., Balasooriya, B. A.I.S., Padmini, W.C., Hara, N., and Fujimoto, Y. (2004). Geranyl chalcone derivatives with antifungal and radical scavenging properties from the leaves of *Artocarpus nobilis*. *Phytochemistry*. **65(9)**, 1287-1290.
41. Wang, L.S., Carmella, S., Keyes, R., Kuo, C.T., Huang, Y.W., Hecht, S.S., and Stoner, G.D. (2012). Anthocyanins and Cancer Prevention. In Sarkar, F.H. (Ed.). *Nutraceuticals and Cancer*, Netherlands: Springer Science. 201-229
42. Sullivan, J. (1998). Anthocyanin. *Carniv. Pl. Newslett.* **27(3)**, 86-88
43. Mann, J. (1987). Secondary metabolism. 2<sup>nd</sup> Ed. USA: Oxford University Press. 173-191.
44. Arora, A., Nair, M.G., and Strasburg, G.M. (1998). Structure-Activity Relationship for Antioxidant Activities of a Series of Flavonoids in a Liposomal System. *Free Radic. Biol. Med.* **24(9)**, 1355-1363.
45. Nowakowska, Z. (2007). A review of anti-infective and anti-inflammatory chalcones. *Eur. J. Med. Chem.* **42(2)**, 125-137.
46. Tsukiyama, R.-I., Katsura, H., Tokuriki, N., and Kobayashi, M. (2002). Antibacterial Activity of Licochalcone A against Spore-Forming Bacteria. *Antimicrob. Agents Ch.* **46(5)**, 1226-1230.
47. Narender, T., Shweta, S., Tanvir, K., Srinivasa Rao, M., Srivastava, K., and Puri, S. K. (2005). Prenylated chalcones isolated from *Crotalaria* genus inhibits in vitro growth of the human malaria parasite *Plasmodium falciparum*. *Bioorg. Med. Chem. Lett.* **15(10)**, 2453-2455.
48. Ram, V.J., Saxena, A.S., Srivastava, S., and Chandra, S. (2000). Oxygenated chalcones and bischalcones as potential antimalarial agents. *Bioorg. Med. Chem. Lett.* **10(19)**, 2159-2161.
49. Troeberg, L., Chen, X., Flaherty, T.M., Morty, R.E., Cheng, M., Hua, H., Springer, C., McKerrow, J.H., Kenyon, G.L., Lonsdale-Eccles, J.D., Coetzer, T.H., and Cohen, F.E. (2000). Chalcone, acyl hydrazide, and related amides kill cultured *Trypanosoma brucei brucei*. *Mol. Med.* **6(8)**, 660-669.
50. Okwu, D.E., and Ukanwa, N. (2010). Isolation and characterization of flavonoids, chalcones and anthocynidines from *Bridelia ferruginea* Benth. *Der Chemica Sinica.* **1(2)**, 21-28.

51. Baviskar, B.A., Baviskar, B., Shiradkar, M.R., Deokate, U.A., and Khadabadi, S.S. (2009). Synthesis and Antimicrobial Activity of Some Novel Benzimidazolyl Chalcones. *J. Chem.* **6(1)**, 196-200.
52. Bohm, B.A. (1998). Introduction to flavonoids. Amsterdam: Harwood Academic. 339-364
53. Daskiewicz, J.B., Comte, G., Barron, D., Di Pietro, A., and Thomasson, F. (1998). Organolithium mediated synthesis of prenylchalcones as potential inhibitors of chemoresistance. *Tetrahedron Lett.* **40(39)**, 7095-7098.
54. Sebti, S., Solhy, A., Tahir, R., Boulaajaj, S., Mayoral, J.A., Faraile, J.M., Kossir, A., and Oumimoun, H. (2001). Calcined sodium nitrate/natural phosphate: an extremely active catalyst for the easy synthesis of chalcones in heterogeneous media. *Tetrahedron Lett.* **42(45)**, 7953-7955.
55. Hamlin, T.A., and Leadbeater, N.E. (2013). Raman spectroscopy as a tool for monitoring mesoscale continuous-flow organic synthesis: Equipment interface and assessment in four medicinally-relevant reactions. *J. Org. Chem.* **9**, 1843-1852.
56. Palleros, D.R. (2004). Solvent-Free Synthesis of Chalcones. *J. Chem. Educ.* **81(9)**, 1345-1347.
57. Petrov, O., Ivanova, Y., & Gerova, M. (2008).  $\text{SOCl}_2/\text{EtOH}$ : Catalytic system for synthesis of chalcones. *Catal. Commun.* **9(2)**, 315-316.
58. Narender, T., and Reddy, K.P. (2007). A simple and highly efficient method for the synthesis of chalcones by using borontrifluoride-etherate. *Tetrahedron Lett.* **48(18)**, 3177-3180.
59. Geissman, T.A., and Fukushima, D.K. (1948). Flavonones and related compounds. V. Oxidation of 2'-hydroxychalcones with alkaline hydrogen peroxide. *J. Am. Chem. Soc.* **70(5)**, 1686-1689.
60. Bennett, M., Burke, A.J., and O'Sullivan, W.I. (1998). Aspects of the Algar-Flynn-Oyamada (AFO) reaction. *Tetrahedron.* **54(34)**, 9911-9911.
61. Bois, F., Beney, C., Boumendjel, A., Mariotte, A.M., Conseil, G., and Di Pietro, A. (1998). Halogenated chalcones with high-affinity binding to P-glycoprotein: potential modulators of multidrug resistance. *Med. Chem.* **41(21)**, 4161-4164.

62. Ivanova, A.B., Batovska, D.I., Todovora, I.T., Stamboliyska, B.A., Serly, J., and Molnar, J. (2011). Comparative study on the MDR reversal effects of selected chalcones. *Int. J. Med. Chem.* **2011**, 1-7.
63. Eddarir, S., Cotelle, N., Bakkour, Y., and Rolando, C. (2003). An efficient synthesis of chalcones based on the Suzuki reaction. *Tetrahedron Lett.* **44(28)**, 5359-5363.
64. Edwards, M.L., Stemerick, D.M., and Sunkara, P.S.J. (1990). Chalcones: a new class of antimitotic agents. *Med. Chem.* **33(7)**, 1948-1954.
65. Liu, M., Wilairat, P., and Go, M.L. (2001). Antimalarial alkoxyated and hydroxylated chalcones: Structure-activity relationship analysis. *Med. Chem.* **44(25)**, 4443-4452.
66. Nielsen, S.F., Christensen, S.B., Cruzciani, G., Kharazmi, A., and Liljefors, T. (1998). Antileishmanial chalcones: statistical design, synthesis, and three-dimensional quantitative structure-activity relationship analysis. *Med. Chem.* **41(24)**, 4819-4832.
67. Lin, Y.M., Zhou, Y., Flavin, M.T., Zhou, L.M., Nie, W., and Chen F.C. (2002). Chalcones and flavonoids as anti-tuberculosis agents. *Bioorg. Med. Chem.* **10(8)**, 2795-2802.
68. Furman, C., Lebeau, J., Fruchart, J., Bernier, J., Duriez, P., Cotelle, N., and Teissier, E. (2001). Di-tert-butylhydroxylated flavonoids protect endothelial cells against oxidized LDL-induced cytotoxicity. *J. Biochem. Mol. Toxicol.* **15(5)**, 270-278.
69. Edward, G., Shahinian, H., Haiduc, I., and Sebe, I. (2011). Synthesis and Characterization of new azo coumarin dyes. *UPB Sci. Bull.* **73(4)**, 154-160.
70. Waksmundzka-Hajnos, M., and Hawrył, M.A. (2008). Application of TLC in the Isolation and Analysis of Coumarins. In Waksmundzka-Hajnos, M., Sherma, J., and Kowalska, T. (Eds.), *Thin Layer Chromatography in Phytochemistry*. Vol. 99. Florida, USA: CRC Press Taylor & Francis Group. 366.
71. IUPAC. (1997). Aglycon (aglycone). In McNaught, A.D., and Wilkinson A., (Eds.), *Compendium of Chemical Terminology*. 2<sup>nd</sup> Ed, Vol. 67. Oxford, UK: Blackwell Scientific Publications. 1307.
72. Glowniak, K., and Widelski, J. (2005). Coumarins: Analysis by TLC. In Cazes J. (Ed), *Encyclopedia of Chromatography*. 2<sup>nd</sup> Ed, Vol. 1. Florida, USA: Taylor & Francis Group. 389.

73. Ahmed, N., and Babu, B.V. (2013). Efficient route to highly functionalized chalcone-based pyranocoumarins via iodine-promoted Michael addition followed by cyclization of 4-hydroxycoumarins. *Synth. Commun.* **43(22)**, 3044-3053.
74. Sieglar, D.S. (1998a). Coumarins. *Plant Secondary Metabolism*. Massachusetts: Kluwer Academic Publishers. 130-138.
75. Ma, T., Liu, L., Xue, H., Li, L., Han, C., Wang, L., Chen, Z., and Liu, G. (2008). Chemical library and structure-activity relationships of 11-demethyl-12-oxo calanolide A analogues as anti-HIV-1 agents. *J. Med. Chem.* **51(5)**, 1432-1446.
76. Appendino, G., Mercalli, E., Fuzzati, N., Arnoldi, L., Stavri, M., Ballero, M., and Maxia, A. (2004). Antimycobacterial coumarins from the sardinian giant fennel (*Ferula communis*). *J. Nat. Prod.* **67(12)**, 2108-2110.
77. Kidane, A.G., Salacinski, H., Tiwari, A., Bruckdorfer, K.F., and Seifalian, A.M. (2004). Anticoagulant and antiplatelet agents: their clinical and device application(s) together with usages to engineer surfaces. *Biomacromolecules.* **5(3)**, 798-813.
78. Kontogiorgis, C., and Hadjipavlou, L.D. (2004). Synthesis and biological evaluation of novel coumarin derivatives with a 7-azomethine linkage. *Bioorg. Med. Chem. Lett.* **14(3)**, 611-614.
79. Sashidhara, K.V., Rosiah, J.N., Kumar, A., Bhatia, G., and Khanna, A.K. (2010). Synthesis of novel benzocoumarin derivatives as lipid lowering agents. *Bioorg. Med. Chem. Lett.* **20(10)**, 3065-3069.
80. Basile, A., Sorbo, S., Spadaro, V., Bruno, M., Maggio, A., Faraone, N., and Rosselli, S. (2009). Antimicrobial and Antioxidant Activities of Coumarins from the Roots of *Ferulago campestris* (Apiaceae). *Molecules.* **14**, 939-952.
81. Perkin, W. H. (1868). On the hydride of aceto-salicyl. *J. Chem. Soc.* **21**, 181-186.
82. Pechmann, H.v., and Duisberg, C. (1883). Ueber die Verbindungen der Phenole mit Acetessigäther. *Ber. Dtsch. Chem. Ges.* **16(2)**, 2119-2128.
83. Maheshwara, M., Siddaiah, V., Lakishmi, G., Damu, V., Rao, Y.K., and Rao, C.V. (2006). A solvent-free synthesis of coumarins via Pechmann condensation using heterogeneous catalyst. *J. Mol. Catal. A: Chem.* **255(1-2)**, 49-52.

84. Hathaway, B.R. (2006). *Barron's Organic Chemistry*. New York, USA: Barron's Educational Series, Inc. 273-275.
85. McDonald, I.M. (2009). Knoevenagel Reaction. In Li, J.J. (Ed.), *Name Reaction for Homologations, Part I*, Hoboken, New Jersey: John Wiley & Sons, Inc. 474-501.
86. Verdia, P., Santamarta, F., and Tojo, E. (2011). Knoevenagel Reaction in [MMIm][MSO<sub>4</sub>]: Synthesis of Coumarins. *Molecules*. **16(6)**, 4379-4388.
87. Musa, M.A., Cooperwood, J.S., and Khan, M.O.F. (2008). A Review of Coumarin Derivatives in Pharmacotherapy of Breast Cancer. *Curr. Med. Chem.* **15(26)**, 2664-2679.
88. Muregi, F.W., and Ishih, A. (2010). Next generation antimalarial drugs: hybrid molecules as a new strategy in drug design. *Drug Dev. Res.* **71(1)**, 20-32.
89. Nepali, K., Sharma, S., Sharma, M., Bedi, P.M.S., and Dhar, K.L. (2014). Rational approaches, design strategies, structure activity relationship and mechanistic insights for anticancer hybrids. *Eur. J. Med. Chem.* **77**, 422-487.
90. Bombardelli, E., and Valenti, P. (2001). 8-(Arylpropenoyl)coumarins as Antiproliferative Agents. Indena, SpA, Italy: PCT.
91. Patel, K., Karthikeyan, C., Moorthy, N.S.H.N., Deora, G.S., Solomon, V.R., Lee, H., and Trivedi, P. (2012). Design, synthesis and biological evaluation of some novel 3-cinnamoyl-4-hydroxy-2H-chromen-2-ones as antimalarial agents. *Med. Chem. Res.* **21(8)**, 1780-1784.
92. Pérez-Cruz, F., Vazquez-Rodriguez, S., Matos, M.J., Herrera-Morales, A., Villamena, F.A., Das, A., Gopalakrishnan, B., Olea-Azar, C., Santana, L., and Uriarte, E. (2013). Synthesis and electrochemical and biological studies of novel coumarin-chalcone hybrid compounds. *J. Med. Chem.* **56(15)**, 6136-6145.
93. Vazquez-Rodriguez, S., Matos, M.J., Santana, L., Borges, E.F., Kachler, S., and Klotz, K.N. (2013). Chalcone-based derivatives as new scaffolds for hA<sub>3</sub> adenosine receptor antagonists. *J. Pharm. Pharmacol.* **65(5)**, 697-703.
94. Choudhary, S., Kini, S.G., and Mubeen, M. (2013). Antioxidant activity of novel coumarin substituted benzothiazole derivatives. *Der Pharma Chem.* **5(4)**, 213-222.

95. Zeng, H. (2010). Human Nutrition-Macronutrients. In *Nutrition Optimization for Health and Longevity*, Bloomington, IN, USA: iUniverse. 11-23.
96. Rolfes, S.R., Pinna, K., and Whitney, E. (2012). Antioxidant Nutrients in Disease Prevention. In Cossio, Y. (Ed.), *Understanding normal and clinical nutrition*, Belmont, CA, USA: Cengage Learning. 376-383.
97. Kaur, C., and Kapoor, H.C. (2001). Antioxidants in fruits and vegetables-the Millenium's health. *Int. J. Food Sci. Tech.* **36(7)**, 703-725.
98. Nigam, D. (2015). Free Radicals and Oxidative Stress in Neurodegenerative Disorders. In Rani, V. and Yadav, U.C.S. (Eds.), *Free Radicals in Human Health and Disease*, India: Springer. 143-159.
99. Newman, D.J., Cragg, G.M., and Snader, K.M. (2003). Natural Products as sources of new drugs over the period 1981-2002. *J. Nat. Prod.* **66**, 1022-1037.
100. Shaikh, R., Pund, M., Dawane, A., and Iliyas, S. (2014). Evaluation of Anticancer, Antioxidant an Possible Anti-inflammatory Properties of Selected Medicinal Plants Used in Indian Traditional Medication. *J. Trad. Complement. Med.* **4(4)**, 253-257.
101. McLain, L. (2013). Antioxidant: Eat All Your Colors! In *Natural Antioxidants: Discover The Cutting Edge Antioxidant Program That Will Make You Healthy, Thin, And Beautiful As You Learn Antioxidant & Anticancer, Antioxidant Weight Loss And The Benefits Antioxidant Foods*. Canada: JDSims Publishing. 16.
102. Prabhakar, E.R, Suchitra, M., Seshadri, V.R., Aparna, R.B., and Sivakumar, V., (2010). Ferric Reducing Ability of Plasma and Lipid Peroxidation in Hemodialysis Patients: Intradialytic Changes. *Nephrourol. Mon.* **2(3)**, 414-421.
103. Kaushik, A., Chanderesh, J., Kaushik, J.J., Zeray, R., Ambesajir, A., and Beyene, L. (2012). FRAP (Ferric reducing ability of plasma) assay and effect of *Diplazium esculentum* (Retz) Sw. (a green vegetable of North India) on central nervous system. *Indian J. Nat. Prod. Resour.* **3(2)**, 228-231.
104. Re, R., Pellegrini, N., Proteggente, A., Pannala, A., Yang, M., and Rice-Evans, C. (1999). Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radic. Biol. Med.* **26(9-10)**, 1231-1237.
105. Antolovich, M., Prenzler, P.D., Patsalides, E., McDonald, S., and Robards, K. (2002). Methods for testing antioxidant activity. *R. Soc. Chem.* **127**, 183-198.

106. Miller, N.J., Sampson, J., Candeias, L.P., Bramley, P.M., and Rece-Evans, C.A. (1996). Antioxidant activities of carotenes and xanthophylls. *FEBS Lett.* **384(3)**, 240-242.
107. Garcia, E.J., Oldoni, T.L.C., Alencar, S.M., Reis, A., Loguercio, A.D., and Grande, R.H.M. (2012). Antioxidant Activity by DPPH Assay of Potential Solutions to be Applied on Bleached Teeth. *Braz. Dent. J.* **23(1)**, 22-27.
108. Blois, M. S. (1958). Antioxidant determinations by the use of a stable free radical. *Nature.* **181(4617)**, 1199-1200.
109. Bonina, F., Puglia, C., Tomaino, A., Saija, A., Mulinacci, N., Romani, A., and Vincieri, F.F. (2000). In-vitro antioxidant and in-vivo photoprotective effect of three lyophilized extracts of *Sedum telephium* L. leaves. *J. Pharm. Pharmacol.* **52(10)**, 1279-1285.
110. Moreno, C.S., Larrauri, J.A., and Fulgencio, S.C. (1999). A procedure to measure the antiradical efficiency of polyphenols. *J. Sci. Food Agr.* **76(2)**, 270-276.
111. Stanchev, S., Penkova, A., Retailleau, C., Avdeev, G., and Manolov, I. (2008). Crystal structure of 3-Acetyl-6-methoxy-2H-1-benzopyran-2-one. *Anal. Sci.* **24**, X183-X184.
112. Shahwar, D., Raza, M. A., Bukhari, S., & Bukhari, G. (2012). Ferric reducing antioxidant power of essential oils extracted from Eucalyptus and Curcuma species. *Asian Pac. J. Trop. Biomed.*, **2(3)**, S1633-S1636.
113. Benzie, I.F.F. (1996). An automated, specific, spectrophotometric method for measuring ascorbic acid in plasma (EFTSA). *Clin. Biochem.* **29**, 111-116.
114. Liu, T.Z., Chin, N., Kiser, M.D. and Bigler, W.N. (1982). Specific spectrophotometry of ascorbic acid in serum or plasma by use of ascorbate oxidase. *Clin. Chem.* **28(11)**, 2225-2228.
115. Benzie, I.F.F. and Strain, J.J. (1996). The Ferric Reducing Ability of Plasma (FRAP) as measure of "Antioxidant Power": The FRAP Assay. *Anal. Biochem.* **239**, 70-76.
116. Miller, D.D. (1996). Mineral. In Fennema, O.R. (Ed.). *Food Chemistry*. New York, USA: Marcel Dekker. 618-649.
117. MacDonald-Wicks L.K., Wood, L.G., and Garg M.L. (2006). Methodology for the determination of biological antioxidant capacity in vitro: a review. *J. Agric. Food Chem.* **86**, 2046-2056.



118. Molyneux, P. (2004). The use of the stable free radical diphenylpicrylhydrazyl (DPPH) for estimating antioxidant activity. *Songklanakar J. Sci. Technolo.* **26(2)**, 211-219.
- 119 Foti, M., Piatelli, M., Baratta, M.T., and Ruberto,G. (1996). Flavonoids, Coumarins, and Cinnamic Acids as Antioxidants in a Micellar System. Structure-Activity Relationship. *J. Agric. Food Chem.*, **44**, 497-501.
- 120 Channarong, S., Jutiviboonsuk, A., and Korsanan, S. (2012). Total Reducing Antioxidant Capacity of Thai Herbal Aromatic Powder (Ya-Hom) Measured by FRAP Assay. *Thai Pharm. Health Sci. J.* **7(3)**, 111-114.
- 121 Zou, Y., Chang, S.K.C., Gu, Y., and Qian, S.Y. (2011). Antioxidant Activity and Phenolic Composition of Lentils (*Lens culinaris* var. Morton) Extracts and its Fractions. *J. Agri. Food Chem.* **59(6)**, 2268-2276.
- 122 Fu, R., Zhang, Y., Guo, Y., Liu, F., and Chen F. (2014). Determination of Phenolic Content and Antioxidant of Extracts of *Jatropha curcas* L. Seed Shell, A By-Product A New Source of Natural Antioxidant. *Ind. Crops. Prod.* **58**, 265-270.