

LIGNOCELLULOLYTIC ENZYMES BY *Aspergillus* sp. A1 AND *Bacillus* sp. B1
ISOLATED FROM GUT OF *Bulbitermes* sp. IN SOLID STATE FERMENTATION
USING SAWDUST AS SUBSTRATE

NORATIQAH BINTI KAMSANI

UNIVERSITI TEKNOLOGI MALAYSIA

LIGNOCELLULOLYTIC ENZYMES BY *Aspergillus* sp. A1 AND *Bacillus* sp. B1
ISOLATED FROM GUT OF *Bulbitermes* sp. IN SOLID STATE FERMENTATION
USING SAWDUST AS SUBSTRATE

NORATIQA BINTI KAMSANI

A thesis submitted in fulfilment of the
requirements for the award of the degree of
Doctor of Philosophy (Bioscience)

Faculty of Biosciences and Medical Engineering
Universiti Teknologi Malaysia

JANUARY 2017

*Specially dedicated to supportive families and friends who had been an inspiration to me
to be a better person*

ACKNOWLEDGEMENT

In the name of Allah, Most Gracious, Most Merciful

Alhamdulillah. Thanks to Allah SWT, whom with His willing giving me the opportunity to complete this study.

I am heartily thankful to my supervisor, Assoc. Prof. Dr. Madihah Md. Salleh, co-supervisors, Dr. Adibah Yahya and Dr. Chong Chun Shiong, whose encouragement, guidance and support from the initial to the final level enabled me to develop an understanding of the subject. Their invaluable help of constructive comments and suggestions throughout experimental and reporting of findings have contributed to the success of this research. The compliment also goes to my external co-supervisor, Dr. Musaalbakri Abdul Manan, for his thoughts and comments.

I would also like to show my appreciation to the Ministry of Higher Education (MOHE) and University for financial assistances.

My sincere appreciation to all my colleagues in Biorefinery Technology Research Laboratory, Dr. Ang Siow Kuang, Dr. Rachmawaty Muchtar, Puan Huszalina Hussin, Shankar A/L Ramanathan, Ahmad Fawwaz Mohd Raji, Mohd Roslan Ikubar, Chew Yue Ming, Rohaya Mohd Noor, Zul and Syifaa for their kindness and moral support during my study. My appreciation also goes to Puan Fatimah for her generous assistance throughout this study.

Last but not least, my deepest gratitude goes to my beloved parents; Mr Kamsani Suratman and Mrs Azizah Abu Bakar and also to my sisters and brother, Nana, Yan and Emy for their endless love, prayers and encouragement. To those who indirectly contributed in this research (especially for the musicians, café cooker and Nescafe inventor) your kindness and inventions means a lot to me. Thank you very much.

ABSTRACT

Sawdust is one of the common lignocellulosic waste biomass produced during the process of planning mills, moulding plants and furniture manufacturing. In practice, the sawdust is discarded in landfill areas, causing dust and dirt pollution in nearby localities. Therefore, the need to find an efficient and practical approach to revalorize sawdust as a starting raw material in the production of lignocellulolytic enzymes is essential as a way to manage and turn the residues into value added products. Prospecting for efficient degrading lignocellulose microorganisms is crucial to facilitate the process of lignocellulolytic enzymes production from the lignocellulosic biomass. This study aimed to exploit microorganisms isolated from gut of termite *Bulbitermes* sp. in producing lignocellulolytic enzymes under solid-state fermentation (SSF) system by using untreated sawdust as substrate. Seventeen bacterial and five fungal with positive lignocellulolytic enzymes activities were successfully isolated from the gut of two hundred termites. Four isolates identified as *Aspergillus* sp. A1, *Bacillus* sp. B1, *Bacillus* sp. B2 and *Brevibacillus* sp. Br3 were selected for further characterization. Among the isolates, *Aspergillus* sp. A1 showed highest activities of lignin peroxidase (LiP) (729.12 U/g) and β -glucosidase (22.97 U/g). The highest activities of endoglucanase (138.77 U/g) and manganese peroxidase (MnP) (47.73 U/g) were recorded in *Bacillus* sp. B1. The *Bacillus* sp. B2 produced the highest activities of exoglucanase (32.16 U/g) and laccase (71.18 U/g). The highest xylanase activity (104.96 U/g) was observed in *Brevibacillus* sp. Br3. The production of endoglucanase, β -glucosidase, xylanase, LiP and laccase were approximated 17–93% higher in co-culture compared to individual culture. Compared to other di-, tri- and quad-mixed culture, *Aspergillus* sp. A1 (A1) and *Bacillus* sp. B1 (B1) co-culture produced the highest lignocellulolytic enzymes activities (endoglucanase, 190.1; exoglucanase, 13.5; β -glucosidase, 33.7; xylanase, 202.5; LiP, 713.5; MnP, 23.3 and laccase, 52.1 U/g). The interaction between A1 and B1 is not antagonistic. Study on the effect of SSF operational variables showed that the use of unsieved sawdust produced significantly higher activities of exoglucanase, xylanase, LiP and laccase compared to that of sieved sawdust. In addition, temperature, pH and moisture content significantly impacted lignocellulolytic enzymes production. In comparing to control, moistening the unsieved sawdust with Mandel basal medium (pH 8) to 1:2.5 (solid:moisture) ratio, and incubation at 35 °C for 9 days produced 1.2–49.4 fold higher lignocellulolytic enzymes activities. Endoglucanase, β -glucosidase and xylanase could be classified as moderately thermostable enzymes with better stability in acidic pH range. Meanwhile, ligninases possessed thermophilic and alkaliphilic characteristics. The co-culture produced 1.9–11.8 fold higher reducing sugars than those yielded by single cultures in the enzymatic degradation of sawdust. The use of co-culture enzymes also produced 3.6–85.4% higher reducing sugars as well as 1.3–2.3 times higher raffinose, cellobiose, maltose, glucose and xylose concentrations compared to that of commercial cellulase (Celluclast) solution. As conclusion, this work has generated a microbial co-culture that could be used for improved lignocellulolytic enzymes and reducing sugars production using untreated sawdust as substrate.

ABSTRAK

Hampas kayu merupakan salah satu sisa biojisim lignoselulosa yang dihasilkan semasa proses pengilangan terancang, loji pengacuan dan pembuatan perabot. Kebiasaannya, hampas kayu dibuang di kawasan pelupusan sampah, mengakibatkan pencemaran habuk dan debu di kawasan setempat yang berhampiran. Oleh itu, mencari pendekatan yang efisien dan praktikal untuk meningkatkan nilai hampas kayu adalah penting sebagai cara untuk mengurus dan menukar sisa ini kepada produk berguna dengan menggunakannya sebagai bahan asas dalam penghasilan enzim lignoselulolitik. Pengenalpastiaan mikroorganisma yang boleh menguraikan lignoselulosa secara efisien adalah penting untuk memudahkan proses penghasilan enzim lignoselulolitik dari biojisim lignoselulosik. Matlamat kajian ini adalah untuk mengeksploitasi mikroorganisma yang dipencilkan daripada usus anai-anai *Bulbitermes* sp. dalam menghasilkan enzim lignoselulolitik di bawah sistem penapaian keadaan pepejal (SSF) menggunakan hampas kayu yang tidak dirawat sebagai substrat. Tujuh belas bakteria dan lima kulat dengan aktiviti enzim lignoselulolitik yang positif telah berjaya dipencilkan daripada usus dua ratus anai-anai. Empat pencilan dikenalpasti sebagai *Aspergillus* sp. A1, *Bacillus* sp. B1, *Bacillus* sp. B2 dan *Brevibacillus* sp. Br3 telah dipilih untuk pencirian yang lebih lanjut. Diantara pencilan-pencilan tersebut, *Aspergillus* sp. A1 menunjukkan aktiviti lignin peroksidase (LiP) (729.12 U/g) dan β -glukosidase tertinggi (22.97 U/g). Aktiviti endoglukanase (138.77 U/g) dan manganese peroksidase (MnP) (47.73 U/g) tertinggi telah direkodkan oleh *Bacillus* sp. B1. *Bacillus* sp. B2 menghasilkan aktiviti eksoglukanase (32.16 U/g) dan lakase (71.18 U/g) tertinggi. Aktiviti xilanase tertinggi (104.96 U/g) dicatatkan oleh *Brevibacillus* sp. Br3. Penghasilan endoglukanase, β -glukosidase, xilanase, LiP dan lakase adalah dianggarkan 17–93% lebih tinggi dalam kultur bersama berbanding dengan kultur tunggal. Perbandingan antara dwi-, tri- dan kuad-kultur bercampur menunjukkan, kultur bersama *Aspergillus* sp. A1 (A1) dan *Bacillus* sp. B1 (B1) menghasilkan aktiviti enzim lignoselulolitik tertinggi (endoglukanase, 190.1; eksoglukanase, 13.5; β -glukosidase, 33.7; xilanase, 202.5; LiP, 713.5; MnP, 23.3 dan lakase, 52.1 U/g). A1 dan B1 mempunyai hubungan tidak antagonis. Kajian mengenai kesan parameter operasi SSF menunjukkan hampas kayu yang tidak diayak menghasilkan aktiviti eksoglukanase, xilanase, LiP dan lakase jauh lebih tinggi berbanding dengan hampas kayu yang diayak. Selain itu, suhu, pH dan kandungan kelembapan memberi kesan yang signifikan terhadap penghasilan enzim lignoselulolitik. Hampas kayu tidak diayak yang dilembapkan dengan medium Mandel asas (pH 8) kepada nisbah 1:2.5 (pepejal:kelembapan) pada suhu 35 °C untuk 9 hari menghasilkan 1.2–49.4 kali ganda lebih tinggi aktiviti enzim lignoselulolitik berbanding dengan eksperimen kawalan. Endoglukanase, β -glukosidase dan xilanase boleh dikategorikan sebagai enzim stabil haba sederhana dan mereka juga lebih stabil dalam pH berasid. Manakala, ligninase mempunyai ciri-ciri stabil haba dan stabil alkali. Kultur bersama menghasilkan 1.9–11.8 kali ganda lebih tinggi gula terturun daripada yang dihasilkan oleh kultur tunggal dalam proses penguraian berenzim hampas kayu. Penggunaan enzim kultur bersama juga menghasilkan 3.6–85.4% lebih tinggi gula terturun dan juga 1.3–2.3 kali ganda kepekatan rafinosa, selobiosa, maltosa, glukosa dan xilosa berbanding dengan menggunakan larutan enzim selulase komersial (Celluclast). Kesimpulannya, kajian ini telah menghasilkan mikrob kultur bersama yang boleh digunakan untuk meningkatkan penghasilan enzim lignoselulolitik dan gula terturun menggunakan hampas kayu yang tidak dirawat sebagai substrat.

TABLE OF CONTENTS

CHAPTER	TITLE	PAGE
	DECLARATION	ii
	DEDICATION	iii
	ACKNOWLEDGEMENT	iv
	ABSTRACT	v
	ABSTRAK	vi
	TABLE OF CONTENTS	vii
	LIST OF TABLES	xvii
	LIST OF FIGURES	xxii
	LIST OF ABBREVIATIONS	xxviii
	LIST OF APPENDICES	xxx
1	INTRODUCTION	1
	1.1 Background of Research	1
	1.2 Objectives	6
	1.3 Scope of Research	7
	1.4 Significance of the Research	8
	1.5 Thesis Organization	9
2	LITERATURE REVIEW	11
	2.1 Lignocellulosic Biomass	11
	2.2 The Composition of Lignocellulosic Material	13

2.2.1 Cellulose	14
2.2.2 Hemicellulose	16
2.2.3 Lignin	18
2.3 Selection of Lignocellulosic Biomass for Production of Biobased Products	20
2.3.1 Sawdust as Raw Material for Production of High Value Products	21
2.4 Lignocellulose Degradation	24
2.4.1 Acid Degradation	27
2.4.2 Enzymatic Degradation: (Hemi) cellulases	29
2.4.3 Lignin Degrading Enzymes: Ligninases	33
2.5 Lignocellulolytic Enzymes Microbial Producer	36
2.6 Sources of Lignocellulolytic Degrading Microorganisms	42
2.6.1 The Termite	43
2.6.2 Lignocellulolytic Microorganisms from The Gut of Termite	45
2.7 Submerged Fermentation (SmF)	50
2.8 Solid-State Fermentation (SSF)	51
2.8.1 Advantages and Challenges in SSF	54
2.9 SSF Process Conditions for Lignocellulolytic Enzymes Production	56
2.9.1 Particle Size	56
2.9.2 Temperature	57
2.9.3 pH	58
2.9.4 Moisture Content	59
2.10 Co-Culture of Lignocellulolytic Enzymes Microbial Producers	62
2.11 Application of Lignocellulolytic Enzymes	66
2.11.1 Cellulases	66
2.11.2 Xylanase	68

2.11.3	Ligninases	69
3	MATERIALS AND METHODS	72
3.1	Research Design	72
3.1.1	Experimental Design	74
3.2	Collection of Termites	76
3.3	Isolation of Microorganisms	76
3.4	Inocula Preparation	77
3.5	Screening of Lignocellulolytic Enzymes Microbial Producers	78
3.5.1	Qualitative Screening	78
3.6	Solid-State Fermentation (SSF) of Sawdust	80
3.6.1	Solid Substrate	80
3.6.2	Inoculum Preparation and Inoculation	81
3.6.3	Enzymes Production in SSF	81
3.6.4	Enzymes Extraction	83
3.7	Reducing Sugar Assay	83
3.8	Enzymes Assay	84
3.8.1	Endoglucanase	84
3.8.2	Exoglucanase	84
3.8.3	β -glucosidase	85
3.8.4	Xylanase	85
3.8.5	Lignin Peroxidase (LiP)	86
3.8.6	Manganese Peroxidase (MnP)	86
3.8.7	Laccase	87
3.9	Protein Assay	87
3.10	Glucosamine Assay	88

4	CULTIVATION AND SELECTION OF LIGNOCELLULOLYTIC MICROORGANISMS FROM THE GUT OF <i>Bulbitermes</i> sp. TERMITES IN SOLID-STATE FERMENTATION OF CHEMICALLY UNTREATED SAWDUST	89
	4.1 Introduction	89
	4.2 Materials and Methods	91
	4.2.1 Collection of Termites	91
	4.2.2 Isolation of Lignocellulolytic Enzymes Producer from <i>Bulbitermes</i> sp. Termite Gut	91
	4.2.3 Gram Staining	91
	4.2.4 Preparation of Bacterial and Fungal Inocula	92
	4.2.5 Qualitative Screening	92
	4.2.6 Substrate Procurement	92
	4.2.7 Quantitative Screening	93
	4.2.8 Enzyme Assays	93
	4.2.9 Identification of Selected Lignocellulolytic Enzymes-Producing Microorganisms	93
	4.2.9.1 DNA Extraction	93
	4.2.9.2 Gel Electrophoresis	95
	4.2.9.3 Polymerase Chain Reaction (PCR)	95
	4.2.9.4 Carbon source Utilization Pattern of Selected Enzymes-Producing Microorganisms	96
	4.2.10 Preparation of Different Cellular Fractions for Enzyme Distribution Studies	97
	4.3 Results and Discussion	98
	4.3.1 Isolation of Microorganisms from Termite Gut	98
	4.3.2 Qualitative Screening for Lignocellulolytic Microorganisms	98

4.3.3	Quantitative Screening for Lignocellulolytic Microorganisms Under SSF condition	102
4.3.3.1	Cellulases Activities	102
4.3.3.2	Xylanase Activities	104
4.3.3.3	Ligninases Activities	106
4.3.4	Selection of Lignocellulolytic Microorganisms for Further Identification and Characterization Study	108
4.3.4.1	Screening for Intracellular Lignocellulolytic Enzymes Production in Selected Microorganisms	116
4.3.5	Comparative Studies	118
4.4	Conclusion	120
5	PRODUCTION OF LIGNOCELLULOLYTIC ENZYMES BY CO-CULTURES OF SELECTED MICROORGANISMS FROM <i>Bulbitermes</i> sp. TERMITE GUT IN SOLID-STATE FERMENTATION OF UNTREATED SAWDUST	121
5.1	Introduction	121
5.2	Materials and Methods	123
5.2.1	Microorganisms and Inocula Preparation	123
5.2.2	Compatibility Test	123
5.2.3	Enzymes Production in Solid-State Fermentation (SSF) of Sawdust	123
5.2.4	Enzymes Assay	124
5.2.5	Statistical Analysis	124
5.3	Results and Discussion	124
5.3.1	Compatibility Test	124
5.3.2	Lignocellulolytic Enzymes Production in Solid- State Fermentation System	125

5.3.2.1	Single Culture	125
5.3.2.2	Co-Culture	126
5.3.2.2.1	Cellulases	126
5.3.2.2.2	Xylanase	130
5.3.2.2.3	Ligninases	131
5.3.2.3	Volumetric Productivity	135
5.4	Conclusion	137
6	CO-CULTIVATION OF <i>Aspergillus</i> sp. A1 AND <i>Bacillus</i> sp. B1 FOR LIGNOCELLULOLYTIC ENZYMES PRODUCTION IN SOLID-STATE FERMENTATION	138
6.1	Introduction	138
6.2	Materials and Methods	139
6.2.1	Strains of Lignocellulolytic Microorganisms	139
6.2.2	Inocula Preparation	139
6.2.3	Production of Lignocellulolytic Enzymes by Single and Co-cultures under SSF Condition	140
6.2.4	Enzyme Assays	140
6.2.5	Glucosamine Assay	140
6.2.6	Scanning Electron Microscopy (SEM)	140
6.2.7	Compatibility Tests	141
6.2.8	Exopolysaccharide (EPS) Determination	141
6.2.8.1	EPS Extraction	141
6.2.8.2	EPS Quantification	142
6.2.9	Spore Staining with Malachite Green-Safranin	142
6.2.10	Sodium Dodecyl Sulfate Polyacrylamide Gel Electrophoresis (SDS-PAGE)	143
6.2.11	Protein Assay	143
6.3	Results and Discussion	144

6.3.1	Effect of Inocula on Lignocellulolytic Enzymes Production	144
6.3.2	Compatibility Evaluation	146
6.3.3	Endospore and Exopolysaccharides Production	151
6.3.4	Effect of Co-Cultivation on Overall Protein Profiles	156
6.3.5	Effects of Single and Co-Culture on Lignocellulolytic Enzymes System	158
6.3.5.1	Cellulases	158
6.3.5.2	Xylanase	164
6.3.5.3	Ligninases	166
6.3.6	Effects of Single and Co-Culture on N-Acetyl-D-Glucosamine (NAG) Production	173
6.4	Conclusion	175
7	EFFECTS OF OPERATING CONDITIONS ON LIGNOCELLULOLYTIC ENZYMES PRODUCTION IN SOLID-STATE FERMENTATION OF SAWDUST	177
7.1	Introduction	177
7.2	Materials and Methods	178
7.2.1	Preparation of Solid Substrate	178
7.2.2	Microorganisms and Inocula Preparation	178
7.2.3	Lignocellulolytic Enzymes Production by Co-Culture under solid-state fermentation (SSF) condition	179
7.2.4	Enzyme Assays	179
7.2.5	Glucosamine Assay	179
7.2.6	Determination of Moisture Content	180
7.2.7	Improving Process Parameters for Lignocellulolytic Enzymes Production under SSF Using One-Factor-At-a-Time (OFAT) Method	180

7.2.8	Statistical Analysis	181
7.3	Results and Discussion	181
7.3.1	Effect of Sawdust Particle Size on Lignocellulolytic Enzymes Production	181
7.3.2	Effect of Incubation Temperature on Lignocellulolytic enzymes Production	186
7.3.3	Effect of initial medium pH on Lignocellulolytic Enzymes Production	190
7.3.4	Effect of Moisture Content on Lignocellulolytic Enzymes Production	194
7.3.5	Comparison of Different SSF Conditions for Lignocellulolytic Enzymes Production	198
7.3.6	Changes in the Activities of Lignocellulolytic Enzymes	200
7.4	Conclusion	207
8	CHARACTERIZATION OF LIGNOCELLULOLYTIC ENZYMES PRODUCED BY <i>Aspergillus</i> sp. A1 AND <i>Bacillus</i> sp. B1 CO-CULTURE USING UNTREATED SAWDUST AS SUBSTRATE UNDER SOLID-STATE FERMENTATION CONDITION	209
8.1	Introduction	209
8.2	Materials and Methods	210
8.2.1	Preparation of Solid Substrate	210
8.2.2	Microorganisms and Inocula Preparation	210
8.2.3	Lignocellulolytic Enzymes Production by Co- Culture under Solid-State Fermentation (SSF) condition	210
8.2.4	Enzyme Assays	211
8.2.5	Characterization of the Crude Lignocellulolytic Enzymes	211

8.2.5.1	Optimum Temperature and Thermal Stability	211
8.2.5.2	Optimum pH and pH stability	211
8.2.6	Sodium Dodecyl Sulfate Polyacrylamide Gel Electrophoresis (SDS-PAGE) and Zymogram Analysis	212
8.3	Results and Discussion	213
8.3.1	Optimum Temperature and Thermal Stability of Lignocellulolytic Enzymes	213
8.3.2	Optimum pH and pH Stability of Lignocellulolytic Enzymes	223
8.3.3	SDS-PAGE and Zymogram of Crude Lignocellulolytic Enzymes	234
8.4	Conclusion	236
9	DEGRADATION OF SAWDUST BASED ON SSF STRATEGY FOR PRODUCTION OF REDUCING SUGARS	238
9.1	Introduction	238
9.2	Materials and Methods	242
9.2.1	Microorganisms and Inoculum Preparation	242
9.2.2	Sawdust-Based Biorefining Strategy	242
9.2.2.1	Solid-State Fermentation (SSF)	242
9.2.2.2	Enzymatic Degradation	243
9.2.3	Analytical Methods	244
9.2.3.1	Enzyme Assays	244
9.2.3.2	Protein	244
9.2.3.3	Analysis of Sawdust Sample Composition	244
9.2.3.3.1	Determination of NDF	245
9.2.3.3.2	Determination of ADF	245
9.2.3.3.3	Determination of ADL	245

9.2.3.4	Fourier Transform Infrared (FTIR)	246
9.2.3.5	Scanning Electron Microscopy (SEM)	246
9.2.3.6	Reducing Sugar Assay	246
9.2.3.7	High Performance Liquid Chromatography (HPLC)	247
9.2.3.8	Total Phenolic and Lignin Content	248
9.2.3.9	Statistical Analysis	248
9.3	Results and Discussion	249
9.3.1	Lignocellulolytic Enzymes Production by Single and Co-Culture of <i>Aspergillus</i> sp. A1 and <i>Bacillus</i> sp. B1	249
9.3.2	Compositional Analysis of Sawdust after Solid- State Fermentation (SSF)	251
9.3.3	Analysis of Sawdust Chemical Structure	253
9.3.4	Microscopic Analysis	257
9.3.5	Enzymatic Degradation of Fermented Sawdust	259
9.3.5.1	Total Phenolic Content and Lignin Concentration	267
9.4	Conclusion	270
10	CONCLUSIONS AND RECOMMENDATIONS	272
10.1	Conclusions	272
10.2	Recommendations	275
	REFERENCES	277
	Appendices A - E	336-349

LIST OF TABLES

TABLE NO.	TITLE	PAGE
2.1	Quantity of biomass produced in Malaysia in 2007	12
2.2	Chemical compositions of lignocellulosic biomass in Malaysia	13
2.3	The comparison of crystalline and amorphous structure of cellulose	15
2.4	Proximate composition of sawdust	22
2.5	Sawdust-based product derived from physical-chemical process	23
2.6	Bioconversion processes of sawdust into various value added products	25
2.7	Reports on the use of dilute acid degradation to different type of lignocellulosic biomass	28
2.8	Enzymes involved in the degradation of complex heteroarabinoxylans and galactoglucomannan structure	32
2.9	Mechanism of lignin biodegradation process	36
2.10	Cellulases producing microorganisms	38
2.11	Xylanase producing microorganisms	39
2.12	Ligninases producing microorganisms	41
2.13	Insects with reported lignocellulolytic microorganisms inside their gut	42

2.14	Lignocellulolytic microorganisms isolated from the gut of termites	48
2.15	Lignocellulosic biomass used as a substrate for production of lignocellulolytic enzymes in SSF and SmF	53
2.16	Advantages and disadvantages of SSF over SmF	54
2.17	Process conditions applied in SSF for lignocellulolytic enzymes production by fungi and bacteria	61
2.18	Compilation of lignocellulolytic enzyme production in fermentation employing single and co-cultures cultivation on different substrate	65
2.19	Application of cellulases	67
2.20	Application of xylanase	69
2.21	Application of ligninases	70
3.1	Composition of Medium 1	77
3.2	Composition of Medium 2	77
3.3	Composition of CMC agar plates	79
3.4	Composition of Birchwood xylan agar plates	79
3.5	Composition of lignin agar plates	80
3.6	Composition of Production Medium	82
3.7	Trace elements	82
4.1	Summary of primers for PCR amplification	96
4.2	Characteristics of thirty bacterial and seven fungal isolates from the gut of <i>Bulbitermes</i> sp. termites	99
4.3	Qualitative screening for cellulolytic, xylanolytic and ligninolytic activities	100
4.4	Species of bacteria and fungi determined by amplification of 16S rRNA and ITS primer pairs respectively	111
4.5	Sole carbon sources utilization patterns of <i>Aspergillus</i> sp. A1, <i>Bacillus</i> sp. B1, <i>Bacillus</i> sp. B2 and <i>Brevibacillus</i> sp. Br3	115

4.6	Highest enzymatic activities detected in different fractions of <i>Aspergillus</i> sp. A1, <i>Bacillus</i> sp. B1, <i>Bacillus</i> sp. B2 and <i>Brevibacillus</i> sp. Br3	117
4.7	Cellulases, xylanase and ligninases production from bacteria and fungi isolated from/associated with the guts of termite under different substrate and fermentation system	119
5.1	Enzymes produced by <i>Aspergillus</i> sp. A1, <i>Bacillus</i> sp. B1, <i>Bacillus</i> sp. B2 and <i>Brevibacillus</i> sp. Br3 in SSF using saw dust as substrate	126
5.2	Comparison of volumetric productivity of cellulases, xylanase and ligninases from different fungal co-cultures under SSF	136
6.1	Comparison of growth (mg NAG/g) and lignocellulolytic enzyme activities (U/g) between single and co-culture after 9 days of SSF	145
6.2	Volumetric productivity of endoglucanase by single and co-culture of <i>Aspergillus</i> sp. A1 and <i>Bacillus</i> sp. B1	160
6.3	Volumetric productivity of exoglucanase by single and co-culture of <i>Aspergillus</i> sp. A1 and <i>Bacillus</i> sp. B1	162
6.4	Volumetric productivity of β -glucosidase by single and co-culture of <i>Aspergillus</i> sp. A1 and <i>Bacillus</i> sp. B1	164
6.5	Volumetric productivity of xylanase by single and co-culture of <i>Aspergillus</i> sp. A1 and <i>Bacillus</i> sp. B1	166
6.6	Volumetric productivity of LiP by single and co-culture of <i>Aspergillus</i> sp. A1 and <i>Bacillus</i> sp. B1	168
6.7	Volumetric productivity of MnP by single and co-culture of <i>Aspergillus</i> sp. A1 and <i>Bacillus</i> sp. B1	170
6.8	Volumetric productivity of laccase by single and co-culture of <i>Aspergillus</i> sp. A1 and <i>Bacillus</i> sp. B1	172
6.9	Rate of N-acetyl-D-glucosamine production by single and co-culture of <i>Aspergillus</i> sp. A1 and <i>Bacillus</i> sp. B1	174

7.1	Variation of process factors for lignocellulolytic enzymes production	181
7.2	Effect of sawdust particle sizes on NAG and final moisture content	186
7.3	Effect of incubation temperature on NAG and final moisture content	190
7.4	Effect of pH on NAG and final moisture content	193
7.5	Effect of initial moisture content on NAG and final moisture content	197
7.6	Lignocellulolytic enzymes activities in different SSF condition by A1B1 co-culture	199
8.1	Optimum temperature of lignocellulolytic enzymes from <i>Aspergillus</i> sp. A1 and <i>Bacillus</i> sp. B1 co-culture grown in SSF of sawdust	214
8.2	Temperature stability for lignocellulolytic enzymes of <i>Aspergillus</i> sp. A1 and <i>Bacillus</i> sp. B1 grown under SSF of sawdust .	220
8.3	Optimum pH of lignocellulolytic enzymes from <i>Aspergillus</i> sp. A1 and <i>Bacillus</i> sp. B1 co-culture grown under SSF of sawdust	223
8.4	pH stability for lignocellulolytic enzymes of <i>Aspergillus</i> sp. A1 and <i>Bacillus</i> sp. B1 grown in SSF of sawdust	230
8.5	Comparison of optimum temperature, pH and stability of cellulases, xylanase and ligninases produced by lignocellulolytic fungal and bacteria.	233
8.6	Molecular mass of endoglucanase, xylanase and laccase from <i>Aspergillus</i> sp. A1 and <i>Bacillus</i> sp. B1 co-culture.	236
9.1	Retention time of sugar standards	247
9.2	Lignocellulolytic enzymes activities produced by single and co-cultures of <i>Aspergillus</i> sp. A1 and <i>Bacillus</i> sp. B1 under SSF condition	250

9.3	Characteristics of the band assignments and wavenumbers in FTIR analysis	254
9.4	Ratios of the intensity of lignin, cellulose and hemicellulose bands for fermented and non-fermented samples before and after SSF	257
9.5	Total reducing sugar produced during enzymatic degradation of fermented and non-fermented sawdust using commercial cellulase of Celluclast and on site-crude enzyme extract from single and co-cultures of <i>Aspergillus</i> sp. A1 and <i>Bacillus</i> sp. B1	261
9.6	Comparison of lignocellulolytic enzymes activities of differently sourced enzymes used in enzymatic degradation of sawdust	262
9.7	Comparison of soluble protein concentration of differently sourced enzymes used in enzymatic degradation of sawdust	263

LIST OF FIGURES

FIGURE NO.	TITLE	PAGE
2.1	Diagrammatic illustration of the framework of lignocellulose	14
2.2	Structure of cellulose	15
2.3	Structure of hemicellulose	17
2.4	Structure of lignin	19
2.5	Overview of lignocellulose degradation via acidic and enzymatic approach	26
2.6	Schematic presentations of cellulases sites of action on the cellulose polymer liberating glucose	30
2.7	Specific attack sites for xylanolytic enzymes on the structure of xylan	31
2.8	Enzymatic attack on galactoglucomannan structure	33
2.9	Lignin biodegradation process	35
2.10	Castes for termites (Isoptera)	44
3.1	Outline of this study	73
3.2	Experimental design	75
4.1	Screening of cellulases producing bacteria isolated from <i>Bulbitermes</i> sp. termite gut	103
4.2	Screening of cellulases producing fungi isolated from <i>Bulbitermes</i> sp. termite gut	103
4.3	Screening of xylanase producing bacteria isolated from <i>Bulbitermes</i> sp. termite gut	105
4.4	Screening of xylanase producing fungi isolated from <i>Bulbitermes</i> sp. termite gut	105

4.5	Screening of ligninases producing bacteria isolated from <i>Bulbitermes</i> sp. termite gut	107
4.6	Screening of ligninases producing fungi isolated from <i>Bulbitermes</i> sp. termite gut	107
4.7	Gel electrophoresis of PCR product	108
4.8	Partial internal transcribing spacer sequence of fungal isolate A1	109
4.9	Partial 16S rRNA gene sequence of bacterial isolate B1	109
4.10	Partial 16S rRNA gene sequence of bacterial isolate B2	110
4.11	Partial 16S rRNA gene sequence of bacterial isolate Br3	110
4.12	Neighbour-joining phylogenetic tree	113
5.1	Compatibility evaluation	125
5.2	Activities of endoglucanase produced in monocultures and co-cultures in SSF	127
5.3	Activities of exoglucanase produced in monocultures and co-cultures in solid-state fermentation	128
5.4	Activities of β -glucosidase produced in monocultures and co-cultures in solid-state fermentation.	129
5.5	Activities of xylanase produced in monocultures and co-cultures in solid-state fermentation	131
5.6	Activities of LiP produced in monocultures and co-cultures in solid-state fermentation	132
5.7	Activities of MnP produced in monocultures and co-cultures in solid-state fermentation	134
5.8	Activities of laccase produced in monocultures and co-cultures in solid-state fermentation	134
6.1	Images of <i>Aspergillus</i> sp. A1 and <i>Bacillus</i> sp. B1 by SEM	147
6.2	Image of the fermented sawdust	148
6.3	Co-cultivation of <i>Aspergillus</i> sp. A1 and <i>Bacillus</i> sp. B1 on solid basal medium + 2% saw dust	150
6.4	SEM micrographs of sawdust under 5000 \times magnification	152

6.5	Malachite green staining	153
6.6	Time course of EPSs production with single culture and co-culture	154
6.7	Protein profiles of <i>Bacillus</i> sp. B1, <i>Aspergillus</i> sp. A1 and combination of both after growth for 9 days in SSF of saw dust	156
6.8	Time course of total protein with single culture and co-culture	157
6.9	Time course of endoglucanase activities with single culture and co-culture	159
6.10	Time course of exoglucanase activities with single culture and co-culture	161
6.11	Time course of β -glucosidase activities with single culture and co-culture	163
6.12	Time course of xylanase activities with single culture and co-culture	165
6.13	Time course of LiP activities with single culture and co-culture	167
6.14	Time course of MnP activities with single culture and co-culture	169
6.15	Time course of laccase activities with single culture and co-culture	171
6.16	Time course of N-acetyl-D-glucosamine with single culture and co-culture	173
7.1	Effect of sawdust particle sizes on cellulases and xylanase activities under SSF by A1B1 co-culture	182
7.2	Effect of sawdust particle sizes on ligninases under SSF by A1B1 co-culture	184
7.3	Effect of incubation temperature on cellulases and xylanase activities under SSF by A1B1 co-culture	187

7.4	Effect of incubation temperature on ligninases under SSF by A1B1 co-culture	189
7.5	Effect of initial medium pH on cellulases and xylanase activities under SSF by A1B1 co-culture	192
7.6	Effect of initial medium pH on ligninases under SSF by A1B1 co-culture	192
7.7	Effect of initial moisture content on cellulases and xylanase activities under SSF by A1B1 co-culture	195
7.8	Effect of initial moisture content on ligninases under SSF by A1B1 co-culture	196
7.9	Summary on the effect of co-culture and improvement of operating conditions on endoglucanase activity	200
7.10	Summary on the effect of co-culture and improvement of operating conditions on exoglucanase activity	201
7.11	Summary on the effect of co-culture and improvement of operating conditions on β -glucosidase activity	202
7.12	Summary on the effect of co-culture and improvement of operating conditions on xylanase activity	203
7.13	Summary on the effect of co-culture and improvement of operating conditions on LiP activity	204
7.14	Summary on the effect of co-culture and improvement of operating conditions on MnP activity	205
7.15	Summary on the effect of co-culture and improvement of operating conditions on laccase activity	206
8.1	Effect of temperature on the activities of endoglucanase, exoglucanase, β -glucosidase and xylanase produced by <i>Aspergillus</i> sp. A1 and <i>Bacillus</i> sp. B1 co-culture	214
8.2	Effect of temperature on the activities of LiP, MnP and laccase produced by <i>Aspergillus</i> sp. A1 and <i>Bacillus</i> sp. B1 co-culture	215

8.3	Effect of temperature on the stability of endoglucanase produced by <i>Aspergillus</i> sp. A1 and <i>Bacillus</i> sp. B1 co-culture	216
8.4	Effect of temperature on the stability of exoglucanase produced by <i>Aspergillus</i> sp. A1 and <i>Bacillus</i> sp. B1 co-culture	217
8.5	Effect of temperature on the stability of β -glucosidase produced by <i>Aspergillus</i> sp. A1 and <i>Bacillus</i> sp. B1 co-culture	218
8.6	Effect of temperature on the stability of xylanase produced by <i>Aspergillus</i> sp. A1 and <i>Bacillus</i> sp. B1 co-culture	218
8.7	Effect of temperature on the stability of LiP produced by <i>Aspergillus</i> sp. A1 and <i>Bacillus</i> sp. B1 co-culture	219
8.8	Effect of temperature on the stability of MnP produced by <i>Aspergillus</i> sp. A1 and <i>Bacillus</i> sp. B1 co-culture	221
8.9	Effect of temperature on the stability of laccase produced by <i>Aspergillus</i> sp. A1 and <i>Bacillus</i> sp. B1 co-culture	222
8.10	Effect of pH on the activities of endoglucanase, exoglucanase, β -glucosidase and xylanase produced by <i>Aspergillus</i> sp. A1 and <i>Bacillus</i> sp. B1 co-culture	224
8.11	Effect of pH on the activities of LiP, MnP and laccase produced by <i>Aspergillus</i> sp. A1 and <i>Bacillus</i> sp. B1 co-culture	225
8.12	Effect of pH on the stability of endoglucanase produced by <i>Aspergillus</i> sp. A1 and <i>Bacillus</i> sp. B1 co-culture	226
8.13	Effect of pH on the stability of exoglucanase produced by <i>Aspergillus</i> sp. A1 and <i>Bacillus</i> sp. B1 co-culture	227
8.14	Effect of pH on the stability of β -glucosidase produced by <i>Aspergillus</i> sp. A1 and <i>Bacillus</i> sp. B1 co-culture	228
8.15	Effect of pH on the stability of xylanase produced by <i>Aspergillus</i> sp. A1 and <i>Bacillus</i> sp. B1 co-culture	228

8.16	Effect of pH on the stability of LiP produced by <i>Aspergillus</i> sp. A1 and <i>Bacillus</i> sp. B1 co-culture	229
8.17	Effect of pH on the stability of MnP produced by <i>Aspergillus</i> sp. A1 and <i>Bacillus</i> sp. B1 co-culture	231
8.18	Effect of pH on the stability of laccase produced by <i>Aspergillus</i> sp. A1 and <i>Bacillus</i> sp. B1 co-culture	232
8.19	SDS-PAGE of the crude enzyme	235
9.1	The schematic diagram of sawdust-SSF based biorefining process	241
9.2	Chemical composition of raw, non-fermented and fermented sawdust after 9 days of SSF (dry matter basis)	251
9.3	FTIR spectra of sawdust	255
9.4	Images of sawdust by SEM	258
9.5	Sugars standard chromatogram	265
9.6	HPLC analysis of sugars content in enzymatic degradation extract of single and co-culture fermented sawdust using <i>Aspergillus</i> sp. A1 and <i>Bacillus</i> sp. B1 co-culture crude enzyme	266
9.7	HPLC analysis of sugars content in enzymatic degradation extract of single and co-culture fermented sawdust using commercial cellulase of Celluclast	266
9.8	Total phenolic content and lignin concentration in enzymatic degradation extract of single and co-culture fermented sawdust using <i>Aspergillus</i> sp. A1 and <i>Bacillus</i> sp. B1 co-culture crude enzyme	268
9.9	Total phenolic content and lignin concentration in enzymatic degradation extract of single and co-culture fermented sawdust using commercial cellulase of Celluclast	269

LIST OF ABBREVIATIONS

ADF	-	Acid Detergent Fibre
BSA	-	Bovine Serum Albumin
CMC	-	Carboxymethyl cellulose
DNS	-	Dinitrosalicylic acid
EPS	-	Exopolysaccharides
FTIR	-	Fourier Transform Infrared Spectroscopy
g	-	Gram
h	-	Hour
H ₂ SO ₄	-	Sulphuric acid
HCl	-	Hydrochloric acid
H ₂ O ₂	-	Hydrogen Peroxides
HPLC	-	High Performance Liquid Chromatography
kDa	-	Kilo Dalton
L	-	Liter
LiP	-	Lignin peroxidase
min	-	Minute
mL	-	Milliliter
mm	-	Millimeter
MnP	-	Manganese peroxidase
MW	-	Molecular Weight
NaOH	-	Sodium hydroxide
NA	-	Not available
NAG	-	N-Acetyl-D-Glucosamine
NDF	-	Neutral Detergent Fibre

nm	-	Nanometer
°C	-	Degree Celsius
PAGE	-	Polyacrylamide Gel Electrophoresis
PDA	-	Potato Dextrose Agar
pNPG	-	p-nitrophenyl β -D-glucoside
RID	-	Refractive Index Detector
rpm	-	Rotation per minute
SEM	-	Scanning Electron Microscopy
SDS	-	Sodium Dodecyl Sulfate
SmF	-	Submerged Fermentation
SSF	-	Solid-State Fermentation
U/g	-	Unit of enzyme per gram
U//Lh	-	Unit per litre per hour
v/v	-	Volume per volume
w/v	-	Weight per volume
μ L	-	Micro liter
μ m	-	Micro meter

LIST OF APPENDICES

APPENDIX	TITLE	PAGE
A	Reducing Sugar Assay – DNS Method	336
B	Determination of Endoglucanase Activity	338
C	Determination of Exoglucanase Activity	341
D	Determination of β -glucosidase Activity	344
E	Determination of Xylanase Activity	347
F	Determination of LiP Activity	350
G	Determination of MnP Activity	352
H	Determination of Laccase Activity	354
I	Determination of Protein Content	356
J	Glucosamine Assay	358
K	Quantification of Exopolysaccharide (EPS)	361
L	Buffer Composition	363
M	Determination of Neutral Detergent Fibre (NDF)	365
N	Determination of Acid Detergent Fibre (ADF)	366
O	Determination of Acid Detergent Lignin (ADL)	367
P	HPLC analysis	368
Q	Total Phenolic Content (TPC) Assay	369
R	Determination of Lignin Concentration	371
S	Spore count Using Hemocytometer	373
T	Publication	375

CHAPTER 1

INTRODUCTION

1.1 Background of Research

Wood-based industry in Malaysia began in the early 1900s (Ramasamy *et al.*, 2015). Starting with only to meet the domestic demand at the time, wood-based activities in Malaysia such as logging, sawmilling, primary and secondary manufacturing, have played an important role in the economic development of the country, in which they contributed 2% of the Malaysian Gross Domestic Product (GDP) and 2.7% of the country's total merchandise exports (Malaysian Timber Council, 2014). In year 2014, Malaysia has produced 3,218,515 m³ of logs, 1,893,949 m³ of sawn timber and 3,099,371 m³ of plywood, with Japan, USA and India are the top three leading export destinations for these local timber products. The export of Malaysian wood-based products has recorded a positive growth of 5.1% with total exports of RM 20.5 billion (Malaysian Timber Council, 2014). Wooden furniture remained as the biggest export item contributing RM 6.3 billion, followed by plywood (RM 5.2 billion), sawntimber (RM 2.5 billion), logs (RM 2.1 billion) and Builders' Carpentry and Joinery (BCJ) (RM 1 billion). While these wood-based industries generate profits, they also yielded a huge amount of wood wastes, which can potentially give rise to environmentally sensitive disposal issues. The issues are particularly obvious in sawmills where most of the manufacturing technology in used is old and obsolete (Tye *et al.*, 2011).

It was reported that the generation of wood wastes in the sawmilling sector of Peninsular Malaysia was approximately 45 to 50% of the total volume of saw-log input (Ramasamy *et al.*, 2015). The production of the wood waste can be found in the form of off-cuts, slabs, shavings, bark and sawdust (Mekhilef *et al.*, 2011). As one of the most common residues found in wood-manufacturing entities, sawdust is largely produced during the process of planing mills, moulding plants and furniture manufacturing (Rafiqul and Sakinah, 2012). In practise, the residues are left accumulated or discarded in landfill areas, causing environmental pollution through the generation of dust and dirt. Moreover, dumping sawdust to landfills involves additional cost due to its handling and transportation, which is another burden for the industries. Burning had also been applied as one of the economical method to dispose sawdust. However, the high sulphur content of wood may result in the formation of sulphur dioxide during incineration, thereby aggravating air pollution and decreasing air quality in the vicinity (Buraimoh *et al.*, 2015). In view of these issues, research on the utilization of sawdust to turn into value-added products is of high interest as a way to manage wood residues, especially in the country like Malaysia which has a total of 3975 wood-based manufacturing entities operating within the country (Ramasamy *et al.*, 2015).

Sawdust had been used as a raw material in the derivation of biochar (Ghani *et al.*, 2013), commercial mineral-bonded cement composites (Frybort *et al.*, 2008) and as bulking agent in the composting systems (Zhou *et al.*, 2014). The utilization of sawdust also includes as a source of fuel for the cyclone gasification system (Miskam *et al.*, 2009) and for energy generation in the boilers (Ramasamy *et al.*, 2015). Sawdust gained another credit in biomass research area for being classified as lignocellulosic material with significant proportion of cellulose, hemicellulose and lignin constituted in its chemical composition. On a dry basis, sawdust contains cellulose (31.99%), hemicellulose (13.33%) and lignin (44.36%) with the rest consisting of extractives and ash (Belewu, 2006). Several potential value-added products could be derived from biodegradation of these lignocellulose components. Degradation of cellulose and hemicellulose polymers could produce hexose or pentose sugars which served as important raw material for ethanol production, while lignin degradation has huge potential for the synthesis of a number of useful

chemicals such as vanillin, phenol, quinone and acetic acids (Hamid *et al.*, 2014). Biodegradation of cellulose, hemicellulose and lignin from lignocellulosic residue is very much associated to the efficiencies of lignocellulolytic enzymes to degrade the lignocellulose components (Sánchez, 2009). The effectiveness of the enzymatic mixture is highly dependent on their specific functionality to degrade specific type of lignocellulosic material. The use of same material for enzymes production and degradation was suggested to produce enzymes composition that might be then tailored for degradation functionality of that specific material (Pensupa *et al.*, 2013). It is therefore logical and necessary to produce on-site, tailor-made lignocellulolytic enzymes that are optimized for biodegradation of specific lignocellulosics material.

Due to their degrading capabilities, lignocellulolytic enzymes find application in various type of industrial field such as textile, detergent, food, animal feed, pulp and paper (Niladevi, 2007; Singh *et al.*, 2007). The field of industrial enzymes production represents the heart of biotechnology. It was estimated that the global market for commercial enzymes reached \$3.3 billion in 2010, with the annual growth rate of 6% over 5-year forecast period (Thomas *et al.*, 2013). One of the major issues faced by the global enzymes manufacturing companies is the high cost of raw material, which contributes 40–60% of the total production cost (Singhania *et al.*, 2010). Therefore, efforts were made to reduce the cost of production by using cheaper and abundantly available substrates to produce enzymes with high activity (Alam *et al.*, 2009a; Jabasingh and Nachiyar, 2011; Bansal *et al.*, 2012; Yoon *et al.*, 2014). At present, there are limited studies that describe the utilization of sawdust as a substrate for the production of lignocellulolytic enzymes (Liu *et al.*, 2006; Poorna and Prema, 2007; Bansal *et al.*, 2012). The sawdust was either used as the minor substrate or been chemically pretreated prior to fermentations. None have focused on the use of untreated sawdust as a sole substrate for enzymes production. The use of untreated substrate is preferred because additional pre-treatment process with either acidic or alkaline solvents may eventually produce by-products such as furfural, 5-hydroxymethyl-2 furfural, acetic acid, phenols, heavy metals, levulinic acids and formic acids, with inhibitory effects to the microbial growth and respiration (Ang *et al.*, 2013).

Lignocellulosic materials are fermented by lignocellulolytic microorganisms in the process to produce lignocellulolytic enzymes. The fermentation can be conducted via two different fermentation approaches, submerged fermentation or solid state fermentation (Hansen *et al.*, 2015). Submerged fermentation (SmF) has been the most popular and conventional fermentation technology used by enzymes manufacturing companies such as Novozymes and Genencor (Singhania *et al.*, 2010). However, in nature, the growth and lignocellulose utilization of microorganisms secreting lignocellulolytic enzymes are more closely resemble solid-state fermentation (SSF) condition than the presence of excess water provided by SmF (Hansen *et al.*, 2015). SSF is the fermentation method that is carried out without apparent presence of water, but with sufficient moisture to support the growth of microorganisms on the solid matrix (Pandey, 2003). One of the most added advantages offered by SSF is that enzymes titers are higher than those obtained from SmF (Couto and Sanromán, 2005). This advantage has been associated with a larger biomass and lower product breakdown as observed in SSF process (Viniegra-González *et al.*, 2003). In addition, energy expenditure is lower for SSF compared to SmF since less water is needed, no mechanical mixing and less energy requirement in downstream processing (Hansen *et al.*, 2015). Furthermore, higher concentration of products can be obtained from SSF, making purification works such as concentration and freeze drying are undesirable (Zhuang *et al.*, 2007).

The lignocellulolytic enzymes production also depends on the type of microbial strain and the strains giving higher activities on lignocellulosic material in SSF condition are important. *Aspergillus*, *Trichoderma*, *Rhizopus*, *Fusarium* and *Penicillium* are some of the fungi genera reported able to produce lignocellulolytic enzymes in SSF (Hansen *et al.*, 2015). For bacteria, *Bacillus* and *Streptomyces* are the most common been reported (Krishna, 1999; Niladevi *et al.*, 2007). The fungal and bacterial strains were isolated from substrata containing lignocellulosic carbon source such as residues from different agricultural sectors, soil and debris from cereal production (Jabasingh and Nachiyar, 2011; Irfan *et al.*, 2012; Ang *et al.*, 2013). Another interesting source to prospect for lignocellulolytic microorganisms is from the guts of insects. Some insects relied upon their gut microbial community to degrade lignocellulosic material as their nutrient sources. One of these insects is

termite. Termites were reported to harbouring diverse array of lignocellulolytic microorganisms inside their gut (Dheeran *et al.*, 2012). Several lignocellulolytic microorganisms such as *Pseudomonas*, *Bacillus*, *Enterobacter*, *Streptomyces*, *Paenibacillus*, *Aspergillus* and *Sporothrix* had been successfully isolated from termite species of *Coptotermes curvignathus* (Ramin *et al.*, 2009), *Reticulitermes santonensis* (Matkar *et al.*, 2013) and *Amitermes hastatus* (Le Roes-Hill *et al.*, 2011). However, the capability of the microorganisms originated from termite's gut to produce lignocellulolytic enzymes have only been studied in culture employing SmF technique. The potential of microorganisms isolated from termite gut in producing lignocellulolytic enzymes under SSF remained to be addressed.

Earlier reports are available for the production of lignocellulolytic enzymes by single culture of bacteria and fungi from termite gut (Wenzel *et al.*, 2002; Ramin *et al.*, 2009; Le Roes-Hill *et al.*, 2011; Dheeran *et al.*, 2012). However, a single microorganism cannot produce all the enzymes necessary for complete bioconversion of lignocellulose and different microorganisms are normally co-exist symbiotically on solid substrates in nature (Yoon *et al.*, 2014). Thus, co-culturing of microorganisms which act synergistically for rapid bioconversion of lignocellulosic biomass under SSF, is attractive (Wang *et al.*, 2006; Kumar *et al.*, 2008a). Co-culture defined as inoculation of different specified microbial strains under aseptic conditions, had been used to achieve improved production of biologically active compounds such as organic acids, vitamins and antibiotics (Bader *et al.*, 2010). Similarly, co-culture is beneficial for production of lignocellulolytic enzymes during biodegradation of lignocellulosic substrate (Brijwani *et al.*, 2010; Dhillon *et al.*, 2011; Kolasa *et al.*, 2014) as they offer higher productivity of enzymes and better adaptability compared to single culture (Dashtban *et al.*, 2010). Hence, it is hypothesized that through co-culture techniques, improved level of lignocellulolytic enzymes produced by synergistic interactions between different microorganisms may be achieved in single process. It may further eliminates the requirement to cultivate multiple single cultures separately, followed by enzymes blending which then increases the cost of double equipment needed (Kolasa *et al.*, 2014). As termite gut was known to contain dense population of microbiota that work co-operatively in

lignocellulosic material decomposition, co-culturing microorganisms originated from such sources remained as an interesting topic to be further explored.

1.2 Objectives

The objectives of this research are as follows:

1. To isolate, screen and identify the termite guts microorganisms with the capability to produce lignocellulolytic enzymes in SSF using untreated sawdust as substrate.
2. To construct and to evaluate the compatibility of the members in microbial co-culture with promising level of lignocellulolytic enzymes activities. The profile of lignocellulolytic enzymes produced by both single and microbial co-culture and its relation with exopolysaccharides production, N-acetyl-D-glucosamine and protein concentration will be analysed.
3. To study the effect of sawdust particle size, incubation temperature, pH and moisture content on the production of lignocellulolytic enzymes by varying one variable at a time.
4. To characterize the lignocellulolytic enzymes produced by a selected microbial co-culture in terms of its optimum pH, optimum temperature, pH stability and thermal stability. A sawdust-based biorefining strategy for reducing sugars production will be developed.

1.3 Scope of the Research

This study focused in investigating the capability of termite gut's microorganisms to produce lignocellulolytic enzymes under SSF by using untreated sawdust as solid substrate. The bacterial and fungal isolates from *Bulbitermes* sp. termite gut were screened through qualitative approach by using plate base technique and the lignocellulolytic activities were assessed quantitatively in SSF condition. Lignocellulolytic activities were calculated based on the endoglucanase, exoglucanase, β -glucosidase, xylanase, lignin peroxidase, manganese peroxidase and laccase activities. The isolates with highest lignocellulolytic activities were selected, identified and further used for the development of microbial co-culture.

The effect of sawdust particle size, incubation temperature, pH of the medium and moisture content on lignocellulolytic enzymes production were studied. An optimal condition for the enzymes production was set. Lignocellulolytic enzymes obtained from the optimal SSF condition were characterized by means of determination of their optimal temperature, pH, and stability.

The biodegradation potential by single and microbial co-culture cultivated under SSF were studied. A sawdust-based biorefining strategy was developed by extracting the lignocellulolytic enzymes produced from SSF process and then used to hydrolyse the fermented sawdust. The amounts of reducing sugars obtained after the hydrolysis process were measured.

1.4 Significance of the Research

As Malaysia has significant amount of woody-based activities such as logging and saw-milling, the mass generation of sawdust as the most common wood residues produced by the forestry related industries, can potentially give rise to environmentally sensitive disposal issues (Hoi, 2003). This had urged the need to properly utilize the sawdust to turn into various value-added products including as raw material for the production of lignocellulolytic enzymes. Below are several identified issues which make the current research is significance:

- i. Improper management of wood residues including sawdust can give an adverse effect towards the air quality, which remained an issue to be solved by the parties involved including the local community, the wood-based industries themselves and the government enforcement bodies. Sawdust is therefore proposed as an alternative raw material that serves as substrate for the production of lignocellulolytic enzymes.
- ii. The afore mentioned suggestion is in line with the Malaysian government effort in exploiting the country's biomass resources up to its optimum level as outlined in National Biomass Strategy 2020 (Agensi Inovasi Malaysia, 2011). From Malaysian perspectives as an important global exporter for wood-based products, the use of sawdust for lignocellulolytic enzymes production is a promising technology to add more value to the wood residues as well as providing more opportunities to achieve economic advancement for the industrial player.
- iii. The cost of raw material contributes for about 40–60 % of the total enzyme production cost. A cheaper alternative substrate can be prospected as a way to reduce the production cost by using the raw untreated sawdust as sole substrate in the process of lignocellulolytic enzymes production. The lack of chemical or/and physical pre-treatment step during the substrate preparation stage could further reduce the cost of overall enzyme production. Furthermore, enzymes production in SSF can also facilitates a lower capital operating cost due to less water requirements and lower energy expenditures. This study is regarded as the first to describe the use of untreated sawdust as a sole solid support in SSF for lignocellulolytic enzymes production.

- iv. Investigation to find new isolates from habitats containing lignocellulosic substrates with the capability to produce lignocellulolytic enzymes are relatively simple strategies to obtain higher titre of enzymatic activities in facilitating the biomass degradation process. Since termite gut stands as a rich source to prospect for diverse and efficient lignocellulose degrading microorganisms, the current study described the potential of termite gut microorganisms in producing lignocellulolytic enzymes and also degrading untreated sawdust under SSF condition.
- v. Although several studies have described the capability of termite gut microorganisms to exhibit lignocellulolytic activities, none has reported the effect when the microorganisms are co-cultured together. As termite gut holds a dense population of microorganisms, co-culturing may provide an insight into types of interactions existed between the guts microbiota.
- vi. Knowledge on self-production of lignocellulolytic enzymes is essential as tailor-made enzymatic mixtures that are optimized for the degradation of specific type of lignocellulosics remains as a strategic issue to be considered during the development of a sustainable biomass-biorefinery process. The use of same material for enzyme production and degradation process could be a way to obtain optimal degradation results of that specific material. Therefore, cultivation of microorganisms on sawdust was projected to produce lignocellulolytic enzymes with specific functionality to degrade sawdust and simultaneously promoting a greener technology as a way to manage woody residues in Malaysia.

1.5 Thesis Organization

This thesis is organized into ten chapters. **Chapter 2** covers relevant literatures on the availability of lignocellulosic biomass in Malaysia, structure of lignocellulose and the potential of sawdust to be used as raw material for production of high value products. This chapter provides an overview of lignocellulose degradation via acidic and enzymatic approach, and the role played by lignocellulolytic enzymes (cellulases, xylanase and ligninases) in the enzymatic degradation process. The source to prospect for lignocellulolytic microorganisms

and the plausibility of termite gut to serve as a good reservoir for isolation of such microorganisms were explained. This chapter also deals with information on SmF and SSF as well as important SSF process variables related to the production of lignocellulolytic enzymes. The positive role of microbial co-culture in lignocellulolytic enzymes production was also reviewed. Literatures related to application of lignocellulolytic enzymes in various industries are briefly summarized.

Chapter 3 describes the general experimental procedures performed in this research. All common methods and procedures are placed in this chapter and be referred to in specific chapters, respectively.

The results and discussions are divided into six main chapters. **Chapter 4** describes the isolation, screening and identification of lignocellulolytic microorganisms from *Bulbitermes* sp. termite gut. **Chapter 5** presents the development of microbial co-culture from the selected lignocellulolytic microorganisms in order to improve the enzymatic activities. In **Chapter 6**, a thorough comparison was made between the lignocellulolytic enzymes activities produced by microbial co-culture with its respective single culture member. The profile of lignocellulolytic activities together with its relation with exopolysaccharides production, N-acetyl-D-glucosamine and protein concentration was described. **Chapter 7** provided the evaluation of the effect of SSF operating parameters on lignocellulolytic enzymes activities. **Chapter 8** presents the characteristics of lignocellulolytic enzymes produced by microbial co-culture in terms of optimal temperature and pH, temperature and pH stability. Activity staining and molecular mass of lignocellulolytic enzymes on SDS-PAGE gel were determined. The capability of single and microbial co-culture of termite gut's microorganisms to degrade lignocellulose and the development of sawdust-based biorefinery strategy were presented in **Chapter 9**.

The conclusions from this research are given in **Chapter 10**. This chapter also states specific achievements, problems and some recommendations for future work.

10.2 Recommendations

The utilization of sawdust as a raw material for the production of highly valuable lignocellulolytic enzymes and fermentable sugars will not only fetch valuable remuneration for wood-based industries, but also help mitigate environmental pollution. In addition, through the study of microorganisms and their enzymatic activities, the mechanisms of efficient lignocellulose degradation in the termite gut may then could be elucidated, findings which have significant potential in biorefinery industries. The defined microbial co-culture also stands as a useful technique to improve the titre of lignocellulolytic enzymes activities and therefore worthy of future study. Some recommendations for future studies are outlined as follows:

- i. As sawdust was observed to contain high content of lignin, it is very interesting to extract the lignin through enzymatic or biological approach as this natural polymer can serve as a base for different materials application in the fields of bioplastics, (nano) composites and nanoparticles.
- ii. Since termite is one of the insects with a dense population of microorganisms living symbiotically inside its guts it is highly expected that the termite gut also resides microorganism that could influence the performance of the fermentation system, including hydrogen yield. Biohydrogen is regarded as one of potentially advantageous alternative energy to minimise or even eliminate the dependability on fossil fuels. Future research should consider prospecting and characterising hydrogen-producing microorganisms from the guts of termites.
- iii. To obtain the best production of enzymes, identification of optimal ratio between the two microorganisms in a co-cultivation is necessary. The addition of termite extract into the medium or substrate can also be considered as a strategy to enhance the growth of microorganisms isolated from the termite gut. It is also feasible to construct an efficient

lignocellulolytic enzymes producing–co-culture for reducing sugars preparation from lignocellulosic biomass by adjusting the microbial constituent proportions in the consortium.

- iv. The present study was able to show that reducing sugars can be produced from the enzymatic degradation of sawdust. Future studies should focus to investigate whether the reducing sugars can be further fermented by microorganisms to make ethanol from sawdust.

- v. Static tray fermentation is often used for large-scale production of enzymes, as it offers potential benefits over bioreactors, such as simple technique, trays can be stacked over one another in shelves and higher yields. Solid-state tray fermentation could be possibly used to achieve higher yield of lignocellulolytic enzymes due to the capacity to put high substrate loading, large area for microorganism to grow and easy handling bioreactor as compared to immersion, packed-bed and rotating drum bioreactor. A more comprehensive study is needed to provide information about the production of lignocellulolytic enzymes in solid-state tray fermentation employing co-culture of selected microorganisms.

REFERENCES

- Adams, L. and Boopathy, R. (2005). Isolation and characterization of enteric bacteria from the hindgut of Formosan termite. *Bioresource Technology*. 96, 1592-1598.
- Aden, A. and Foust, T. (2009). Technoeconomic analysis of the dilute sulfuric acid and enzymatic hydrolysis process for the conversion of corn stover to ethanol. *Cellulose*. 16, 535-545.
- Aditiya, H., Chong, W., Mahlia, T., Sebayang, A., Berawi, M. and Nur, H. (2016). Second generation bioethanol potential from selected Malaysia's biodiversity biomasses: A review. *Waste Management*. 47, 46-61.
- Ahamed, A. and Vermette, P. (2008). Enhanced enzyme production from mixed cultures of *Trichoderma reesei* RUT-C30 and *Aspergillus niger* LMA grown as fed batch in a stirred tank bioreactor. *Biochemical Engineering Journal*. 42, 41-46.
- Ahn, Y., Song, Y., Kwak, S. Y. and Kim, H. (2016). Highly ordered cellulose II crystalline regenerated from cellulose hydrolyzed by 1-butyl-3-methylimidazolium chloride. *Carbohydrate Polymers*. 137, 321-327.
- Aidoo, K. E., Hendry, R. and Wood, B. (1981). Estimation of fungal growth in a solid state fermentation system. *European Journal of Applied Microbiology and Biotechnology*. 12, 6-9.
- Agensi Inovasi Malaysia. (2011). National Biomass Strategy 2020: New wealth creation for Malaysia's palm oil industry.
- Akocak, P. B., Churey, J. J. and Worobo, R. W. (2015). Antagonistic effect of chitinolytic *Pseudomonas* and *Bacillus* on growth of fungal hyphae and spores of aflatoxigenic *Aspergillus flavus*. *Food Bioscience*. 10, 48-58.
- Akpınar, O., Erdogan, K. and Bostancı, S. (2009). Enzymatic production of xylooligosaccharide from selected agricultural wastes. *Food and Bioproducts Processing*. 87, 145-151.

- Aktaş, N. and Tanyolaç, A. (2003). Reaction conditions for laccase catalyzed polymerization of catechol. *Bioresource Technology*. 87, 209-214.
- Alam, M. Z., Fakhru'l-Razi, A., Abd-Aziz, S. and Molla, A. H. (2003). Optimization of compatible mixed cultures for liquid state bioconversion of municipal wastewater sludge. *Water, Air and Soil Pollution*. 149, 113-126.
- Alam, M. Z., Mamun, A. A., Qudsieh, I. Y., Muyibi, S. A., Salleh, H. M. and Omar, N. M. (2009a). Solid state bioconversion of oil palm empty fruit bunches for cellulase enzyme production using a rotary drum bioreactor. *Biochemical Engineering Journal*. 46, 61-64.
- Alam, M. Z., Mansor, M. F. and Jalal, K. (2009b). Optimization of lignin peroxidase production and stability by *Phanerochaete chrysosporium* using sewage-treatment-plant sludge as substrate in a stirred-tank bioreactor. *Journal of Industrial Microbiology and Biotechnology*. 36, 757-764.
- Alam, M. Z., Muhammad, N. and Mahmat, M. E. (2005). Production of cellulase from oil palm biomass as substrate by solid state bioconversion. *American Journal of Applied Science*. 2, 569-572.
- Amel, B. D., Nawel, B., Khelifa, B., Mohammed, G., Manon, J., Salima, K. G., Farida, N., Hocine, H., Bernard, O. and Jean-Luc, C. (2016). Characterization of a purified thermostable xylanase from *Caldicoprobacter algeriensis* sp. nov. strain TH7C1 T. *Carbohydrate Research*. 419, 60-68.
- An, J., Xie, Y., Zhang, Y., Tian, D., Wang, S., Yang, G. and Feng, Y. (2015). Characterization of a thermostable, specific GH10 xylanase from *Caldicellulosiruptor bescii* with high catalytic activity. *Journal of Molecular Catalysis B: Enzymatic*. 117, 13-20.
- Anand, A. A. P., Vennison, S. J., Sankar, S. G., Prabhu, D. I. G., Vasani, P. T., Raghuraman, T., Geoffrey, C. J. and Vendan, S. E. (2010). Isolation and characterization of bacteria from the gut of *Bombyx mori* that degrade cellulose, xylan, pectin and starch and their impact on digestion. *Journal of Insect Science*. 10, 107.
- Ang, S. K., Shaza, E., Adibah, Y., Suraini, A. A. and Madihah, M. (2013). Production of cellulases and xylanase by *Aspergillus fumigatus* SK1 using untreated oil palm trunk through solid state fermentation. *Process Biochemistry*. 48, 1293-1302.

- Ang, S. K., Yahya, A., Abd Aziz, S. and Md Salleh, M. (2015a). Isolation, screening and identification of potential cellulolytic and xylanolytic producers for biodegradation of untreated oil palm trunk and its application in saccharification of lemongrass leaves. *Preparative Biochemistry and Biotechnology*. 45, 279-305.
- Ang Siow Kuang (2015). *Cellulases and Xylanase Production by Aspergillus fumigatus SK1 through Solid State Fermentation for Ethanol Fermentation*. PhD Thesis Page 91, Universiti Teknologi Malaysia, Skudai.
- Ang, S. K., Yahya, A., Abd Aziz, S. and Md Salleh, M. (2015b). Potential uses of xylanase-rich lignocellulolytic enzymes cocktail for oil palm trunk (OPT) degradation and lignocellulosic ethanol production. *Energy and Fuels*. 29, 5103-5116.
- Angelis, S., Novak, A., Sydney, E., Soccol, V., Carvalho, J., Pandey, A., Nosedá, M., Tholozan, J. L., Lorquin, J. and Soccol, C. (2012). Co-culture of microalgae, cyanobacteria, and macromycetes for exopolysaccharides production: process preliminary optimization and partial characterization. *Applied Biochemistry and Biotechnology*. 167, 1092-1106.
- Antoine, A. A., Jacqueline, D. and Thonart, P. (2010). Xylanase production by *Penicillium canescens* on soya oil cake in solid-state fermentation. *Applied Biochemistry and Biotechnology*. 160, 50-62.
- Arias, M. E., Arenas, M., Rodríguez, J., Soliveri, J., Ball, A. S. and Hernández, M. (2003). Kraft pulp biobleaching and mediated oxidation of a nonphenolic substrate by laccase from *Streptomyces cyaneus* CECT 3335. *Applied and Environmental Microbiology*. 69, 1953-1958.
- Ariffin, H., Abdullah, N., Umi Kalsom, M., Shirai, Y. and Hassan, M. (2006). Production and characterization of cellulase by *Bacillus pumilus* EB3. *International Journal of Engineering and Technology*. 3, 47-53.
- Ariffin, H., Hassan, M. A., Shah, U. K. M., Abdullah, N., Ghazali, F. M. and Shirai, Y. (2008). Production of bacterial endoglucanase from pretreated oil palm empty fruit bunch by *Bacillus pumilus* EB3. *Journal of Bioscience and Bioengineering*. 106, 231-236.
- Arora, A., Nain, L. and Gupta, J. (2005). Solid-state fermentation of wood residues by *Streptomyces griseus* B1, a soil isolate, and solubilization of lignins. *World Journal of Microbiology and Biotechnology*. 21, 303-308.

- Arora, D. S., Chander, M. and Gill, P. K. (2002). Involvement of lignin peroxidase, manganese peroxidase and laccase in degradation and selective ligninolysis of wheat straw. *International Biodeterioration and Biodegradation*. 50, 115-120.
- Arora, D. S. and Gill, P. K. (2001). Comparison of two assay procedures for lignin peroxidase. *Enzyme and Microbial Technology*. 28, 602-605.
- Arora, D. S. and Sharma, R. K. (2010). Ligninolytic fungal laccases and their biotechnological applications. *Applied Biochemistry and Biotechnology*. 160, 1760-1788.
- Arora, D. S. and Sharma, R. K. (2011). Effect of different supplements on bioprocessing of wheat straw by *Phlebia brevispora*: changes in its chemical composition, in vitro digestibility and nutritional properties. *Bioresource Technology*. 102, 8085-8091.
- Asgher, M., Ahmed, N. and Iqbal, H. M. N. (2011). Hyperproductivity of extracellular enzymes from indigenous white rot fungi (*Phanerochaete chrysosporium*) by utilizing agro-wastes. *BioResources*. 6, 4454-4467.
- Asgher, M., Asad, M. and Legge, R. (2006). Enhanced lignin peroxidase synthesis by *Phanerochaete chrysosporium* in solid state bioprocessing of a lignocellulosic substrate. *World Journal of Microbiology and Biotechnology*. 22, 449-453.
- Asgher, M., Iqbal, H. M. N. and Irshad, M. (2012). Characterization of purified and xerogel immobilized novel lignin peroxidase produced from *Trametes versicolor* IBL-04 using solid state medium of corncobs. *BMC Biotechnology*. 12, 46.
- Asgher, M., Wahab, A., Bilal, M. and Iqbal, H. M. N. (2016). Lignocellulose degradation and production of lignin modifying enzymes by *Schizophyllum commune* IBL-06 in solid-state fermentation. *Biocatalysis and Agricultural Biotechnology*. 6, 195-201.
- Asha Poorna, C. and Prema, P. (2007). Production of cellulase-free endoxylanase from novel alkalophilic thermotolerant *Bacillus pumilus* by solid-state fermentation and its application in wastepaper recycling. *Bioresource Technology*. 98, 485-490.

- Assareh, R., Zahiri, H. S., Noghabi, K. A. and Aminzadeh, S. (2012). Characterization of the newly isolated *Geobacillus* sp. T1, the efficient cellulase-producer on untreated barley and wheat straws. *Bioresource Technology*. 120, 99-105.
- Auxenfans, T., Buchoux, S., Larcher, D., Husson, G., Husson, E. and Sarazin, C. (2014). Enzymatic saccharification and structural properties of industrial wood sawdust: recycled ionic liquids pretreatments. *Energy Conversion and Management*. 88, 1094-1103.
- Azadi, P., Inderwildi, O. R., Farnood, R. and King, D. A. (2013). Liquid fuels, hydrogen and chemicals from lignin: A critical review. *Renewable and Sustainable Energy Reviews*. 21, 506-523.
- Bader, J., Mast-Gerlach, E., Popović, M., Bajpai, R. and Stahl, U. (2010). Relevance of microbial co-culture fermentations in biotechnology. *Journal of Applied Microbiology*. 109, 371-387.
- Bahrin, E. K., Seng, P. Y. and Abd Aziz, S. (2011). Effect of oil palm empty fruit bunch particle size on cellulase production by *Botryosphaeria* sp. under solid state fermentation. *Australian Journal of Basic and Applied Sciences*. 5, 276-280.
- Bajaj, B. K., Sharma, M. and Sharma, S. (2011). Alkalistable endo- β -1, 4-xylanase production from a newly isolated alkalitolerant *Penicillium* sp. SS1 using agro-residues. *3 Biotech*. 1, 83-90.
- Bajpai, P. (1999). Application of enzymes in the pulp and paper industry. *Biotechnology Progress*. 15, 147-157.
- Bakar, N. K. A., Zanirun, Z., Abd Aziz, S., Ghazali, F. M. and Hassan, M. A. (2012). Production of fermentable sugars from oil palm empty fruit bunch using crude cellulase cocktails with *Trichoderma asperellum* UPM1 and *Aspergillus fumigatus* UPM2 for bioethanol production. *BioResources*. 7, 3627-3639.
- Baker, R. A. and Wicker, L. (1996). Current and potential applications of enzyme infusion in the food industry. *Trends in Food Science and Technology*. 7, 279-284.
- Baldrian, P. (2004). Increase of laccase activity during interspecific interactions of white-rot fungi. *FEMS Microbiology Ecology*. 50, 245-253.

- Bansal, N., Tewari, R., Soni, R. and Soni, S. K. (2012). Production of cellulases from *Aspergillus niger* NS-2 in solid state fermentation on agricultural and kitchen waste residues. *Waste Management*. 32, 1341-1346.
- Bashir, Z., Kondapalli, V. K., Adlakha, N., Sharma, A., Bhatnagar, R. K., Chandel, G. and Yazdani, S. S. (2013). Diversity and functional significance of cellulolytic microbes living in termite, pill-bug and stem-borer guts. *Scientific Reports*. 3.
- Batool, S., Asgher, M., Sheikh, M. and Rahma, S. (2013). Optimization of physical and nutritional factors for enhanced production of lignin peroxidase by *Ganoderma lucidum* IBL-05 in solid state culture of wheat straw. *Journal of Animal and Plant Sciences*. 23, 1166-1176.
- Bauer, C. G., Kühn, A., Gajovic, N., Skorobogatko, O., Holt, P. J., Bruce, N. C., Makower, A., Lowe, C. R. and Scheller, F. W. (1999). New enzyme sensors for morphine and codeine based on morphine dehydrogenase and laccase. *Fresenius' Journal of Analytical Chemistry*. 364, 179-183.
- Beg, Q., Kapoor, M., Mahajan, L. and Hoondal, G. (2001). Microbial xylanases and their industrial applications: A review. *Applied Microbiology and Biotechnology*. 56, 326-338.
- Belewu, M. (2006). Conversion of masonia tree sawdust and cotton plant by product into feed by white rot fungus. *African Journal of Biotechnology*. 5, 1763-1764.
- Belinky, P., Lasser, H. and Dosoretz, C. (2014). U.S. Patent No. 8, 691, 194. Washington DC: U.S. Patent and Trademark Office.
- Bending, G. D. and Read, D. J. (1997). Lignin and soluble phenolic degradation by ectomycorrhizal and ericoid mycorrhizal fungi. *Mycological Research*. 101, 1348-1354.
- Bermek, H., Yazıcı, H., Öztürk, H., Tamerler, C., Jung, H., Li, K., Brown, K. M., Ding, H. and Xu, F. (2004). Purification and characterization of manganese peroxidase from wood-degrading fungus *Trichophyton rubrum* LSK-27. *Enzyme and Microbial Technology*. 35, 87-92.
- Bhalla, A., Bischoff, K. M. and Sani, R. K. (2015). Highly thermostable xylanase production from a thermophilic *Geobacillus* sp. strain WsUcF1 utilizing lignocellulosic biomass. *Frontiers in Bioengineering and Biotechnology*. 3, 1-8.

- Bhat, M. (2000). Cellulases and related enzymes in biotechnology. *Biotechnology Advances*. 18, 355-383.
- Bholay, A., Borkhataria, B. V., Jadhav, P. U., Palekar, K. S., Dhalkari, M. V. and Nalawade, P. (2012). Bacterial lignin peroxidase: A tool for biobleaching and biodegradation of industrial effluents. *Universal Journal of Environmental Research and Technology*. 2, 58-64.
- Bianchetti, C. M., Brumm, P., Smith, R. W., Dyer, K., Hura, G. L., Rutkoski, T. J. and Phillips, G. N. (2013). Structure, dynamics, and specificity of endoglucanase D from *Clostridium cellulovorans*. *Journal of Molecular Biology*. 425, 4267-4285.
- Bibi, Z., Ansari, A., Zohra, R. R., Aman, A. and Qader, S. A. U. (2014). Production of xylan degrading endo-1, 4- β -xylanase from thermophilic *Geobacillus stearothermophilus* KIBGE-IB29. *Journal of Radiation Research and Applied Sciences*. 7, 478-485.
- Biely, P., Markovič, O. and Mislovičová, D. (1985). Sensitive detection of endo-1, 4- β -glucanases and endo-1, 4- β -xylanases in gels. *Analytical Biochemistry*. 144, 147-151.
- Bilal, M. and Asgher, M. (2015). Sandal reactive dyes decolorization and cytotoxicity reduction using manganese peroxidase immobilized onto polyvinyl alcohol-alginate beads. *Chemistry Central Journal*. 9, 1-14.
- Biolog (2002). *Microlog User's Manual, Release 4.2*. [Brochure]. California: Hayward.
- Blandino, A., Iqbalsyah, T., Pandiella, S., Cantero, D. and Webb, C. (2002). Polygalacturonase production by *Aspergillus awamori* on wheat in solid-state fermentation. *Applied Microbiology and Biotechnology*. 58, 164-169.
- Boer, C. G., Obici, L., De Souza, C. G. M. and Peralta, R. M. (2006). Purification and some properties of Mn peroxidase from *Lentinula edodes*. *Process Biochemistry*. 41, 1203-1207.
- Boonyeun, P., Shotipruk, A., Prommuak, C., Supphantharika, M. and Muangnapoh, C. (2011). Enhancement of amino acid production by two-step autolysis of spent brewer's yeast. *Chemical Engineering Communications*. 198, 1594-1602.

- Boruah, P., Dowarah, P., Hazarika, R., Yadav, A., Barkakati, P. and Goswami, T. (2015). Xylanase from *Penicillium meleagrinum* var. *viridiflavum*-a potential source for bamboo pulp bleaching. *Journal of Cleaner Production*. 116, 259-267.
- Borzani, W., Salomão, G., Martins, J. and Alonso, V. (1999). A simple method to control the moisture content of the fermenting medium during laboratory-scale solid-state fermentation experiments. *Brazilian Journal of Chemical Engineering*. 16, 101-102.
- Botella, C., Diaz, A., De Ory, I., Webb, C. and Blandino, A. (2007). Xylanase and pectinase production by *Aspergillus awamori* on grape pomace in solid state fermentation. *Process Biochemistry*. 42, 98-101.
- Bourbonnais, R., Paice, M., Reid, I., Lanthier, P. and Yaguchi, M. (1995). Lignin oxidation by laccase isozymes from *Trametes versicolor* and role of the mediator 2, 2'-azinobis (3-ethylbenzthiazoline-6-sulfonate) in kraft lignin depolymerization. *Applied and Environmental Microbiology*. 61, 1876-1880.
- Brijwani, K., Oberoi, H. S. and Vadlani, P. V. (2010). Production of a cellulolytic enzyme system in mixed-culture solid-state fermentation of soybean hulls supplemented with wheat bran. *Process Biochemistry*. 45, 120-128.
- Brinchi, L., Cotana, F., Fortunati, E. and Kenny, J. (2013). Production of nanocrystalline cellulose from lignocellulosic biomass: technology and applications. *Carbohydrate Polymers*. 94, 154-169.
- Brown, L., Wolf, J. M., Prados-Rosales, R. and Casadevall, A. (2015). Through the wall: Extracellular vesicles in gram-positive bacteria, mycobacteria and fungi. *Nature Reviews Microbiology*. 13, 620-630.
- Brown, M. E. and Chang, M. C. (2014). Exploring bacterial lignin degradation. *Current Opinion in Chemical Biology*. 19, 1-7.
- Brune, A. (2013). Symbiotic Associations between Termites and Prokaryotes. In Dworkin, M., Falkow, S. and Rosenberg, E. (Eds) *The Prokaryotes* (pp. 545-577). New York: Springer.
- Brune, A. (2014). Symbiotic digestion of lignocellulose in termite guts. *Nature Reviews Microbiology*. 12, 168-180.
- Bugg, T. D., Ahmad, M., Hardiman, E. M. and Singh, R. (2011). The emerging role for bacteria in lignin degradation and bio-product formation. *Current Opinion in Biotechnology*. 22, 394-400.

- Buraimoh, O. M., Ilori, M. O., Amund, O. O., Michel, F. C. and Grewal, S. K. (2015). Assessment of bacterial degradation of lignocellulosic residues (sawdust) in a tropical estuarine microcosm using improvised floating raft equipment. *International Biodeterioration and Biodegradation*. 104, 186-193.
- Cai, Y., Wu, H., Liao, X., Ding, Y., Sun, J. and Zhang, D. (2010). Purification and characterization of novel manganese peroxidase from *Rhizoctonia* sp. SYBC-M3. *Biotechnology and Bioprocess Engineering*. 15, 1016-1021.
- Camarero, S., Garcia, O., Vidal, T., Colom, J., Del Rio, J. C., Gutiérrez, A., Gras, J. M., Monje, R., Martinez, M. J. and Martinez, Á. T. (2004). Efficient bleaching of non-wood high-quality paper pulp using laccase-mediator system. *Enzyme and Microbial Technology*. 35, 113-120.
- Carabajal, M., Levin, L., Albertó, E. and Lechner, B. (2012). Effect of co-cultivation of two *Pleurotus* species on lignocellolytic enzyme production and mushroom fructification. *International Biodeterioration and Biodegradation*. 66, 71-76.
- Carere, C. R., Sparling, R., Cicek, N. and Levin, D. B. (2008). Third generation biofuels via direct cellulose fermentation. *International Journal of Molecular Sciences*. 9, 1342-1360.
- Castoldi, R., Bracht, A., De Moraes, G. R., Baesso, M. L., Correa, R. C. G., Peralta, R. A., Moreira, R. D. F. P. M., De Moraes, M. D. L. T., De Souza, C. G. M. and Peralta, R. M. (2014). Biological pretreatment of *Eucalyptus grandis* sawdust with white-rot fungi: Study of degradation patterns and saccharification kinetics. *Chemical Engineering Journal*. 258, 240-246.
- Chan Cupul, W., Heredia Abarca, G., Martínez Carrera, D. and Rodríguez Vázquez, R. (2014). Enhancement of ligninolytic enzyme activities in a *Trametes maxima*-*Paecilomyces carneus* co-culture: Key factors revealed after screening using a Plackett-Burman experimental design. *Electronic Journal of Biotechnology*. 17, 114-121.
- Chandrasekaran, G., Choi, S. K., Lee, Y. C., Kim, G. J. and Shin, H. J. (2014). Oxidative biodegradation of single-walled carbon nanotubes by partially purified lignin peroxidase from *Sparassis latifolia* mushroom. *Journal of Industrial and Engineering Chemistry*. 20, 3367-3374.

- Chang, A. J., Fan, J. and Wen, X. (2012). Screening of fungi capable of highly selective degradation of lignin in rice straw. *International Biodeterioration and Biodegradation*. 72, 26-30.
- Chang, Y. C., Choi, D., Takamizawa, K. and Kikuchi, S. (2014). Isolation of *Bacillus* sp. strains capable of decomposing alkali lignin and their application in combination with lactic acid bacteria for enhancing cellulase performance. *Bioresource Technology*. 152, 429-436.
- Chapla, D., Divecha, J., Madamwar, D. and Shah, A. (2010). Utilization of agro-industrial waste for xylanase production by *Aspergillus foetidus* MTCC 4898 under solid state fermentation and its application in saccharification. *Biochemical Engineering Journal*. 49, 361-369.
- Chekan, J. R., Kwon, I. H., Agarwal, V., Dodd, D., Revindran, V., Mackie, R. I., Cann, I. and Nair, S. K. (2014). Structural and biochemical basis for mannan utilization by *Caldanaerobius polysaccharolyticus* strain ATCC BAA-17. *Journal of Biological Chemistry*. 289, 34965-34977.
- Chen, H. Z. and He, Q. (2012). Value-added bioconversion of biomass by solid-state fermentation. *Journal of Chemical Technology and Biotechnology*. 87, 1619-1625.
- Chen, M., Zhao, J. and Xia, L. (2008). Enzymatic hydrolysis of maize straw polysaccharides for the production of reducing sugars. *Carbohydrate Polymers*. 71, 411-415.
- Cheng, C. L. and Chang, J. S. (2011). Hydrolysis of lignocellulosic feedstock by novel cellulases originating from *Pseudomonas* sp. CL3 for fermentative hydrogen production. *Bioresource Technology*. 102, 8628-8634.
- Cherry, J. R., and Fidantsef, A. L. (2003). Directed evolution of industrial enzymes: an update. *Current opinion in biotechnology*. 14, 438-443.
- Chi, Y., Hatakka, A. and Maijala, P. (2007). Can co-culturing of two white-rot fungi increase lignin degradation and the production of lignin-degrading enzymes? *International Biodeterioration and Biodegradation*. 59, 32-39.
- Chin, K., H'ng, P., Wong, L., Tey, B. and Paridah, M. (2011). Production of glucose from oil palm trunk and sawdust of rubberwood and mixed hardwood. *Applied Energy*. 88, 4222-4228.

- Christenson, A., Dimcheva, N., Ferapontova, E. E., Gorton, L., Ruzgas, T., Stoica, L., Shleev, S., Yaropolov, A. I., Haltrich, D. and Thorneley, R. N. (2004). Direct electron transfer between ligninolytic redox enzymes and electrodes. *Electroanalysis*. 16, 1074-1092.
- Christov, L., Szakacs, G. and Balakrishnan, H. (1999). Production, partial characterization and use of fungal cellulase-free xylanases in pulp bleaching. *Process Biochemistry*. 34, 511-517.
- Chung, S. Y., Maeda, M., Song, E., Horikoshij, K. and Kudo, T. (1994). A gram-positive polychlorinated biphenyl-degrading bacterium, *Rhodococcus erythropolis* strain TA421, isolated from a termite ecosystem. *Bioscience, Biotechnology and Biochemistry*. 58, 2111-2113.
- Collins, T., Gerday, C. and Feller, G. (2005). Xylanases, xylanase families and extremophilic xylanases. *FEMS Microbiology Reviews*. 29, 3-23.
- Couto, S. R., Domínguez, A. and Sanroman, A. (2001). Utilisation of lignocellulosic wastes for lignin peroxidase production by semi-solid-state cultures of *Phanerochaete chrysosporium*. *Biodegradation*. 12, 283-289.
- Couto, S. R. and Herrera, J. L. T. (2006). Industrial and biotechnological applications of laccases: A review. *Biotechnology Advances*. 24, 500-513.
- Couto, S. R., Moldes, D. and Sanroman, M. A. (2006). Optimum stability conditions of pH and temperature for ligninase and manganese-dependent peroxidase from *Phanerochaete chrysosporium*. Application to in vitro decolorization of Poly R-478 by MnP. *World Journal of Microbiology and Biotechnology*. 22, 607-612.
- Couto, S. R. and Sanromán, M. A. (2005). Application of solid-state fermentation to ligninolytic enzyme production. *Biochemical Engineering Journal*. 22, 211-219.
- Cunha, F. M., Esperança, M. N., Zangirolami, T. C., Badino, A. C. and Farinas, C. S. (2012). Sequential solid-state and submerged cultivation of *Aspergillus niger* on sugarcane bagasse for the production of cellulase. *Bioresource Technology*. 112, 270-274.
- Czaczyk, K. and Myszka, K. (2007). Biosynthesis of extracellular polymeric substances (EPS) and its role in microbial biofilm formation. *Polish Journal of Environmental Studies*. 16, 799.

- Da Silva, C. R. and Koblitz, M. G. B. (2010). Partial characterization and inactivation of peroxidases and polyphenol-oxidases of Umbu-Cajá (*Spondias* spp.). *Ciência e Tecnologia de Alimentos*. 30, 11.
- Da Silva Delabona, P., Lima, D. J., Robl, D., Rabelo, S. C., Farinas, C. S. and Da Cruz Pradella, J. G. (2016). Enhanced cellulase production by *Trichoderma harzianum* by cultivation on glycerol followed by induction on cellulosic substrates. *Journal of Industrial Microbiology and Biotechnology*. 1-10.
- Da Silva Delabona, P., Pirola, R. D. P. B., Codima, C. A., Tremacoldi, C. R., Rodrigues, A. and Farinas, C. S. (2013). Effect of initial moisture content on two Amazon rainforest *Aspergillus* strains cultivated on agro-industrial residues: Biomass-degrading enzymes production and characterization. *Industrial Crops and Products*. 42, 236-242.
- Da Silveira, C. L., Mazutti, M. A. and Salau, N. P. (2014). Modeling the microbial growth and temperature profile in a fixed-bed bioreactor. *Bioprocess and Biosystems Engineering*. 37, 1945-1954.
- Da Silveira, C. L., Mazutti, M. A. and Salau, N. P. (2015). Solid-state fermentation process model reparametrization procedure for parameters estimation using particle swarm optimization. *Journal of Chemical Technology and Biotechnology*. 91, 762-768.
- Dabhi, B., Vyas, R. and Shelat, H. (2014). Use of banana waste for the production of cellulolytic enzymes under solid substrate fermentation using bacterial consortium. *International Journal of Current Microbiology and Applied Sciences*. 3, 337-346.
- Dai, D. and Fan, M. (2015). Preparation of bio-composite from wood sawdust and gypsum. *Industrial Crops and Products*. 74, 417-424.
- Dar, M. A., Pawar, K. D., Jadhav, J. P. and Pandit, R. S. (2015). Isolation of cellulolytic bacteria from the gastro-intestinal tract of *Achatina fulica* (Gastropoda: Pulmonata) and their evaluation for cellulose biodegradation. *International Biodeterioration and Biodegradation*. 98, 73-80.

- Das, A., Paul, T., Jana, A., Halder, S. K., Ghosh, K., Maity, C., Mohapatra, P. K. D., Pati, B. R. and Mondal, K. C. (2013). Bioconversion of rice straw to sugar using multizyme complex of fungal origin and subsequent production of bioethanol by mixed fermentation of *Saccharomyces cerevisiae* MTCC 173 and *Zymomonas mobilis* MTCC 2428. *Industrial Crops and Products*. 46, 217-225.
- Dashtban, M., Schraft, H., Syed, T. A. and Qin, W. (2010). Fungal biodegradation and enzymatic modification of lignin. *International Journal of Biochemistry Molecular Biology*. 1, 36-50.
- De Almeida, M. N., Falkoski, D. L., Guimarães, V. M., Ramos, H. J. D. O., Visser, E. M., Maitan-Alfenas, G. P. and De Rezende, S. T. (2013). Characteristics of free endoglucanase and glycosidases multienzyme complex from *Fusarium verticillioides*. *Bioresource Technology*. 143, 413-422.
- De Souza, B. (2013). Microbial degradation of lignocellulosic biomass. *Sustainable Degradation of Lignocellulosic Biomass-Techniques, Applications and Commercialization*. 207-247.
- Delabona, P. D. S., Farinas, C. S., Da Silva, M. R., Azzoni, S. F. and Da Cruz Pradella, J. G. (2012). Use of a new *Trichoderma harzianum* strain isolated from the Amazon rainforest with pretreated sugar cane bagasse for on-site cellulase production. *Bioresource Technology*. 107, 517-521.
- Deswal, D., Gupta, R., Nandal, P. and Kuhad, R. C. (2014). Fungal pretreatment improves amenability of lignocellulosic material for its saccharification to sugars. *Carbohydrate Polymers*. 99, 264-269.
- Deswal, D., Khasa, Y. P. and Kuhad, R. C. (2011). Optimization of cellulase production by a brown rot fungus *Fomitopsis* sp. RCK2010 under solid state fermentation. *Bioresource Technology*. 102, 6065-6072.
- Devi, M. C. and Kumar, M. S. (2012). Isolation and screening of lignocellulose hydrolytic saprophytic fungi from dairy manure soil. *Annals of Biological Research*. 3, 1145-1152.
- Dhar, B. R., Elbeshbishy, E., Hafez, H. and Lee, H. S. (2015). Hydrogen production from sugar beet juice using an integrated biohydrogen process of dark fermentation and microbial electrolysis cell. *Bioresource Technology*. 198, 223-230.

- Dheeran, P., Nandhagopal, N., Kumar, S., Jaiswal, Y. K. and Adhikari, D. K. (2012). A novel thermostable xylanase of *Paenibacillus macerans* IIPSP3 isolated from the termite gut. *Journal of Industrial Microbiology and Biotechnology*. 39, 851-860.
- Dhillon, A., Gupta, J., Jauhari, B. and Khanna, S. (2000). A cellulase-poor, thermostable, alkalitolerant xylanase produced by *Bacillus circulans* AB 16 grown on rice straw and its application in biobleaching of eucalyptus pulp. *Bioresource Technology*. 73, 273-277.
- Dhillon, G. S., Brar, S. K., Kaur, S., Metahni, S. and M'hamdi, N. (2012). Lactoserum as a moistening medium and crude inducer for fungal cellulase and hemicellulase induction through solid-state fermentation of apple pomace. *Biomass and Bioenergy*. 41, 165-174.
- Dhillon, G. S., Kaur, S. and Brar, S. K. (2013). Perspective of apple processing wastes as low-cost substrates for bioproduction of high value products: A review. *Renewable and Sustainable Energy Reviews*. 27, 789-805.
- Dhillon, G. S., Oberoi, H. S., Kaur, S., Bansal, S. and Brar, S. K. (2011). Value-addition of agricultural wastes for augmented cellulase and xylanase production through solid-state tray fermentation employing mixed-culture of fungi. *Industrial Crops and Products*. 34, 1160-1167.
- Dhillon, S. S., Gill, R. K., Gill, S. S. and Singh, M. (2004). Studies on the utilization of citrus peel for pectinase production using fungus *Aspergillus niger*. *International Journal of Environmental Studies*. 61, 199-210.
- Dien, B. S., Sarath, G., Pedersen, J. F., Sattler, S. E., Chen, H., Funnell-Harris, D. L., Nichols, N. N. and Cotta, M. A. (2009). Improved sugar conversion and ethanol yield for forage sorghum (*Sorghum bicolor* L. Moench) lines with reduced lignin contents. *BioEnergy Research*. 2, 153-164.
- Dinis, M. J., Bezerra, R. M., Nunes, F., Dias, A. A., Guedes, C. V., Ferreira, L. M., Cone, J. W., Marques, G. S., Barros, A. R. and Rodrigues, M. A. (2009). Modification of wheat straw lignin by solid state fermentation with white-rot fungi. *Bioresource Technology*. 100, 4829-4835.
- Dong, X. Q., Yang, J. S., Zhu, N., Wang, E. T. and Yuan, H. L. (2013). Sugarcane bagasse degradation and characterization of three white-rot fungi. *Bioresource Technology*. 131, 443-451.

- Donot, F., Fontana, A., Baccou, J. and Schorr-Galindo, S. (2012). Microbial exopolysaccharides: Main examples of synthesis, excretion, genetics and extraction. *Carbohydrate Polymers*. 87, 951-962.
- Dorado, M. P., Lin, S. K. C., Koutinas, A., Du, C., Wang, R. and Webb, C. (2009). Cereal-based biorefinery development: Utilisation of wheat milling by-products for the production of succinic acid. *Journal of Biotechnology*. 143, 51-59.
- Dos Santos Barbosa, E., Perrone, D., Do Amaral Vendramini, A. L. and Leite, S. G. F. (2008). Vanillin production by *Phanerochaete chrysosporium* grown on green coconut agro-industrial husk in solid state fermentation. *BioResources*. 3, 1042-1050.
- Du, C., Lin, S. K. C., Koutinas, A., Wang, R., Dorado, P. and Webb, C. (2008). A wheat biorefining strategy based on solid-state fermentation for fermentative production of succinic acid. *Bioresource Technology*. 99, 8310-8315.
- Duan, X., Liu, S., Zhang, W., Zhang, Q. and Gao, P. (2004). Volumetric productivity improvement for endoglucanase of *Trichoderma pseudokoingii* S-38. *Journal of Applied Microbiology*. 96, 772-776.
- Dubois, M., Gilles, K. A., Hamilton, J. K., Rebers, P. and Smith, F. (1956). Colorimetric method for determination of sugars and related substances. *Analytical Chemistry*. 28, 350-356.
- Dutta, T., Sahoo, R., Sengupta, R., Ray, S. S., Bhattacharjee, A. and Ghosh, S. (2008). Novel cellulases from an extremophilic filamentous fungi *Penicillium citrinum*: Production and characterization. *Journal of Industrial Microbiology and Biotechnology*. 35, 275-282.
- Dwivedi, P., Vivekanand, V., Pareek, N., Sharma, A. and Singh, R. P. (2011a). Co-cultivation of mutant *Penicillium oxalicum* SAU E-3.510 and *Pleurotus ostreatus* for simultaneous biosynthesis of xylanase and laccase under solid-state fermentation. *New Biotechnology*. 28, 616-626.
- Dwivedi, U. N., Singh, P., Pandey, V. P. and Kumar, A. (2011b). Structure–function relationship among bacterial, fungal and plant laccases. *Journal of Molecular Catalysis B: Enzymatic*. 68, 117-128.
- Ekperigin, M. (2007). Preliminary studies of cellulase production by *Acinetobacter anitratus* and *Branhamella* sp. *African Journal of Biotechnology*. 6.

- El-Wafa, A. S., Shalash, S., Selim, N., Radwan, A. and Abdel-Salam, A. (2013). Response of broiler chicks to xylanase supplementation of corn/rye containing diets varying in metabolizable energy. *International Journal of Poultry Science*. 12, 705.
- El-Zawawy, W. K., Ibrahim, M. M., Abdel-Fattah, Y. R., Soliman, N. A. and Mahmoud, M. M. (2011). Acid and enzyme hydrolysis to convert pretreated lignocellulosic materials into glucose for ethanol production. *Carbohydrate Polymers*. 84, 865-871.
- Elisashvili, V. and Kachlishvili, E. (2009). Physiological regulation of laccase and manganese peroxidase production by white-rot Basidiomycetes. *Journal of Biotechnology*. 144, 37-42.
- Elyas, K., Mathew, A., Sukumaran, R. K., Ali, P. M., Sapna, K., Kumar, S. R. and Mol, K. R. (2010). Production optimization and properties of beta glucosidases from a marine fungus *Aspergillus*-SA 58. *New Biotechnology*. 27, 347-351.
- Endo, K., Hakamada, Y., Takizawa, S., Kubota, H., Sumitomo, N., Kobayashi, T. and Ito, S. (2001). A novel alkaline endoglucanase from an alkaliphilic *Bacillus* isolate: Enzymatic properties, and nucleotide and deduced amino acid sequences. *Applied Microbiology and Biotechnology*. 57, 109-116.
- Eyini, M., Babitha, S. and Lee, M. W. (2002). Cellulose utilization and protein productivity of some cellulolytic fungal co-cultures. *Mycobiology*. 30, 166-169.
- Fang, H., Zhao, C., Song, X. Y., Chen, M., Chang, Z. and Chu, J. (2013a). Enhanced cellulolytic enzyme production by the synergism between *Trichoderma reesei* RUT-C30 and *Aspergillus niger* NL02 and by the addition of surfactants. *Biotechnology and Bioprocess Engineering*. 18, 390-398.
- Fang, H., Zhao, C. and Song, X. Y. (2010). Optimization of enzymatic hydrolysis of steam-exploded corn stover by two approaches: Response surface methodology or using cellulase from mixed cultures of *Trichoderma reesei* RUT-C30 and *Aspergillus niger* NL02. *Bioresource Technology*. 101, 4111-4119.
- Fang, Y., Ahmed, S., Liu, S., Wang, S., Lu, M. and Jiao, Y. (2013b). Optimization of antioxidant exopolysaccharides production by *Bacillus licheniformis* in solid state fermentation. *Carbohydrate Polymers*. 98, 1377-1382.

- Farani de Souza, D., Kirst Tychanowicz, G., Giatti Marques de Souza, C. and Peralta, R. M. (2006). Co-production of ligninolytic enzymes by *Pleurotus pulmonarius* on wheat bran solid state cultures. *Journal of Basic Microbiology*. 46, 126-134.
- Farinas, C. S. (2015). Developments in solid-state fermentation for the production of biomass-degrading enzymes for the bioenergy sector. *Renewable and Sustainable Energy Reviews*. 52, 179-188.
- Faure, D., Bouillant, M. and Bally, R. (1995). Comparative study of substrates and inhibitors of *Azospirillum lipoferum* and *Pyricularia oryzae* laccases. *Applied and Environmental Microbiology*. 61, 1144-1146.
- Feng, C., Zeng, G., Huang, D., Hu, S., Zhao, M., Lai, C., Huang, C., Wei, Z. and Li, N. (2011). Effect of ligninolytic enzymes on lignin degradation and carbon utilization during lignocellulosic waste composting. *Process Biochemistry*. 46, 1515-1520.
- Ferapontova, E. E., Shleev, S., Ruzgas, T., Stoica, L., Christenson, A., Tkac, J., Yaropolov, A. I. and Gorton, L. (2005). Direct electrochemistry of proteins and enzymes. *Perspectives in Bioanalysis*. 1, 517-598.
- Foreman, P. K., Brown, D., Dankmeyer, L., Dean, R., Diener, S., Dunn-Coleman, N. S., Goedegebuur, F., Houfek, T. D., England, G. J. and Kelley, A. S. (2003). Transcriptional regulation of biomass-degrading enzymes in the filamentous fungus *Trichoderma reesei*. *Journal of Biological Chemistry*. 278, 31988-31997.
- Forootanfar, H., Faramarzi, M. A., Shahverdi, A. R. and Yazdi, M. T. (2011). Purification and biochemical characterization of extracellular laccase from the ascomycete *Paraconiothyrium variable*. *Bioresource Technology*. 102, 1808-1814.
- Frybort, S., Mauritz, R., Teischinger, A. and Müller, U. (2008). Cement bonded composites—A mechanical review. *BioResources*. 3, 602-626.
- Galante, Y., De Conti, A. and Monteverdi, R. (1998). Application of *Trichoderma* enzymes in the food and feed industries. *Trichoderma and Gliocladium*. 2, 327-342.
- Galbe, M. and Zacchi, G. (2002). A review of the production of ethanol from softwood. *Applied Microbiology and Biotechnology*. 59, 618-628.

- Garcia, N. F. L., Santos, F. R. D. S., Gonçalves, F. A., Paz, M. F. D., Fonseca, G. G. and Leite, R. S. R. (2015). Production of β -glucosidase on solid-state fermentation by *Lichtheimia ramosa* in agroindustrial residues: Characterization and catalytic properties of the enzymatic extract. *Electronic Journal of Biotechnology*. 18, 314-319.
- Gautam, S., Bundela, P., Pandey, A., Khan, J., Awasthi, M. and Sarsaiya, S. (2011). Optimization for the production of cellulase enzyme from municipal solid waste residue by two novel cellulolytic fungi. *Biotechnology Research International*. 2011, 1-8.
- Gàmez, S., González-Cabriales, J. J., Ramírez, J. A., Garrote, G. and Vázquez, M. (2006). Study of the hydrolysis of sugar cane bagasse using phosphoric acid. *Journal of Food Engineering*. 74, 78-88.
- Gervais, P. and Molin, P. (2003). The role of water in solid-state fermentation. *Biochemical Engineering Journal*. 13, 85-101.
- Gessesse, A. and Gashe, B. (1997). Production of alkaline xylanase by an alkaliphilic *Bacillus* sp. isolated from an alkalinesoda lake. *Journal of Applied Microbiology*. 83, 402-406.
- Gessesse, A. and Mamo, G. (1999). High-level xylanase production by an alkaliphilic *Bacillus* sp. by using solid-state fermentation. *Enzyme and Microbial Technology*. 25, 68-72.
- Ghani, W. A. W. A. K., Mohd, A., Da Silva, G., Bachmann, R. T., Taufiq-Yap, Y. H., Rashid, U. and Ala'a, H. (2013). Biochar production from waste rubber-wood-sawdust and its potential use in C sequestration: Chemical and physical characterization. *Industrial Crops and Products*. 44, 18-24.
- Ghodake, G. S., Kalme, S. D., Jadhav, J. P. and Govindwar, S. P. (2009). Purification and partial characterization of lignin peroxidase from *Acinetobacter calcoaceticus* NCIM 2890 and its application in decolorization of textile dyes. *Applied Biochemistry and Biotechnology*. 152, 6-14.
- Ghose, T. (1987). Measurement of cellulase activities. *Pure and Applied Chemistry*. 59, 257-268.
- Ghose, T. and Bisaria, V. S. (1987). Measurement of hemicellulase activities: Part I xylanases. *Pure and Applied Chemistry*. 59, 1739-1751.

- Glenn, J. K. and Gold, M. H. (1985). Purification and characterization of an extracellular Mn (II)-dependent peroxidase from the lignin-degrading Basidiomycete, *Phanerochaete chrysosporium*. *Archives of Biochemistry and Biophysics*. 242, 329-341.
- Goering, H. K. and Van Soest, P. J. (1970). *Forage Fiber Analyses (Apparatus, Reagents, Procedures, and Some Applications)*. Washington D. C.: U. S. Department of Agriculture Handbook.
- Goh, C. S., Tan, K. T., Lee, K. T. and Bhatia, S. (2010). Bio-ethanol from lignocellulose: Status, perspectives and challenges in Malaysia. *Bioresource Technology*. 101, 4834-4841.
- Gonçalves, F. A., Ruiz, H. A., Dos Santos, E. S., Teixeira, J. A. and De Macedo, G. R. (2015). Bioethanol production from coconuts and cactus pretreated by autohydrolysis. *Industrial Crops and Products*. 77, 1-12.
- Gottschalk, L. M. F., Oliveira, R. A. and Da Silva Bon, E. P. (2010). Cellulases, xylanases, β -glucosidase and ferulic acid esterase produced by *Trichoderma* and *Aspergillus* act synergistically in the hydrolysis of sugarcane bagasse. *Biochemical Engineering Journal*. 51, 72-78.
- Gowdhaman, D., Manaswini, V., Jayanthi, V., Dhanasri, M., Jeyalakshmi, G., Gunasekar, V., Sugumaran, K. and Ponnusami, V. (2014). Xylanase production from *Bacillus aerophilus* KGJ2 and its application in xylooligosaccharides preparation. *International Journal of Biological Macromolecules*. 64, 90-98.
- Gowthaman, M., Krishna, C. and Moo-Young, M. (2001). Fungal solid state fermentation—an overview. *Applied Mycology and Biotechnology*. 1, 305-352.
- Grujić, M., Dojnov, B., Potočnik, I., Duduk, B. and Vujčić, Z. (2015). Spent mushroom compost as substrate for the production of industrially important hydrolytic enzymes by fungi *Trichoderma* spp. and *Aspergillus niger* in solid state fermentation. *International Biodeterioration and Biodegradation*. 104, 290-298.
- Guan, G., Zhang, Z., Ding, H., Li, M., Shi, D., Zhu, M. and Xia, L. (2015). Enhanced degradation of lignin in corn stalk by combined method of *Aspergillus oryzae* solid state fermentation and H₂O₂ treatment. *Biomass and Bioenergy*. 81, 224-233.

UNIVERSITI TEKNOLOGI MALAYSIA

DECLARATION OF THESIS / ~~UNDERGRADUATE PROJECT PAPER~~ AND COPYRIGHT

Author's full name : **NORATIQA BINTI KAMSANI**

Date of birth : **13TH SEPTEMBER 1987**

Title : **LIGNOCELLULOLYTIC ENZYMES BY *Aspergillus* sp. A1 AND *Bacillus* sp. B1 ISOLATED FROM GUT OF *Bulbitermes* sp. IN SOLID STATE FERMENTATION USING SAWDUST AS SUBSTRATE**

Academic Session : **2016/2017 (1)**

I declare that this thesis is classified as :

- CONFIDENTIAL** (Contains confidential information under the Official Secret Act 1972)*
- RESTRICTED** (Contains restricted information as specified by the organization where research was done)*
- OPEN ACCESS** I agree that my thesis to be published as online open access (full text)

I acknowledged that Universiti Teknologi Malaysia reserves the right as follows:


1. The thesis is the property of Universiti Teknologi Malaysia.
2. The Library of Universiti Teknologi Malaysia has the right to make copies for the purpose of research only.
3. The Library has the right to make copies of the thesis for academic exchange.



SIGNATURE
870913-05-5364
(NEW IC NO. /PASSPORT NO.)

Date : 17th January 2017

Certified by :



SIGNATURE OF SUPERVISOR
P.M. DR MADIHAH MD. SALLEH
NAME OF SUPERVISOR

Date : 17th January 2017

NOTES : * If the thesis is CONFIDENTIAL or RESTRICTED, please attach with the letter from the organization with period and reasons for confidentiality or restriction.

- Guerra-Rodríguez, E., Portilla-Rivera, O. M., Jarquín-Enríquez, L., Ramírez, J. A. and Vázquez, M. (2012). Acid hydrolysis of wheat straw: A kinetic study. *Biomass and Bioenergy*. 36, 346-355.
- Gupta, N., Yadav, K. K. and Kumar, V. (2015a). A review on current status of municipal solid waste management in India. *Journal of Environmental Sciences*. 37, 206-217.
- Gupta, R., Mehta, G., Khasa, Y. P. and Kuhad, R. C. (2011). Fungal delignification of lignocellulosic biomass improves the saccharification of celluloses. *Biodegradation*. 22, 797-804.
- Gupta, V., Garg, S., Capalash, N., Gupta, N. and Sharma, P. (2015b). Production of thermo-alkali-stable laccase and xylanase by co-culturing of *Bacillus* sp. and *Bacillus halodurans* for biobleaching of kraft pulp and deinking of waste paper. *Bioprocess and Biosystems Engineering*. 38, 947-956.
- Gutierrez-Correa, M., Portal, L., Moreno, P. and Tengerdy, R. P. (1999). Mixed culture solid substrate fermentation of *Trichoderma reesei* with *Aspergillus niger* on sugar cane bagasse. *Bioresource Technology*. 68, 173-178.
- Gutierrez-Correa, M. and Tengerdy, R. P. (1997). Production of cellulase on sugar cane bagasse by fungal mixed culture solid substrate fermentation. *Biotechnology Letters*. 19, 665-667.
- Gutierrez-Correa, M. and Tengerdy, R. P. (1998). Xylanase production by fungal mixed culture solid substrate fermentation on sugar cane bagasse. *Biotechnology Letters*. 20, 45-47.
- Haki, G. and Rakshit, S. (2003). Developments in industrially important thermostable enzymes: A review. *Bioresource Technology*. 89, 17-34.
- Hamid, S. B. A., Shilpy, M. Z. and Ali, M. (2014). Green catalytic approach for the synthesis of platform chemicals from palm tree lignin. *Advanced Materials Research*. 62-66.
- Hamzah, F., Idris, A. and Shuan, T. K. (2011). Preliminary study on enzymatic hydrolysis of treated oil palm (*Elaeis*) empty fruit bunches fibre (EFB) by using combination of cellulase and β -1,4-glucosidase. *Biomass and Bioenergy*. 35, 1055-1059.

- Han, W. and He, M. (2010). The application of exogenous cellulase to improve soil fertility and plant growth due to acceleration of straw decomposition. *Bioresource Technology*. 101, 3724-3731.
- Han, Y. and Chen, H. (2008). Characterization of β -glucosidase from corn stover and its application in simultaneous saccharification and fermentation. *Bioresource Technology*. 99, 6081-6087.
- Hansen, G. H., Lübeck, M., Frisvad, J. C., Lübeck, P. S. and Andersen, B. (2015). Production of cellulolytic enzymes from ascomycetes: Comparison of solid state and submerged fermentation. *Process Biochemistry*. 50, 1327-1341.
- Haq, I., Kumar, S., Kumari, V., Singh, S. K. and Raj, A. (2016). Evaluation of bioremediation potentiality of ligninolytic *Serratia liquefaciens* for detoxification of pulp and paper mill effluent. *Journal of Hazardous Materials*. 305, 190-199.
- Harazono, K., Yamashita, N., Shinzato, N., Watanabe, Y., Fukatsu, T. and Kurane, R. (2003). Isolation and characterization of aromatics-degrading microorganisms from the gut of the lower termite *Coptotermes formosanus*. *Bioscience, Biotechnology, and Biochemistry*. 67, 889-892.
- Hariharan, S. and Nambisan, P. (2012). Optimization of lignin peroxidase, manganese peroxidase and laccase production from *Ganoderma lucidum* under solid state fermentation of pineapple leaf. *BioResources*. 8, 250-271.
- Harman, G. E. and Kubicek, C. P. (2002). *Trichoderma and Gliocladium: Enzymes, Biological Control and Commercial Applications*. London, U. K.: Taylor & Francis.
- Harris, A. D. and Ramalingam, C. (2010). Xylanases and its application in food industry: A review. *Journal of Experimental Sciences*. 1.
- Harrold, Z. R., Hertel, M. R. and Gorman-Lewis, D. (2011). Optimizing *Bacillus subtilis* spore isolation and quantifying spore harvest purity. *Journal of Microbiological Methods*. 87, 325-329.
- Heck, J. X., Flôres, S. H., Hertz, P. F. and Ayub, M. A. Z. (2005). Optimization of cellulase-free xylanase activity produced by *Bacillus coagulans* BL69 in solid-state cultivation. *Process Biochemistry*. 40, 107-112.
- Heller, A. (2004). Miniature biofuel cells. *Physical Chemistry Chemical Physics*. 6, 209-216.

- Hendriks, A. T. W. M. and Zeeman, G. (2009). Pretreatments to enhance the digestibility of lignocellulosic biomass. *Bioresource Technology*. 100, 10-18.
- Henry, R. J. (2010). Evaluation of plant biomass resources available for replacement of fossil oil. *Plant Biotechnology Journal*. 8, 288-293.
- Heo, H. S., Park, H. J., Park, Y. K., Ryu, C., Suh, D. J., Suh, Y. W., Yim, J. H. and Kim, S. S. (2010). Bio-oil production from fast pyrolysis of waste furniture sawdust in a fluidized bed. *Bioresource Technology*. 101, 91-96.
- Himmel, M. E., Ding, S. Y., Johnson, D. K., Adney, W. S., Nimlos, M. R., Brady, J. W. and Foust, T. D. (2007). Biomass recalcitrance: Engineering plants and enzymes for biofuels production. *Science*. 315, 804-807.
- Hodge, D. B., Karim, M. N., Schell, D. J. and McMillan, J. D. (2008). Soluble and insoluble solids contributions to high-solids enzymatic hydrolysis of lignocellulose. *Bioresource Technology*. 99, 8940-8948.
- Hoi, W. (2003). *Wood Waste to Energy-from Waste to Wealth with Special Reference to Malaysia*. [Brochure] Kuala Lumpur: Forest Research Institute of Malaysia.
- Hölker, U. and Lenz, J. (2005). Solid-state fermentation—are there any biotechnological advantages? *Current Opinion in Microbiology*. 8, 301-306.
- Hopkins, J. D. (2003). *Subterranean Termite Identification and Biology*. United States: University of Arkansas.
- Hossain, S. M. and Anantharaman, N. (2006). Activity enhancement of ligninolytic enzymes of *Trametes versicolor* with bagasse powder. *African Journal of Biotechnology*. 5, 189.
- Howell, J. and Stuck, J. (1975). Kinetics of solka floc cellulose hydrolysis by *Trichoderma viride* cellulase. *Biotechnology and Bioengineering*. 17, 873-893.
- Hsieh, J. W., Wu, H. S., Wei, Y. H. and Wang, S. S. (2007). Determination and kinetics of producing glucosamine using fungi. *Biotechnology Progress*. 23, 1009-1016.
- Hsu, T. C., Guo, G. L., Chen, W. H. and Hwang, W. S. (2010). Effect of dilute acid pretreatment of rice straw on structural properties and enzymatic hydrolysis. *Bioresource Technology*. 101, 4907-4913.

- Hu, H., Van den Brink, J., Gruben, B., Wösten, H., Gu, J. D. and De Vries, R. (2011). Improved enzyme production by co-cultivation of *Aspergillus niger* and *Aspergillus oryzae* and with other fungi. *International Biodeterioration and Biodegradation*. 65, 248-252.
- Huang, S., Sheng, P. and Zhang, H. (2012). Isolation and identification of cellulolytic bacteria from the gut of *Holotrichia parallela* larvae (Coleoptera: Scarabaeidae). *International Journal of Molecular Sciences*. 13, 2563-2577.
- Hung, F. C. (2016). Linking forest naturalness and human wellbeing—a study on public's experiential connection to remnant forests within a highly urbanized region in Malaysia. *Urban Forestry and Urban Greening*. 16, 13-24.
- Hung, K. S., Liu, S. M., Tzou, W. S., Lin, F. P., Pan, C. L., Fang, T. Y., Sun, K. H. and Tang, S. J. (2011). Characterization of a novel GH10 thermostable, halophilic xylanase from the marine bacterium *Thermoanaerobacterium saccharolyticum* NTOU1. *Process Biochemistry*. 46, 1257-1263.
- Husaini, A., Fisol, F. A., Yun, L. C., Hussain, M. H. and Roslan, H. A. (2011). Lignocellulolytic enzymes produced by tropical white rot fungi during biopulping of *Acacia mangium* wood chips. *Journal of Biochemical Technology*. 3, 245-250.
- Hyodo, F., Inoue, T., Azuma, J. I., Tayasu, I. and Abe, T. (2000). Role of the mutualistic fungus in lignin degradation in the fungus-growing termite *Macrotermes gilvus* (Isoptera; Macrotermitinae). *Soil Biology and Biochemistry*. 32, 653-658.
- Ibrahim, N., El-Badry, K., Eid, B. and Hassan, T. (2011). A new approach for biofinishing of cellulose-containing fabrics using acid cellulases. *Carbohydrate Polymers*. 83, 116-121.
- Ikram-ul-Haq, M. M. J., Khan, S. and Siddiq, T. Z. (2005). Cotton saccharifying activity of cellulases produced by co-culture of *Aspergillus niger* and *Trichoderma viride*. *Research Journal of Agriculture and Biological Sciences*. 1, 241-245.
- Inoue, T., Murashima, K., Azuma, J. I., Sugimoto, A. and Slaytor, M. (1997). Cellulose and xylan utilisation in the lower termite *Reticulitermes speratus*. *Journal of Insect Physiology*. 43, 235-242.

- Irfan, M., Nadeem, M. and Syed, Q. (2014). One-factor-at-a-time (OFAT) optimization of xylanase production from *Trichoderma viride*-IR05 in solid-state fermentation. *Journal of Radiation Research and Applied Sciences*. 7, 317-326.
- Irfan, M., Safdar, A., Syed, Q. and Nadeem, M. (2012). Isolation and screening of cellulolytic bacteria from soil and optimization of cellulase production and activity. *Turkish Journal of Biochemistry*. 37, 287-293.
- Irshad, M. and Asgher, M. (2013). Production and optimization of ligninolytic enzymes by white rot fungus *Schizophyllum commune* IBL-06 in solid state medium banana stalks. *African Journal of Biotechnology*. 10, 18234-18242.
- Ivanka, S., Albert, K. and Veselin, S. (2010). Properties of crude laccase from *Trametes versicolor* produced by solid-substrate fermentation. *Advances in Bioscience and Biotechnology*. 1, 208-215.
- Jabasingh, S. A. and Nachiyar, C. V. (2011). Utilization of pretreated bagasse for the sustainable bioproduction of cellulase by *Aspergillus nidulans* MTCC344 using response surface methodology. *Industrial Crops and Products*. 34, 1564-1571.
- Janusz, G., Czurylo, A., Frąc, M., Rola, B., Sulej, J., Pawlik, A., Siwulski, M. and Rogalski, J. (2015). Laccase production and metabolic diversity among *Flammulina velutipes* strains. *World Journal of Microbiology and Biotechnology*. 31, 121-133.
- Jayaraman, S. and Gunasekaran, M. (1990). Influence of solvent treatment of alkaline peat extracts on the growth and the enzyme activities of *Bacillus subtilis*. *Resources, Conservation and Recycling*. 4, 297-304.
- Jing, L., Zhao, S., Xue, J. L., Zhang, Z., Yang, Q., Xian, L. and Feng, J. X. (2015). Isolation and characterization of a novel *Penicillium oxalicum* strain Z1-3 with enhanced cellobiohydrolase production using cellulase-hydrolyzed sugarcane bagasse as carbon source. *Industrial Crops and Products*. 77, 666-675.
- Jørgensen, H., Kristensen, J. B. and Felby, C. (2007). Enzymatic conversion of lignocellulose into fermentable sugars: Challenges and opportunities. *Biofuels, Bioproducts and Biorefining*. 1, 119-134.

- Juhasz, T., Szengyel, Z., Reczey, K., Siika-Aho, M. and Viikari, L. (2005). Characterization of cellulases and hemicellulases produced by *Trichoderma reesei* on various carbon sources. *Process Biochemistry*. 40, 3519-3525.
- Juturu, V. and Wu, J. C. (2014). Microbial cellulases: Engineering, production and applications. *Renewable and Sustainable Energy Reviews*. 33, 188-203.
- Kallel, F., Driss, D., Chaari, F., Zouari-Ellouzi, S., Chaabouni, M., Ghorbel, R. and Chaabouni, S. E. (2016). Statistical optimization of low-cost production of an acidic xylanase by *Bacillus mojavensis* UEB-FK: Its potential applications. *Biocatalysis and Agricultural Biotechnology*. 5, 1-10.
- Kalogeris, E., Iniotaki, F., Topakas, E., Christakopoulos, P., Kekos, D. and Macris, B. (2003). Performance of an intermittent agitation rotating drum type bioreactor for solid-state fermentation of wheat straw. *Bioresource Technology*. 86, 207-213.
- Kalyani, D., Lee, K. M., Kim, T. S., Li, J., Dhiman, S. S., Kang, Y. C. and Lee, J. K. (2013). Microbial consortia for saccharification of woody biomass and ethanol fermentation. *Fuel*. 107, 815-822.
- Kamble, R. D. and Jadhav, A. R. (2012). Isolation, purification and characterization of xylanase produced by a new species of *Bacillus* in solid state fermentation. *International Journal of Microbiology*. 2012, 1-8.
- Kamsani, N., Salleh, M. M., Yahya, A. and Chong, C. S. (2016). Production of lignocellulolytic enzymes by microorganisms isolated from *Bulbitermes* sp. termite gut in solid-state fermentation. *Waste and Biomass Valorization*. 7, 357-371.
- Kana, E. G., Oloke, J., Lateef, A. and Adesiyun, M. (2012). Modeling and optimization of biogas production on saw dust and other co-substrates using artificial neural network and genetic algorithm. *Renewable Energy*. 46, 276-281.
- Kanayama, N., Tohru, S. and Keiichi, K. (2002). Purification and characterization of an alkaline manganese peroxidase from *Aspergillus terreus* LD-1. *Journal of Bioscience and Bioengineering*. 93, 405-410.
- Kang, S., Park, Y., Lee, J., Hong, S. and Kim, S. (2004). Production of cellulases and hemicellulases by *Aspergillus niger* KK2 from lignocellulosic biomass. *Bioresource Technology*. 91, 153-156.

- Karimi, K., Kheradmandinia, S. and Taherzadeh, M. J. (2006). Conversion of rice straw to sugars by dilute-acid hydrolysis. *Biomass and Bioenergy*. 30, 247-253.
- Karmakar, M. and Ray, R. (2011). Current trends in research and application of microbial cellulases. *Research Journal of Microbiology*. 6, 41.
- Karp, S. G., Faraco, V., Amore, A., Birolo, L., Giangrande, C., Soccol, V. T., Pandey, A. and Soccol, C. R. (2012). Characterization of laccase isoforms produced by *Pleurotus ostreatus* in solid state fermentation of sugarcane bagasse. *Bioresource Technology*. 114, 735-739.
- Kato, K., Kozaki, S. and Sakuranaga, M. (1998). Degradation of lignin compounds by bacteria from termite guts. *Biotechnology Letters*. 20, 459-462.
- Kausar, H., Sariah, M., Mohd Saud, H., Zahangir Alam, M. and Razi Ismail, M. (2010). Development of compatible lignocellulolytic fungal consortium for rapid composting of rice straw. *International Biodeterioration and Biodegradation*. 64, 594-600.
- Khan, M. I. M., Sajjad, M., Sadaf, S., Zafar, R., Niazi, U. H. and Akhtar, M. W. (2013). The nature of the carbohydrate binding module determines the catalytic efficiency of xylanase Z of *Clostridium thermocellum*. *Journal of Biotechnology*. 168, 403-408.
- Khandeparkar, R. and Bhosle, N. (2006). Isolation, purification and characterization of the xylanase produced by *Arthrobacter* sp. MTCC 5214 when grown in solid-state fermentation. *Enzyme and Microbial Technology*. 39, 732-742.
- Khasin, A., Alchanati, I. and Shoham, Y. (1993). Purification and characterization of a thermostable xylanase from *Bacillus stearothermophilus* T-6. *Applied and Environmental Microbiology*. 59, 1725-1730.
- Khelil, O., Choubane, S. and Cheba, B. A. (2015). Co-production of cellulases and manganese peroxidases by *Bacillus* sp. R2 and *Bacillus cereus* 11778 on waste newspaper: Application in dyes decolourization. *Procedia Technology*. 19, 980-987.
- Kheng, P. P. and Omar, I. C. (2005). Xylanase production by a local fungal isolate, *Aspergillus niger* USM A1 via solid state fermentation using palm kernel cake (PKC) as substrate. *Songklanakarin Journal of Science and Technology*. 27, 325-336.

- Kim, O. S., Cho, Y. J., Lee, K., Yoon, S. H., Kim, M., Na, H., Park, S. C., Jeon, Y. S., Lee, J. H. and Yi, H. (2012). Introducing EzTaxon-e: A prokaryotic 16S rRNA gene sequence database with phylotypes that represent uncultured species. *International Journal of Systematic and Evolutionary Microbiology*. 62, 716-721.
- Ko, J. K., Ko, H., Kim, K. H. and Choi, I. G. (2016). Characterization of the biochemical properties of recombinant Xyn10C from a marine bacterium, *Saccharophagus degradans* 2-40. *Bioprocess and Biosystems Engineering*. DOI 10.1007/s00449-016-1548-2.
- Kolasa, M., Ahring, B. K., Lübeck, P. S. and Lübeck, M. (2014). Co-cultivation of *Trichoderma reesei* RutC30 with three black *Aspergillus* strains facilitates efficient hydrolysis of pretreated wheat straw and shows promises for on-site enzyme production. *Bioresource Technology*. 169, 143-148.
- König, H., Fröhlich, J. and Hertel, H. (2006). Diversity and Lignocellulolytic Activities of Cultured Microorganisms. In König, H. and Varma, A. (Eds) *Intestinal Microorganisms of Termites and other Invertebrates* (pp. 271-301). Berlin: Springer.
- Kont, R., Kurašin, M., Teugjas, H. and Väljamäe, P. (2013). Strong cellulase inhibitors from the hydrothermal pretreatment of wheat straw. *Biotechnology and Biofuels*. 6, 135.
- Koroiva, R., Souza, C., Toyama, D., Henrique-Silva, F. and Fonseca-Gessner, A. (2013). Lignocellulolytic enzymes and bacteria associated with the digestive tracts of *Stenochironomus* (Diptera: Chironomidae) larvae. *Genetics and Molecular Research*. 12, 3421-3434.
- Kotchoni, O., Shonukan, O. and Gachomo, W. (2003). *Bacillus pumilus* BpCRI 6, a promising candidate for cellulase production under conditions of catabolite repression. *African Journal of Biotechnology*. 2, 140-146.
- Koyani, R. D., Sanghvi, G. V., Sharma, R. K. and Rajput, K. S. (2013). Contribution of lignin degrading enzymes in decolourisation and degradation of reactive textile dyes. *International Biodeterioration and Biodegradation*. 77, 1-9.
- Krause, M., Beauchemin, K., Rode, L., Farr, B. and Nørgaard, P. (1998). Fibrolytic enzyme treatment of barley grain and source of forage in high-grain diets fed to growing cattle. *Journal of Animal Science*. 76, 2912-2920.

- Krishna, C. (1999). Production of bacterial cellulases by solid state bioprocessing of banana wastes. *Bioresource Technology*. 69, 231-239.
- Krishna, C. and Chandrasekaran, M. (1996). Banana waste as substrate for α -amylase production by *Bacillus subtilis* (CBTK 106) under solid-state fermentation. *Applied Microbiology and Biotechnology*. 46, 106-111.
- Kudo, T. (2009). Termite-microbe symbiotic system and its efficient degradation of lignocellulose. *Bioscience, Biotechnology and Biochemistry*. 73, 2561-2567.
- Kuhad, R. C., Deswal, D., Sharma, S., Bhattacharya, A., Jain, K. K., Kaur, A., Pletschke, B. I., Singh, A. and Karp, M. (2016). Revisiting cellulase production and redefining current strategies based on major challenges. *Renewable and Sustainable Energy Reviews*. 55, 249-272.
- Kuhad, R. C., Gupta, R., Khasa, Y. P. and Singh, A. (2010). Bioethanol production from *Lantanacamura* (red sage): Pretreatment, saccharification and fermentation. *Bioresource Technology*. 101, 8348-8354.
- Kuhad, R. C. and Singh, A. (1993). Lignocellulose biotechnology: Current and future prospects. *Critical Reviews in Biotechnology*. 13, 151-172.
- Kuhar, F., Castiglia, V. and Levin, L. (2015). Enhancement of laccase production and malachite green decolorization by co-culturing *Ganoderma lucidum* and *Trametes versicolor* in solid-state fermentation. *International Biodeterioration and Biodegradation*. 104, 238-243.
- Kumar, A., Gaiind, S. and Nain, L. (2008a). Evaluation of thermophilic fungal consortium for paddy straw composting. *Biodegradation*. 19, 395-402.
- Kumar, R., Mago, G., Balan, V. and Wyman, C. E. (2009). Physical and chemical characterizations of corn stover and poplar solids resulting from leading pretreatment technologies. *Bioresource Technology*. 100, 3948-3962.
- Kumar, R., Singh, S. and Singh, O. V. (2008b). Bioconversion of lignocellulosic biomass: Biochemical and molecular perspectives. *Journal of Industrial Microbiology and Biotechnology*. 35, 377-391.
- Kumar, R. and Wyman, C. E. (2009). Effect of xylanase supplementation of cellulase on digestion of corn stover solids prepared by leading pretreatment technologies. *Bioresource Technology*. 100, 4203-4213.

- Kumar, S., Jain, K. K., Bhardwaj, K. N., Chakraborty, S. and Kuhad, R. C. (2015). Multiple genes in a single host: Cost-effective production of bacterial laccase (cotA), pectate lyase (pel) and endoxylanase (xyl) by simultaneous expression and cloning in single vector in *E. coli*. *PLoS One*. DOI:10.1371/journal.pone.0144379.
- Kumar, V. and Satyanarayana, T. (2014). Production of thermo-alkali-stable xylanase by a novel polyextremophilic *Bacillus halodurans* TSEV1 in cane molasses medium and its applicability in making whole wheat bread. *Bioprocess and Biosystems Engineering*. 37, 1043-1053.
- Kunamneni, A., Ghazi, I., Camarero, S., Ballesteros, A., Plou, F. J. and Alcalde, M. (2008). Decolorization of synthetic dyes by laccase immobilized on epoxy-activated carriers. *Process Biochemistry*. 43, 169-178.
- Kurakake, M., Ide, N. and Komaki, T. (2007). Biological pretreatment with two bacterial strains for enzymatic hydrolysis of office paper. *Current Microbiology*. 54, 424-428.
- Kuwahara, M., Glenn, J. K., Morgan, M. A. and Gold, M. H. (1984). Separation and characterization of two extracellular H₂O₂-dependent oxidases from ligninolytic cultures of *Phanerochaete chrysosporium*. *FEBS Letters*. 169, 247-250.
- Laemmli, U. K. (1970). Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature*. 227, 680-685.
- Lah, N., Rahman, N. and Nama, M. (2012). Cellulase activity and glucose production by *Bacillus cereus* monoculture and co-culture utilizing palm kernel cake (PKC) under solid state fermentation. *International Conference on Environment, Energy and Biotechnology IPCBEE*. 33, 172-177.
- Lakshmi, G. S., Rao, C. S., Rao, R. S., Hobbs, P. J. and Prakasham, R. S. (2009). Enhanced production of xylanase by a newly isolated *Aspergillus terreus* under solid state fermentation using palm industrial waste: A statistical optimization. *Biochemical Engineering Journal*. 48, 51-57.
- Lan, T. Q., Wei, D., Yang, S. T. and Liu, X. (2013). Enhanced cellulase production by *Trichoderma viride* in a rotating fibrous bed bioreactor. *Bioresour. Technol.* 133, 175-182.

- Lasure, L. L. and Zhang, M. (2004). Bioconversion and biorefineries of the future. *Draft Report from the Pacific Northwest National Laboratory and National Renewable Energy Lab.*
- Laureano-Perez, L., Teymouri, F., Alizadeh, H. and Dale, B. E. (2005). Understanding factors that limit enzymatic hydrolysis of biomass. *Applied Biochemistry and Biotechnology*. 124, 1081-1099.
- Le Roes-Hill, M., Rohland, J. and Burton, S. (2011). Actinobacteria isolated from termite guts as a source of novel oxidative enzymes. *Antonie van Leeuwenhoek*. 100, 589-605.
- Lee, J., Gwak, K., Park, J., Park, M., Choi, D., Kwon, M. and Choi, I. (2007). Biological pretreatment of softwood *Pinus densiflora* by three white rot fungi. *Journal of Microbiology Seoul*. 45, 485.
- Lee, J., Lee, S., Hong, E., Jeung, E., Kang, H., Kim, M. and Choi, I. (2006). Estrogenic reduction of styrene monomer degraded by *Phanerochaete chrysosporium* KFRI 20742. *Journal of Microbiology Seoul*. 44, 177.
- Lenartovicz, V., Marques De Souza, C. G., Moreira, F. G. and Peralta, R. M. (2002). Temperature effect in the production of multiple xylanases by *Aspergillus fumigatus*. *Journal of Basic Microbiology*. 42, 388-395.
- Lenting, H. and Warmoeskerken, M. (2001). Mechanism of interaction between cellulase action and applied shear force, an hypothesis. *Journal of Biotechnology*. 89, 217-226.
- Levin, L., Herrmann, C. and Papinutti, V. L. (2008). Optimization of lignocellulolytic enzyme production by the white-rot fungus *Trametes trogii* in solid-state fermentation using response surface methodology. *Biochemical Engineering Journal*. 39, 207-214.
- Lewis, V. (2008). Isoptera. In Resh, V. H. and Cardé R. T. (Eds.) *Encyclopedia of Insects*. (pp. 531-534). United Kingdom: Academic Press.
- Li, A. H., Lin, C. W. and Tran, D. T. (2011). Optimizing the response surface for producing ethanol from avicel by *Brevibacillus* strain AHPC8120. *Journal of The Taiwan Institute of Chemical Engineers*. 42, 787-792.
- Li, C., Wang, Q. and Zhao, Z. K. (2008). Acid in ionic liquid: An efficient system for hydrolysis of lignocellulose. *Green Chemistry*. 10, 177-182.

- Li, C., Yang, Z., Zhang, R. H. C., Zhang, D., Chen, S. and Ma, L. (2013). Effect of pH on cellulase production and morphology of *Trichoderma reesei* and the application in cellulosic material hydrolysis. *Journal of Biotechnology*. 168, 470-477.
- Li, H., Xue, Y., Wu, J., Wu, H., Qin, G., Li, C., Ding, J., Liu, J., Gan, L. and Long, M. (2016). Enzymatic hydrolysis of hemicelluloses from *Miscanthus* to monosaccharides or xylo-oligosaccharides by recombinant hemicellulases. *Industrial Crops and Products*. 79, 170-179.
- Liang, Y., Yesuf, J., Schmitt, S., Bender, K. and Bozzola, J. (2009). Study of cellulases from a newly isolated thermophilic and cellulolytic *Brevibacillus* sp. strain JXL. *Journal of Industrial Microbiology and Biotechnology*. 36, 961-970.
- Liao, H., Fan, X., Mei, X., Wei, Z., Raza, W., Shen, Q. and Xu, Y. (2015). Production and characterization of cellulolytic enzyme from *Penicillium oxalicum* GZ-2 and its application in lignocellulose saccharification. *Biomass and Bioenergy*. 74, 122-134.
- Liew, C., Husaini, A., Hussain, H., Muid, S., Liew, K. and Roslan, H. (2011). Lignin biodegradation and ligninolytic enzyme studies during biopulping of *Acacia mangium* wood chips by tropical white rot fungi. *World Journal of Microbiology and Biotechnology*. 27, 1457-1468.
- Lim, J. S., Manan, Z. A., Alwi, S. R. W. and Hashim, H. (2012). A review on utilisation of biomass from rice industry as a source of renewable energy. *Renewable and Sustainable Energy Reviews*. 16, 3084-3094.
- Lin, H., Wang, B., Zhuang, R., Zhou, Q. and Zhao, Y. (2011). Artificial construction and characterization of a fungal consortium that produces cellulolytic enzyme system with strong wheat straw saccharification. *Bioresource Technology*. 102, 10569-10576.
- Lin, Y., Zhang, Z., Tian, Y., Zhao, W., Zhu, B., Xu, Z., Peng, R. and Yao, Q. (2013). Purification and characterization of a novel laccase from *Coprinus cinereus* and decolorization of different chemically dyes. *Molecular Biology Reports*. 40, 1487-1494.
- Ling, T. P., Hassan, O., Badri, K., Maskat, M. Y. and Mustapha, W. A. W. (2013). Sugar recovery of enzymatic hydrolysed oil palm empty fruit bunch fiber by chemical pretreatment. *Cellulose*. 20, 3191-3203.

- Liu, D., Zhang, R., Yang, X., Wu, H., Xu, D., Tang, Z. and Shen, Q. (2011). Thermostable cellulase production of *Aspergillus fumigatus* Z5 under solid-state fermentation and its application in degradation of agricultural wastes. *International Biodeterioration and Biodegradation*. 65, 717-725.
- Liu, J., Yuan, X., Zeng, G., Shi, J. and Chen, S. (2006). Effect of biosurfactant on cellulase and xylanase production by *Trichoderma viride* in solid substrate fermentation. *Process Biochemistry*. 41, 2347-2351.
- Lo, Y. C., Lu, W. C., Chen, C. Y., Chen, W. M. and Chang, J. S. (2010). Characterization and high-level production of xylanase from an indigenous cellulolytic bacterium *Acinetobacter junii* F6-02 from southern Taiwan soil. *Biochemical Engineering Journal*. 53, 77-84.
- Lochner, A., Giannone, R. J., Rodriguez, M., Shah, M. B., Mielenz, J. R., Keller, M., Antranikian, G., Graham, D. E. and Hettich, R. L. (2011). Use of label-free quantitative proteomics to distinguish the secreted cellulolytic systems of *Caldicellulosiruptor bescii* and *Caldicellulosiruptor obsidiansis*. *Applied and Environmental Microbiology*. 77, 4042-4054.
- Lončar, N., Gligorijević, N., Božić, N. and Vujčić, Z. (2014). Congo red degrading laccases from *Bacillus amyloliquefaciens* strains isolated from salt spring in Serbia. *International Biodeterioration and Biodegradation*. 91, 18-23.
- Lopretti, M., Mathias, A. and Rodrigues, A. (1993). Activity of ligninase peroxidase from *Acinetobacter anitratus* and the degradation of *Pinus pinaster* lignin. *Process Biochemistry*. 28, 543-547.
- Lowry, O. H., Rosebrough, N. J., Farr, A. L. and Randall, R. J. (1951). Protein measurement with the Folin phenol reagent. *Journal of Biological Chemistry*. 193, 265-275.
- Lykidis, A., Mavromatis, K., Ivanova, N., Anderson, I., Land, M., DiBartolo, G., Martinez, M., Lapidus, A., Lucas, S. and Copeland, A. (2007). Genome sequence and analysis of the soil cellulolytic actinomycete *Thermobifida fusca* YX. *Journal of Bacteriology*. 189, 2477-2486.
- Lynd, L. R., Laser, M. S., Bransby, D., Dale, B. E., Davison, B., Hamilton, R., Himmel, M., Keller, M., McMillan, J. D. and Sheehan, J. (2008). How biotech can transform biofuels. *Nature Biotechnology*. 26, 169-172.

- Lynd, L. R., Weimer, P. J., Van Zyl, W. H. and Pretorius, I. S. (2002). Microbial cellulose utilization: Fundamentals and biotechnology. *Microbiology and Molecular Biology Reviews*. 66, 506-577.
- Ma, F., Wang, J., Zeng, Y., Yu, H., Yang, Y. and Zhang, X. (2011). Influence of the co-fungal treatment with two white rot fungi on the lignocellulosic degradation and thermogravimetry of corn stover. *Process Biochemistry*. 46, 1767-1773.
- Ma, K. and Ruan, Z. (2015). Production of a lignocellulolytic enzyme system for simultaneous bio-delignification and saccharification of corn stover employing co-culture of fungi. *Bioresource Technology*. 175, 586-593.
- Ma, L., Yang, W., Meng, F., Ji, S., Xin, H. and Cao, B. (2015). Characterization of an acidic cellulase produced by *Bacillus subtilis* BY-4 isolated from gastrointestinal tract of Tibetan pig. *Journal of The Taiwan Institute of Chemical Engineers*. 56, 67-72.
- Ma, R. J., Wang, C. Y., Liu, Y. W., Sivakumar, T. R., Ren, Z. X., Fang, Y., Jia, J. Q. and Gui, Z. Z. (2014). Identification and characterization of a novel endoglucanase (CMCase) isolated from the larval gut of *Bombyx mori*. *Journal of Asia-Pacific Entomology*. 17, 67-71.
- Madlala, A. M., Bissoon, S., Singh, S. and Christov, L. (2001). Xylanase-induced reduction of chlorine dioxide consumption during elemental chlorine-free bleaching of different pulp types. *Biotechnology Letters*. 23, 345-351.
- Magalhães, P. O. and Milagres, A. M. F. (2009). Biochemical properties of a β -mannanase and a β -xylanase produced by *Ceriporiopsis subvermispora* during biopulping conditions. *International Biodeterioration and Biodegradation*. 63, 191-195.
- Mahanta, N., Gupta, A. and Khare, S. (2008). Production of protease and lipase by solvent tolerant *Pseudomonas aeruginosa* PseA in solid-state fermentation using *Jatropha curcas* seed cake as substrate. *Bioresource Technology*. 99, 1729-1735.
- Maki, M., Leung, K. T. and Qin, W. (2009). The prospects of cellulase-producing bacteria for the bioconversion of lignocellulosic biomass. *International Journal of Biological Sciences*. 5, 500-516.

- Malaysian Timber Industry Board (2012). *Malaysian Timber Statistics 2009-2011*. [Brochure]. Kuala Lumpur: Percetakan Nasional Malaysia Berhad.
- Malaysian Timber Council (2014). Annual report 2014. *Malaysian Timber Council*. Retrieved September 7, 2015, from <http://mtccomy/wp-content/uploads/2015/11/MTC-Annual-Report-2014pdf>.
- Mandels, M. and Weber, J. (1969). The production of cellulases. *Advances in Chemistry*. 95, 391-414.
- Manfredi, A. P., Perotti, N. I. and Martínez, M. A. (2015). Cellulose degrading bacteria isolated from industrial samples and the gut of native insects from Northwest of Argentina. *Journal of Basic Microbiology*. 55, 1384-1393.
- Marjamaa, K., Toth, K., Bromann, P. A., Szakacs, G. and Kruus, K. (2013). Novel *Penicillium* cellulases for total hydrolysis of lignocellulosics. *Enzyme and Microbial Technology*. 52, 358-369.
- Marlatt, C., Ho, C. T. and Chien, M. (1992). Studies of aroma constituents bound as glycosides in tomato. *Journal of Agricultural and Food Chemistry*. 40, 249-252.
- Martin, M. M. and Martin, J. S. (1978). Cellulose digestion in the midgut of the fungus-growing termite *Macrotermes natalensis*: The role of acquired digestive enzymes. *Science*. 199, 1453-1455.
- Martins, L. G. O., Soares, C. M., Pereira, M. M., Teixeira, M., Costa, T., Jones, G. H. and Henriques, A. O. (2002). Molecular and biochemical characterization of a highly stable bacterial laccase that occurs as a structural component of the *Bacillus subtilis* endospore coat. *Journal of Biological Chemistry*. 277, 18849-18859.
- Massadeh, M. I., Yusoff, W. M. W., Omar, O. and Kader, J. (2001). Synergism of cellulase enzymes in mixed culture solid substrate fermentation. *Biotechnology Letters*. 23, 1771-1774.
- Matkar, K., Chapla, D., Divecha, J., Nighojkar, A. and Madamwar, D. (2013). Production of cellulase by a newly isolated strain of *Aspergillus sydowii* and its optimization under submerged fermentation. *International Biodeterioration and Biodegradation*. 78, 24-33.

- Mattéotti, C., Bauwens, J., Brasseur, C., Tarayre, C., Thonart, P., Destain, J., Francis, F., Haubruge, E., De Pauw, E. and Portetelle, D. (2012). Identification and characterization of a new xylanase from Gram positive bacteria isolated from termite gut (*Reticulitermes santonensis*). *Protein Expression and Purification*. 83, 117-127.
- Maurya, D. P., Singh, D., Pratap, D. and Maurya, J. P. (2012). Optimization of solid state fermentation conditions for the production of cellulase by *Trichoderma reesei*. *Journal of Environmental Biology*. 33, 5-8.
- Mawadza, C., Hatti-Kaul, R., Zvauya, R. and Mattiasson, B. (2000). Purification and characterization of cellulases produced by two *Bacillus* strains. *Journal of Biotechnology*. 83, 177-187.
- McIntyre, M., Berry, D. and McNeil, B. (2000). Role of proteases in autolysis of *Penicillium chrysogenum* chemostat cultures in response to nutrient depletion. *Applied Microbiology and Biotechnology*. 53, 235-242.
- Medina, J. D. C., Woiciechowski, A., Zandona Filho, A., Nosedá, M. D., Kaur, B. S. and Soccol, C. R. (2015). Lignin preparation from oil palm empty fruit bunches by sequential acid/alkaline treatment—A biorefinery approach. *Bioresource Technology*. 194, 172-178.
- Mekhilef, S., Saidur, R., Safari, A. and Mustaffa, W. (2011). Biomass energy in Malaysia: Current state and prospects. *Renewable and Sustainable Energy Reviews*. 15, 3360-3370.
- Membrillo, I., Sánchez, C., Meneses, M., Favela, E. and Loera, O. (2008). Effect of substrate particle size and additional nitrogen source on production of lignocellulolytic enzymes by *Pleurotus ostreatus* strains. *Bioresource Technology*. 99, 7842-7847.
- Menon, V. and Rao, M. (2012). Trends in bioconversion of lignocellulose: Biofuels, platform chemicals and biorefinery concept. *Process in Energy and Combustion Science*. 38, 522-550.
- Michel, F., Dass, S. B., Grulke, E. and Reddy, C. (1991). Role of manganese peroxidases and lignin peroxidases of *Phanerochaete chrysosporium* in the decolorization of kraft bleach plant effluent. *Applied and Environmental Microbiology*. 57, 2368-2375.

- Michelin, M., Ximenes, E., De Moraes, M. D. L. T. and Ladisch, M. R. (2016). Effect of phenolic compounds from pretreated sugarcane bagasse on cellulolytic and hemicellulolytic activities. *Bioresource Technology*. 199, 275-278.
- Mihajlovski, K. R., Carević, M. B., Dević, M. L., Šiler-Marinković, S., Rajilić-Stojanović, M. D. and Dimitrijević-Branković, S. (2015). Lignocellulosic waste material as substrate for Avicelase production by a new strain of *Paenibacillus chitinolyticus* CKS1. *International Biodeterioration and Biodegradation*. 104, 426-434.
- Mikiashvili, N. A., Isikhuemhen, O. S. and Ohimain, E. I. (2011). Lignin degradation, ligninolytic enzymes activities and exopolysaccharide production by *Grifola frondosa* strains cultivated on oak sawdust. *Brazilian Journal of Microbiology*. 42, 1101-1108.
- Miller, G. L. (1959). Use of dinitrosalicylic acid reagent for determination of reducing sugar. *Analytical Chemistry*. 31, 426-428.
- Min, D. Y., Xu, R. S., Hou, Z., Lv, J. Q., Huang, C. X., Jin, Y. C. and Yong, Q. (2015). Minimizing inhibitors during pretreatment while maximizing sugar production in enzymatic hydrolysis through a two-stage hydrothermal pretreatment. *Cellulose*. 22, 1253-1261.
- Miron, J., Ben-Ghedalia, D. and Morrison, M. (2001). Invited review: Adhesion mechanisms of rumen cellulolytic bacteria. *Journal of Dairy Science*. 84, 1294-1309.
- Mishra, D., Kumar, A., Prajapati, C., Singh, A. and Sharma, S. (2013). Identification of compatible bacterial and fungal isolate and their effectiveness against plant disease. *Journal of Environmental Biology*. 34, 183.
- Miskam, A., Zainal, Z. and Yusof, I. (2009). Characterization of sawdust residues for cyclone gasifier. *Journal of Applied Sciences*. 9, 2294-2300.
- Miyazaki, K. (2005). A hyperthermophilic laccase from *Thermus thermophilus* HB27. *Extremophiles*. 9, 415-425.
- Mohanram, S., Rajan, K., Carrier, D. J., Nain, L. and Arora, A. (2015). Insights into biological delignification of rice straw by *Trametes hirsuta* and *Myrothecium roridum* and comparison of saccharification yields with dilute acid pretreatment. *Biomass and Bioenergy*. 76, 54-60.

- Mohtar, S., Busu, T. T. M., Noor, A. M., Shaari, N., Yusoff, N., Bustam, M., Mutalib, M. A. and Mat, H. (2015). Extraction and characterization of lignin from oil palm biomass via ionic liquid dissolution and non-toxic aluminium potassium sulfate dodecahydrate precipitation processes. *Bioresource Technology*. 192, 212-218.
- Mood, S. H., Golfeshan, A. H., Tabatabaei, M., Jouzani, G. S., Najafi, G. H., Gholami, M. and Ardjmand, M. (2013). Lignocellulosic biomass to bioethanol, a comprehensive review with a focus on pretreatment. *Renewable and Sustainable Energy Reviews*. 27, 77-93.
- Moreira, L. (2008). An overview of mannan structure and mannan-degrading enzyme systems. *Applied Microbiology and Biotechnology*. 79, 165-178.
- Mussatto, S. I. and Roberto, I. C. (2004). Alternatives for detoxification of diluted-acid lignocellulosic hydrolyzates for use in fermentative processes: A review. *Bioresource Technology*. 93, 1-10.
- Mussatto, S. I. and Teixeira, J. (2010). Lignocellulose as raw material in fermentation processes. *Current Research*. 2, 897-907.
- Nadeem, M. T., Butt, M. S., Anjum, F. M. and Asgher, M. (2009). Improving bread quality by carboxymethyl cellulase application. *International Journal of Agriculture and Biology*. 11, 727-730.
- Nakamiya, K., Ooi, T. and Kinoshita, S. (1997). Non-heme hydroquinone peroxidase from *Azotobacter beijerinckii* HM121. *Journal of Fermentation and Bioengineering*. 84, 14-21.
- Nakamura, Y., Sawada, T., Sungusia, M. G., Kobayashi, F., Kuwahara, M. and Ito, H. (1997). Lignin peroxidase production by *Phanerochaete chrysosporium* immobilized on polyurethane foam. *Journal of Chemical Engineering of Japan*. 30, 1-6.
- Nakashima, K., Watanabe, H., Saitoh, H., Tokuda, G. and Azuma, J. I. (2002). Dual cellulose-digesting system of the wood-feeding termite, *Coptotermes formosanus* Shiraki. *Insect Biochemistry and Molecular Biology*. 32, 777-784.
- Nanda, S., Azargohar, R., Dalai, A. K. and Kozinski, J. A. (2015). An assessment on the sustainability of lignocellulosic biomass for biorefining. *Renewable and Sustainable Energy Reviews*. 50, 925-941.

- Nandal, P., Ravella, S. R. and Kuhad, R. C. (2013). Laccase production by *Corioloropsis caperata* RCK2011: Optimization under solid state fermentation by Taguchi DOE methodology. *Scientific Reports*. 3, 1-7.
- Nascimento, R., Coelho, R., Marques, S., Alves, L., Girio, F., Bon, E. and Amaral-Collaco, M. (2002). Production and partial characterisation of xylanase from *Streptomyces* sp. strain AMT-3 isolated from Brazilian cerrado soil. *Enzyme and Microbial Technology*. 31, 549-555.
- Nathan, V. K., Rani, M. E., Rathinasamy, G., Dhiraviam, K. N. and Jayavel, S. (2014). Process optimization and production kinetics for cellulase production by *Trichoderma viride* VKF3. *SpringerPlus*. 3, 1-12.
- Nawel, B., Said, B., Estelle, C., Hakim, H. and Duchiron, F. (2011). Production and partial characterization of xylanase produced by *Jonesia denitrificans* isolated in Algerian soil. *Process Biochemistry*. 46, 519-525.
- Ncube, T., Howard, R. L., Abotsi, E. K., Van Rensburg, E. L. J. and Ncube, I. (2012). Jatropha curcas seed cake as substrate for production of xylanase and cellulase by *Aspergillus niger* FGSCA733 in solid-state fermentation. *Industrial Crops and Products*. 37, 118-123.
- Ni, J. and Tokuda, G. (2013). Lignocellulose-degrading enzymes from termites and their symbiotic microbiota. *Biotechnology Advances*. 31, 838-850.
- Nicholson, W. L., Munakata, N., Horneck, G., Melosh, H. J. and Setlow, P. (2000). Resistance of *Bacillus* endospores to extreme terrestrial and extraterrestrial environments. *Microbiology and Molecular Biology Reviews*. 64, 548-572.
- Nigam, P. S. (2013). Microbial enzymes with special characteristics for biotechnological applications. *Biomolecules*. 3, 597-611.
- Niladevi, K. N. (2007). Ligninolytic Enzymes. In Singh nee', P. and Pandey, A (Eds). *Biotechnology for Agro-Industrial Residues Utilisation*. (pp. 397-414). Berlin: Springer.
- Niladevi, K. N. and Prema, P. (2005). Mangrove actinomycetes as the source of ligninolytic enzymes. *Actinomycetology*. 19, 40-47.
- Niladevi, K. N., Sukumaran, R. K. and Prema, P. (2007). Utilization of rice straw for laccase production by *Streptomyces psammoticus* in solid-state fermentation. *Journal of Industrial Microbiology and Biotechnology*. 34, 665-674.

- Ninawe, S., Kapoor, M. and Kuhad, R. C. (2008). Purification and characterization of extracellular xylanase from *Streptomyces cyaneus* SN32. *Bioresource Technology*. 99, 1252-1258.
- Noratiqah, K., Madihah, M., Aisyah, B., Shaza Eva, M., Suraini, A. and Kamarulzaman, K. (2013). Statistical optimization of enzymatic degradation process for oil palm empty fruit bunch (OPEFB) in rotary drum bioreactor using crude cellulase produced from *Aspergillus niger* EFB1. *Biochemical Engineering Journal*. 75, 8-20.
- Norgren, M. and Edlund, H. (2014). Lignin: Recent advances and emerging applications. *Current Opinion in Colloid and Interface Science*. 19, 409-416.
- Oberoi, H. S., Chavan, Y., Bansal, S. and Dhillon, G. S. (2010). Production of cellulases through solid state fermentation using kinnow pulp as a major substrate. *Food and Bioprocess Technology*. 3, 528-536.
- Ohkuma, M. (2003). Termite symbiotic systems: Efficient bio-recycling of lignocellulose. *Applied Microbiology and Biotechnology*. 61, 1-9.
- Oliveira, P. L. D., Duarte, M. C. T., Ponezi, A. N. and Durrant, L. R. (2009). Purification and partial characterization of manganese peroxidase from *Bacillus pumilus* and *Paenibacillus* sp. *Brazilian Journal of Microbiology*. 40, 818-826.
- Ollikka, P., Alhonmäki, K., Leppänen, V. M., Glumoff, T., Raijola, T. and Suominen, I. (1993). Decolorization of azo, triphenyl methane, heterocyclic, and polymeric dyes by lignin peroxidase isoenzymes from *Phanerochaete chrysosporium*. *Applied and Environmental Microbiology*. 59, 4010-4016.
- Pal, A. and Khanum, F. (2010). Production and extraction optimization of xylanase from *Aspergillus niger* DFR-5 through solid-state-fermentation. *Bioresource Technology*. 101, 7563-7569.
- Pal, A. and Khanum, F. (2011). Identification and optimization of critical medium components using statistical experimental designs for enhanced production of xylanase from *Aspergillus flavus* DFR-6. *Food Technology and Biotechnology*. 49, 228-236.
- Pal, S., Banik, S. P. and Khowala, S. (2013). Mustard stalk and straw: A new source for production of lignocellulolytic enzymes by the fungus *Termitomyces clypeatus* and as a substrate for saccharification. *Industrial Crops and Products*. 41, 283-288.

- Palmqvist, E. and Hahn-Hägerdal, B. (2000). Fermentation of lignocellulosic hydrolysates. II: Inhibitors and mechanisms of inhibition. *Bioresource Technology*. 74, 25-33.
- Panagiotou, G., Kekos, D., Macris, B. J. and Christakopoulos, P. (2003). Production of cellulolytic and xylanolytic enzymes by *Fusarium oxysporum* grown on corn stover in solid state fermentation. *Industrial Crops and Products*. 18, 37-45.
- Pandey, A. (1992). Recent process developments in solid-state fermentation. *Process Biochemistry*. 27, 109-117.
- Pandey, A. (2003). Solid-state fermentation. *Biochemical Engineering Journal*. 13, 81-84.
- Pandey, A., Soccol, C. R., Nigam, P. and Soccol, V. T. (2000). Biotechnological potential of agro-industrial residues. I: Sugarcane bagasse. *Bioresource Technology*. 74, 69-80.
- Pandya, J. J. and Gupte, A. (2012). Production of xylanase under solid-state fermentation by *Aspergillus tubingensis* JP-1 and its application. *Bioprocess and Biosystems Engineering*. 35, 769-779.
- Parajó, J. C., Domínguez, H. and Domínguez, J. (1998). Biotechnological production of xylitol. Part 1: Interest of xylitol and fundamentals of its biosynthesis. *Bioresource Technology*. 65, 191-201.
- Parani, K. and Eyini, M. (2012). Production of ligninolytic enzymes during solid state fermentation of coffee pulp by selected fungi. *Science Research Reporter*. 2, 202-209.
- Pasti, M. B. and Belli, M. L. (1985). Cellulolytic activity of actinomycetes isolated from termites (Termitidae) gut. *FEMS Microbiology Letters*. 26, 107-112.
- Patel, M. and Kumar, A. (2016). Production of renewable diesel through the hydroprocessing of lignocellulosic biomass-derived bio-oil: A review. *Renewable and Sustainable Energy Reviews*. 58, 1293-1307.
- Pearl, I. and Benson, H. (1940). The determination of lignin in sulphide pulping liquor. *Paper Trade Journal*. 111, 35-36.
- Peng, J., Bi, X. T., Lim, C. J., Peng, H., Kim, C. S., Jia, D. and Zuo, H. (2015). Sawdust as an effective binder for making torrefied pellets. *Applied Energy*. 157, 491-498.

- Pensupa, N., Jin, M., Kokolski, M., Archer, D. B. and Du, C. (2013). A solid state fungal fermentation-based strategy for the hydrolysis of wheat straw. *Bioresource Technology*. 149, 261-267.
- Perestelo, F., Falcón, M. A., Pérez, M. L., Roig, E. C. and De La Fuente Martin, G. (1989). Bioalteration of kraft pine lignin by *Bacillus megaterium* isolated from compost piles. *Journal of Fermentation and Bioengineering*. 68, 151-153.
- Péros, J., This, P., Confuron, Y. and Chacon, H. (1996). Comparison by isozyme and RAPD analysis of some isolates of the grapevine dieback fungus, *Eutypa lata*. *American Journal of Enology and Viticulture*. 47, 49-56.
- Philippoussis, A., Diamantopoulou, P., Papadopoulou, K., Lakhtar, H., Roussos, S., Parissopoulos, G. and Papanikolaou, S. (2011). Biomass, laccase and endoglucanase production by *Lentinula edodes* during solid state fermentation of reed grass, bean stalks and wheat straw residues. *World Journal of Microbiology and Biotechnology*. 27, 285-297.
- Plácido, J. and Capareda, S. (2015). Ligninolytic enzymes: A biotechnological alternative for bioethanol production. *Bioresources and Bioprocessing*. 2, 1-12.
- Plassard, C. S., Mousain, D. G. and Salsac, L. E. (1982). Estimation of mycelial growth of basidiomycetes by means of chitin determination. *Phytochemistry*. 21, 345-348.
- Pointing, S. B. (1999). Qualitative Methods for the Determination of Lignocellulolytic Enzyme Production by Tropical Fungi. *Fungal Diversity*. 2, 17-33.
- Poorna, C. A. and Prema, P. (2006). Production and partial characterization of endoxylanase by *Bacillus pumilus* using agro industrial residues. *Biochemical Engineering Journal*. 32, 106-112.
- Poorna, C. A. and Prema, P. (2007). Production of cellulase-free endoxylanase from novel alkalophilic thermotolerant *Bacillus pumilus* by solid-state fermentation and its application in wastepaper recycling. *Bioresource Technology*. 98, 485-490.
- Popham, D. L., Helin, J., Costello, C. E. and Setlow, P. (1996). Analysis of the peptidoglycan structure of *Bacillus subtilis* endospores. *Journal of Bacteriology*. 178, 6451-6458.

- Potumarthi, R., Baadhe, R. R., Nayak, P. and Jetty, A. (2013). Simultaneous pretreatment and saccharification of rice husk by *Phanerochete chrysosporium* for improved production of reducing sugars. *Bioresource Technology*. 128, 113-117.
- Pourramezan, Z., Ghezelbash, G. R., Romani, B., Ziaei, S. and Hedayatkah, A. (2012). Screening and identification of newly isolated cellulose-degrading bacteria from the gut of xylophagous termite *Microcerotermes diversus* (Silvestri). *Microbiology*. 81, 736-742.
- Prasad, D. Y., Heitmann, J. A. and Joyce, T. W. (1993). Enzymatic deinking of colored offset newsprint. *Nordic Pulp and Paper Research Journal*. 8, 284.
- Prasad, P., Tanuja and Bedi, S. (2013). Isolation and characterization of cellulose degrading *Brevibacillus brevis* and optimization of its cellulase activity at different physico-chemical parameters. *Research Journal of Biotechnology*. 8, 66-71.
- Puchart, V. R., Katapodis, P., Biely, P., Kremnický, L. R., Christakopoulos, P., Vrřanská, M., Kekos, D., Macris, B. J. and Bhat, M. K. (1999). Production of xylanases, mannanases and pectinases by the thermophilic fungus *Thermomyces lanuginosus*. *Enzyme and Microbial Technology*. 24, 355-361.
- Qi, B., Chen, X., Shen, F., Su, Y. and Wan, Y. (2009). Optimization of enzymatic hydrolysis of wheat straw pretreated by alkaline peroxide using response surface methodology. *Industrial and Engineering Chemistry Research*. 48, 7346-7353.
- Qin, Y., He, H., Li, N., Ling, M. and Liang, Z. (2010). Isolation and characterization of a thermostable cellulase-producing *Fusarium chlamydosporum*. *World Journal of Microbiology and Biotechnology*. 26, 1991-1997.
- Qiu, H., Li, Y., Ji, G., Zhou, G., Huang, X., Qu, Y. and Gao, P. (2009). Immobilization of lignin peroxidase on nanoporous gold: Enzymatic properties and in situ release of H₂O₂ by co-immobilized glucose oxidase. *Bioresource Technology*. 100, 3837-3842.
- Rafiqul, I. and Sakinah, A. M. (2012). Design of process parameters for the production of xylose from wood sawdust. *Chemical Engineering Research and Design*. 90, 1307-1312.

- Raghavarao, K., Ranganathan, T. and Karanth, N. (2003). Some engineering aspects of solid-state fermentation. *Biochemical Engineering Journal*. 13, 127-135.
- Rahman, N. H. A., Aziz, S. A. and Hassan, M. A. (2013). Production of ligninolytic enzymes by newly isolated bacteria from palm oil plantation soils. *BioResources*. 8, 6136-6150.
- Rahman, S., Choudhury, J. and Ahmad, A. (2006). Production of xylose from oil palm empty fruit bunch fiber using sulfuric acid. *Biochemical Engineering Journal*. 30, 97-103.
- Rahman, S., Choudhury, J., Ahmad, A. and Kamaruddin, A. (2007). Optimization studies on acid hydrolysis of oil palm empty fruit bunch fiber for production of xylose. *Bioresource Technology*. 98, 554-559.
- Rai, P., Majumdar, G., Gupta, S. D. and De, S. (2007). Effect of various pretreatment methods on permeate flux and quality during ultrafiltration of mosambi juice. *Journal of Food Engineering*. 78, 561-568.
- Raimbault, M. (1998). General and microbiological aspects of solid substrate fermentation. *Electronic Journal of Biotechnology*. 1, 26-27.
- Rajan, K. and Carrier, D. J. (2014). Effect of dilute acid pretreatment conditions and washing on the production of inhibitors and on recovery of sugars during wheat straw enzymatic hydrolysis. *Biomass and Bioenergy*. 62, 222-227.
- Ramachandra, M., Crawford, D. L. and Hertel, G. (1988). Characterization of an extracellular lignin peroxidase of the lignocellulolytic actinomycete *Streptomyces viridosporus*. *Applied and Environmental Microbiology*. 54, 3057-3063.
- Ramasamy, G., Ratnasingam, J., Bakar, E. S., Halis, R. and Muttiah, N. (2015). Assessment of environmental emissions from sawmilling activity in Malaysia. *BioResources*. 10, 6643-6662.
- Ramin, M., Alimon, A. and Abdullah, N. (2009). Identification of cellulolytic bacteria isolated from the termite *Coptotermes curvignathus* (Holmgren). *Journal of Rapid Methods and Automation in Microbiology*. 17, 103-116.
- Rana, V., Eckard, A. D., Teller, P. and Ahring, B. K. (2014). On-site enzymes produced from *Trichoderma reesei* RUT-C30 and *Aspergillus saccharolyticus* for hydrolysis of wet exploded corn stover and loblolly pine. *Bioresource Technology*. 154, 282-289.

- Rani, G. B., Chiranjeevi, T., Chandel, A. K., Satish, T., Radhika, K., Narasu, M. L. and Uma, A. (2014). Optimization of selective production media for enhanced production of xylanases in submerged fermentation by *Thielaviopsis basicola* MTCC 1467 using L16 orthogonal array. *Journal of Food Science and Technology*. 51, 2508-2516.
- Rashid, M., Rajoka, M., Siddiqui, K. and Shakoori, A. (1997). Kinetic properties of chemically modified β -glucosidase from *Aspergillus niger* 280. *Pakistan Journal of Zoology*. 29, 354-363.
- Ravella, S. R., Quiñones, T. S., Retter, A., Heiermann, M., Amon, T. and Hobbs, P. J. (2010). Extracellular polysaccharide (EPS) production by a novel strain of yeast-like fungus *Aureobasidium pullulans*. *Carbohydrate Polymers*. 82, 728-732.
- Ravindran, R. and Jaiswal, A. K. (2016). A comprehensive review on pre-treatment strategy for lignocellulosic food industry waste: Challenges and opportunities. *Bioresource Technology*. 199, 92-102.
- Razak, D. L. A., Rashid, N. Y. A., Jamaluddin, A., Sharifudin, S. A. and Long, K. (2015). Enhancement of phenolic acid content and antioxidant activity of rice bran fermented with *Rhizopus oligosporus* and *Monascus purpureus*. *Biocatalysis and Agricultural Biotechnology*. 4, 33-38.
- Rekik, H., Nadia, Z. J., Bejar, W., Kourdali, S., Belhoul, M., Hmidi, M., Benkiar, A., Badis, A., Sallem, N. and Bejar, S. (2015). Characterization of a purified decolorizing detergent-stable peroxidase from *Streptomyces griseosporus* SN9. *International Journal of Biological Macromolecules*. 73, 253-263.
- Robinson, T. and Nigam, P. S. (2008). Remediation of textile dye waste water using a white-rot fungus *Bjerkandera adusta* through solid-state fermentation (SSF). *Applied Biochemistry and Biotechnology*. 151, 618-628.
- Robles-Hernández, L., Gonzales-Franco, A., Crawford, D. L. and Chun, W. W. (2008). Review of environmental organopollutants degradation by white-rot basidiomycete mushrooms. *Tecnociencia Chihuahua*. 2, 32-39.
- Rodrigues, T. H., Pinto, G. A. and Gonçalves, L. R. (2008). Effects of inoculum concentration, temperature and carbon sources on tannase production during solid state fermentation of cashew apple bagasse. *Biotechnology and Bioprocess Engineering*. 13, 571-576.

- Rodriguez, S., Santoro, R., Cameselle, C. and Sanroman, A. (1998). Effect of the different parts of the corn cob employed as a carrier on ligninolytic activity in solid state cultures by *Phanerochaete chrysosporium*. *Bioprocess Engineering*. 18, 251-255.
- Rojas-Rejón, O., Cristiani-Urbina, E., Poggi-Varaldo, H., Ramos-Valdivia, A., Martínez-Jiménez, A. and Ponce-Noyola, T. (2010). Production of cellulase and xylanase by *Cellulomonas flavigena* immobilized in sodium alginate in bubble column reactors. *Journal of Biotechnology*. 150, 358.
- Rosenberg, I. M. (2013). *Protein Analysis and Purification: Benchtop Techniques*. Boston: Birkhauser, Springer Science and Business Media.
- Rouland-Lefèvre, C., Inoue, T. and Johjima, T. (2006). Termitomyces/termite interactions. In König, H. and Varma, A. (Eds). *Intestinal Microorganisms of Termites and Other Invertebrates* (pp. 335-350). Berlin: Springer-Verlag.
- Rubilar, O., Diez, M. C. and Gianfreda, L. (2008). Transformation of chlorinated phenolic compounds by white rot fungi. *Critical Reviews in Environmental Science and Technology*. 38, 227-268.
- Ruqayyah, T. I., Jamal, P., Alam, M. Z. and Mirghani, M. E. S. (2013). Biodegradation potential and ligninolytic enzyme activity of two locally isolated *Panus tigrinus* strains on selected agro-industrial wastes. *Journal of Environmental Management*. 118, 115-121.
- Saha, B. C. (2003). Hemicellulose bioconversion. *Journal of Industrial Microbiology and Biotechnology*. 30, 279-291.
- Saha, B. C., Iten, L. B., Cotta, M. A. and Wu, Y. V. (2005). Dilute acid pretreatment, enzymatic saccharification and fermentation of wheat straw to ethanol. *Process Biochemistry*. 40, 3693-3700.
- Saha, B. C., Qureshi, N., Kennedy, G. J. and Cotta, M. A. (2016). Biological pretreatment of corn stover with white-rot fungus for improved enzymatic hydrolysis. *International Biodeterioration and Biodegradation*. 109, 29-35.
- Saini, R., Saini, J. K., Adsul, M., Patel, A. K., Mathur, A., Tuli, D. and Singhania, R. R. (2015). Enhanced cellulase production by *Penicillium oxalicum* for bio-ethanol application. *Bioresource Technology*. 188, 240-246.

- Sajben-Nagy, E., Manczinger, L., Škrbić, B., Živančev, J., Antić, I., Krisch, J. and Vágvölgyi, C. (2014). Characterization of an extracellular laccase of *Leptosphaerulina chartarum*. *World Journal of Microbiology and Biotechnology*. 30, 2449-2458.
- Salakkam, A. (2012). Bioconversion of Biodiesel By-Products to Value-Added Chemicals. PhD Thesis. Page 37, The University of Manchester, United Kingdom.
- Samiullah, T. R., Bakhsh, A., Rao, A. Q., Naz, M. and Saleem, M. (2009). Isolation, purification and characterization of extracellular β -glucosidase from *Bacillus* sp. *Advances in Environmental Biology*. 3, 269-277.
- Sánchez, C. (2009). Lignocellulosic residues: Biodegradation and bioconversion by fungi. *Biotechnology Advances*. 27, 185-194.
- Saritha, M., Arora, A. and Nain, L. (2012). Pretreatment of paddy straw with *Trametes hirsuta* for improved enzymatic saccharification. *Bioresource Technology*. 104, 459-465.
- Saritha, M., Arora, A., Singh, S. and Nain, L. (2013). *Streptomyces griseorubens* mediated delignification of paddy straw for improved enzymatic saccharification yields. *Bioresource Technology*. 135, 12-17.
- Sarrazfadeh, M. H., Guiraud, J. P., Lagneau, C., Gaven, B., Carron, A. and Navarro, J. M. (2005). Growth, sporulation, δ -endotoxins synthesis, and toxicity during culture of *Bacillus thuringiensis* H14. *Current Microbiology*. 51, 75-81.
- Schäfer, A., Konrad, R., Kuhnigk, T., Kämpfer, P., Hertel, H. and König, H. (1996). Hemicellulose-degrading bacteria and yeasts from the termite gut. *Journal of Applied Bacteriology*. 80, 471-478.
- Schlosser, D., Grey, R. and Fritsche, W. (1997). Patterns of ligninolytic enzymes in *Trametes versicolor*. Distribution of extra-and intracellular enzyme activities during cultivation on glucose, wheat straw and beech wood. *Applied Microbiology and Biotechnology*. 47, 412-418.
- Scholl, A. L., Menegol, D., Pitarelo, A. P., Fontana, R. C., Filho, A. Z., Ramos, L. P., Dillon, A. J. P. and Camassola, M. (2015). Ethanol production from sugars obtained during enzymatic hydrolysis of elephant grass (*Pennisetum purpureum*, Schum.) pretreated by steam explosion. *Bioresource Technology*. 192, 228-237.

- Scotti, C. T., Vergoignan, C., Feron, G. and Durand, A. (2001). Glucosamine measurement as indirect method for biomass estimation of *Cunninghamella elegans* grown in solid state cultivation conditions. *Biochemical Engineering Journal*. 7, 1-5.
- Segel, I. (1993). *Enzyme Kinetics: Behaviour and Analysis of Rapid Equilibrium and Steady-State Enzyme Systems*. New York: John Wiley & Sons.
- Sethi, S., Datta, A., Gupta, B. L. and Gupta, S. (2013). Optimization of cellulase production from bacteria isolated from soil. *ISRN Biotechnology*. 2013, 1-7.
- Shah, A. R. and Madamwar, D. (2005). Xylanase production under solid-state fermentation and its characterization by an isolated strain of *Aspergillus foetidus* in India. *World Journal of Microbiology and Biotechnology*. 21, 233-243.
- Shankar, T., Mariappan, V. and Isaiarasu, L. (2011). Screening cellulolytic bacteria from the mid-gut of the popular composting earthworm, *Eudrilus eugeniae* (Kinberg). *World Journal of Zoology*. 6, 142-148.
- Sharma, B., Agrawal, R., Singhanian, R. R., Satlewal, A., Mathur, A., Tuli, D. and Adsul, M. (2015a). Untreated wheat straw: Potential source for diverse cellulolytic enzyme secretion by *Penicillium janthinellum* EMS-UV-8 mutant. *Bioresource Technology*. 196, 518-524.
- Sharma, R., Rawat, R., Bhogal, R. S. and Oberoi, H. S. (2015b). Multi-component thermostable cellulolytic enzyme production by *Aspergillus niger* HN-1 using pea pod waste: Appraisal of hydrolytic potential with lignocellulosic biomass. *Process Biochemistry*. 50, 696-704.
- Sharma, R. K. and Arora, D. S. (2010). Production of lignocellulolytic enzymes and enhancement of in vitro digestibility during solid state fermentation of wheat straw by *Phlebia floridensis*. *Bioresource Technology*. 101, 9248-9253.
- Shastri, Y., Hansen, A., Rodriguez, L. and Ting, K. (2014). Engineering and Science of Biomass Feedstock Production and Provision. In Chaoui, H. and Eckhoff, S. R. (Eds.) *Biomass Feedstock Storage for Quantity and Quality Preservation*. (pp. 165-194). New York: Springer.
- Shi, J., Sharma-Shivappa, R. R., Chinn, M. and Howell, N. (2009). Effect of microbial pretreatment on enzymatic hydrolysis and fermentation of cotton stalks for ethanol production. *Biomass and Bioenergy*. 33, 88-96.

- Shim, S. S. and Kawamoto, K. (2002). Enzyme production activity of *Phanerochaete chrysosporium* and degradation of pentachlorophenol in a bioreactor. *Water Research*. 36, 4445-4454.
- Shrivastava, B., Thakur, S., Khasa, Y. P., Gupte, A., Puniya, A. K. and Kuhad, R. C. (2011). White-rot fungal conversion of wheat straw to energy rich cattle feed. *Biodegradation*. 22, 823-831.
- Sills, D. L. and Gossett, J. M. (2011). Assessment of commercial hemicellulases for saccharification of alkaline pretreated perennial biomass. *Bioresource Technology*. 102, 1389-1398.
- Silva, L., Terrasan, C. R. F. and Carmona, E. C. (2015). Purification and characterization of xylanases from *Trichoderma inhamatum*. *Electronic Journal of Biotechnology*. 18, 307-313.
- Sindhu, R., Binod, P. and Pandey, A. (2016). Biological pretreatment of lignocellulosic biomass—An overview. *Bioresource Technology*. 199, 76-82.
- Sindhu, R., Suprabha, N. G. and Shashidhar, S. (2011). Media engineering for the production of cellulase from *Penicillium* species (SBSS 30) under solid state fermentation. *Biotechnology, Bioinformatics and Bioengineering*. 1, 343-349.
- Singh, A., Kuhad, R. C. and Ward, O. P. (2007). Industrial application of microbial cellulases. *Lignocellulose Biotechnology: Future Prospects*. 345-358.
- Singh, D., Zeng, J., Laskar, D. D., Deobald, L., Hiscox, W. C. and Chen, S. (2011). Investigation of wheat straw biodegradation by *Phanerochaete chrysosporium*. *Biomass and Bioenergy*. 35, 1030-1040.
- Singh, P., Suman, A., Tiwari, P., Arya, N., Gaur, A. and Shrivastava, A. (2008). Biological pretreatment of sugarcane trash for its conversion to fermentable sugars. *World Journal of Microbiology and Biotechnology*. 24, 667-673.
- Singh, S., Moholkar, V. S. and Goyal, A. (2013). Isolation, identification, and characterization of a cellulolytic *Bacillus amyloliquefaciens* strain SS35 from rhinoceros dung. *ISRN microbiology*. 2013, 1-7.
- Singhania, R. R. (2011). Production of Cellulolytic Enzymes for the Hydrolysis of Lignocellulosic Biomass. In Pandey, A. (Ed.). *Biofuels: Alternative Feedstocks and Conversion Processes*. (pp. 177-201). USA: Academic Press.
- Singhania, R. R., Patel, A. K., Soccol, C. R. and Pandey, A. (2009). Recent advances in solid-state fermentation. *Biochemical Engineering Journal*. 44, 13-18.

- Singhania, R. R., Patel, A. K., Sukumaran, R. K., Larroche, C. and Pandey, A. (2013). Role and significance of beta-glucosidases in the hydrolysis of cellulose for bioethanol production. *Bioresource Technology*. 127, 500-507.
- Singhania, R. R., Saini, J. K., Saini, R., Adsul, M., Mathur, A., Gupta, R. and Tuli, D. K. (2014). Bioethanol production from wheat straw via enzymatic route employing *Penicillium janthinellum* cellulases. *Bioresource Technology*. 169, 490-495.
- Singhania, R. R., Sukumaran, R. K. and Pandey, A. (2007). Improved cellulase production by *Trichoderma reesei* RUT C30 under SSF through process optimization. *Applied Biochemistry and Biotechnology*. 142, 60-70.
- Singhania, R. R., Sukumaran, R. K., Patel, A. K., Larroche, C. and Pandey, A. (2010). Advancement and comparative profiles in the production technologies using solid-state and submerged fermentation for microbial cellulases. *Enzyme and Microbial Technology*. 46, 541-549.
- Singhania, R. R., Sukumaran, R. K., Rajasree, K. P., Mathew, A., Gottumukkala, L. and Pandey, A. (2011). Properties of a major β -glucosidase-BGL1 from *Aspergillus niger* NII-08121 expressed differentially in response to carbon sources. *Process Biochemistry*. 46, 1521-1524.
- Sinma, K., Khucharoenphaisan, K., Kitpreechavanich, V. and Tokuyama, S. (2011). Purification and characterization of a thermostable xylanase from *Saccharopolyspora pathumthaniensis* S582 isolated from the gut of a termite. *Bioscience, Biotechnology and Biochemistry*. 75, 1957-1963.
- Slaytor, M. (2000). *Energy Metabolism in The Termite and its Gut Microbiota*. Netherlands: Springer.
- Sluiter, J. B., Ruiz, R. O., Scarlata, C. J., Sluiter, A. D. and Templeton, D. W. (2010). Compositional analysis of lignocellulosic feedstocks. Review and description of methods. *Journal of Agricultural and Food Chemistry*. 58, 9043-9053.
- Smitha, R. B., Jisha, V. N., Pradeep, S., Josh, M. S. and Benjamin, S. (2013). Potato flour mediated solid-state fermentation for the enhanced production of *Bacillus thuringiensis*-toxin. *Journal of Bioscience and Bioengineering*. 116, 595-601.
- Sohail, M., Siddiqi, R., Ahmad, A. and Khan, S. A. (2009). Cellulase production from *Aspergillus niger* MS82: Effect of temperature and pH. *New Biotechnology*. 25, 437-441.

- Song, H. Y., Lim, H. K., Lee, K. I. and Hwang, I. T. (2014). A new bi-modular endo- β -1, 4-xylanase KRICT PX-3 from whole genome sequence of *Paenibacillus terrae* HPL-003. *Enzyme and Microbial Technology*. 54, 1-7.
- Song, J. M. and Wei, D. Z. (2010). Production and characterization of cellulases and xylanases of *Cellulosimicrobium cellulans* grown in pretreated and extracted bagasse and minimal nutrient medium M9. *Biomass and Bioenergy*. 34, 1930-1934.
- Song, L., Yu, H., Ma, F. and Zhang, X. (2013). Biological pretreatment under non-sterile conditions for enzymatic hydrolysis of corn stover. *BioResources*. 8, 3802-3816.
- Srebotnik, E. and Hammel, K. E. (2000). Degradation of nonphenolic lignin by the laccase/1-hydroxybenzotriazole system. *Journal of Biotechnology*. 81, 179-188.
- Sreenath, H. K. and Santhanam, K. (1992). The use of commercial enzymes in white grape juice clarification. *Journal of Fermentation and Bioengineering*. 73, 241-243.
- Sreenath, H. K., Shah, A. B., Yang, V. W., Gharia, M. M. and Jeffries, T. W. (1996). Enzymatic polishing of jute/cotton blended fabrics. *Journal of Fermentation and Bioengineering*. 81, 18-20.
- Sridevi, A., Narasimha, G., Ramanjaneyulu, G., Dileepkumar, K., Reddy, B. R. and Devi, P. S. (2015). Saccharification of pretreated sawdust by *Aspergillus niger* cellulase. *3 Biotech*. 5, 883-892.
- Srinivasan, M. and Rele, M. V. (1999). Microbial xylanases for paper industry. *Current Science*. 77, 137-142.
- Srivastava, N., Singh, J., Ramteke, P. W., Mishra, P. and Srivastava, M. (2015). Improved production of reducing sugars from rice straw using crude cellulase activated with Fe₃O₄/Alginate nanocomposite. *Bioresource Technology*. 183, 262-266.
- Subramaniyan, S. and Prema, P. (2002). Biotechnology of microbial xylanases: Enzymology, molecular biology and application. *Critical Reviews in Biotechnology*. 22, 33-64.
- Subramaniyan, S., Sandhia, G. and Prema, P. (2001). Control of xylanase production without protease activity in *Bacillus* sp. by selection of nitrogen source. *Biotechnology Letters*. 23, 369-371.

- Sudo, S., Kobayashi, S., Kaneko, A., Sato, K. and Oba, T. (1995) Growth of submerged mycelia of *Aspergillus kawachii* in solid-state culture. *Journal of Fermentation and Bioengineering*. 79, 252-256.
- Suhara, H., Kodama, S., Kamei, I., Maekawa, N. and Meguro, S. (2012). Screening of selective lignin-degrading basidiomycetes and biological pretreatment for enzymatic hydrolysis of bamboo culms. *International Biodeterioration and Biodegradation*. 75, 176-180.
- Sukumaran, R. K., Singhanian, R. R., Mathew, G. M. and Pandey, A. (2009). Cellulase production using biomass feed stock and its application in lignocellulose saccharification for bio-ethanol production. *Renewable Energy*. 34, 421-424.
- Sukumaran, R. K., Singhanian, R. R. and Pandey, A. (2005). Microbial cellulases-production, applications and challenges. *Journal of Scientific and Industrial Research*. 64, 832.
- Sun, Y. and Cheng, J. (2002). Hydrolysis of lignocellulosic materials for ethanol production: A review. *Bioresource Technology*. 83, 1-11.
- Taherzadeh, M. J. and Karimi, K. (2007). Enzymatic-based hydrolysis processes for ethanol from lignocellulosic materials: A review. *BioResources*. 2, 707-738.
- Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M. and Kumar, S. (2011). MEGA5: Molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Molecular Biology and Evolution*. 28, 2731-2739.
- Tarayre, C., Brognaux, A., Bauwens, J., Brasseur, C., Mattéotti, C., Millet, C., Destain, J., Vandebol, M., Portetelle, D. and De Pauw, E. (2014). Isolation of amylolytic, xylanolytic and cellulolytic microorganisms extracted from the gut of the termite *Reticulitermes santonensis* by means of a micro-aerobic atmosphere. *World Journal of Microbiology and Biotechnology*. 30, 1655-1660.
- Tartar, A., Wheeler, M. M., Zhou, X., Coy, M. R., Boucias, D. G. and Scharf, M. E. (2009). Parallel metatranscriptome analyses of host and symbiont gene expression in the gut of the termite *Reticulitermes flavipes*. *Biotechnology for Biofuels*. 2, 1.

- Tesfaw, A. and Assefa, F. (2014). Co-culture: A great promising method in single cell protein production. *Biotechnology and Molecular Biology Reviews*. 9, 12-20.
- Thomas, L., Larroche, C. and Pandey, A. (2013). Current developments in solid-state fermentation. *Biochemical Engineering Journal*. 81, 146-161.
- Thomas, L., Ram, H., Kumar, A. and Singh, V. P. (2016). Production, optimization and characterization of organic solvent tolerant cellulases from a lignocellulosic waste-degrading actinobacterium, *Promicromonospora* sp. VP111. *Applied Biochemistry and Biotechnology*. 1-17.
- Tien, M. and Kirk, T. K. (1984). Lignin-degrading enzyme from *Phanerochaete chrysosporium*: Purification, characterization, and catalytic properties of a unique H₂O₂-requiring oxygenase. *Proceedings of the National Academy of Sciences*. 81, 2280-2284.
- Tiwari, R., Pranaw, K., Singh, S., Nain, P. K., Shukla, P. and Nain, L. (2015). Two-step statistical optimization for cold active β -glucosidase production from *Pseudomonas lutea* BG8 and its application for improving saccharification of paddy straw. *Biotechnology and Applied Biochemistry*. DOI:10.1002/bab.1415.
- Tokuda, G., Watanabe, H., Matsumoto, T. and Noda, H. (1997). Cellulose digestion in the wood-eating higher termite, *Nasutitermes takasagoensis* (Shiraki): Distribution of cellulases and properties of endo- β -1, 4-gluconase. *Zoological Science*. 14, 83-93.
- Tortora, G. J., Funke, B. R. and Case, C. L. (2004). *Microbiology An Introduction*. (8th ed.). San Francisco: Pearson Benjamin Cummings.
- Trevorah, R. M. and Othman, M. Z. (2015). Alkali pretreatment and enzymatic hydrolysis of Australian timber mill sawdust for biofuel production. *Journal of Renewable Energy*. 2015. 1-9.
- Trivedi, N., Gupta, V., Kumar, M., Kumari, P., Reddy, C. and Jha, B. (2011). An alkali-halotolerant cellulase from *Bacillus flexus* isolated from green seaweed *Ulva lactuca*. *Carbohydrate Polymers*. 83, 891-897.
- Tuisel, H., Sinclair, R., Bumpus, J. A., Ashbaugh, W., Brock, B. J. and Aust, S. D. (1990). Lignin peroxidase H₂ from *Phanerochaete chrysosporium*: Purification, characterization and stability to temperature and pH. *Archives of Biochemistry and Biophysics*. 279, 158-166.

- Tuncer, M., Kuru, A., Sahin, N., Isikli, M. and Isik, K. (2009). Production and partial characterization of extracellular peroxidase produced by *Streptomyces* sp. F6616 isolated in Turkey. *Annals of Microbiology*. 59, 323-334.
- Tye, Y. Y., Lee, K. T., Abdullah, W. N. W. and Leh, C. P. (2011). Second-generation bioethanol as a sustainable energy source in Malaysia transportation sector: Status, potential and future prospects. *Renewable and Sustainable Energy Reviews*. 15, 4521-4536.
- Tymchyshyn, M. and Xu, C. C. (2010). Liquefaction of bio-mass in hot-compressed water for the production of phenolic compounds. *Bioresource Technology*. 101, 2483-2490.
- Umikalsom, M., Ariff, A., Zulkifli, H., Tong, C., Hassan, M. and Karim, M. (1997). The treatment of oil palm empty fruit bunch fibre for subsequent use as substrate for cellulase production by *Chaetomium globosum* Kunze. *Bioresource Technology*. 62, 1-9.
- Umsza-Guez, M. A., Díaz, A. B., Ory, I. D., Blandino, A., Gomes, E. and Caro, I. (2011). Xylanase production by *Aspergillus awamori* under solid state fermentation conditions on tomato pomace. *Brazilian Journal of Microbiology*. 42, 1585-1597.
- Upadhyaya, S. K., Manandhar, A., Mainali, H., Pokhrel, A. R., Rijal, A., Pradhan, B. and Koirala, B. (2012). Isolation and Characterization of Cellulolytic Bacteria from Gut of Termite. *Rentech Symposium Compendium*, March 2012. 14-18.
- Ürek, R. Ö. and Pazarlioğlu, N. K. (2005). Production and stimulation of manganese peroxidase by immobilized *Phanerochaete chrysosporium*. *Process Biochemistry*. 40, 83-87.
- Valášková, V. and Baldrian, P. (2006a). Degradation of cellulose and hemicelluloses by the brown rot fungus *Piptoporus betulinus*—production of extracellular enzymes and characterization of the major cellulases. *Microbiology*. 152, 3613-3622.
- Valášková, V. and Baldrian, P. (2006b). Estimation of bound and free fractions of lignocellulose-degrading enzymes of wood-rotting fungi *Pleurotus ostreatus*, *Trametes versicolor* and *Piptoporus betulinus*. *Research in Microbiology*. 157, 119-124.

- Van Dyk, J. and Pletschke, B. (2012). A review of lignocellulose bioconversion using enzymatic hydrolysis and synergistic cooperation between enzymes—factors affecting enzymes, conversion and synergy. *Biotechnology Advances*. 30, 1458-1480.
- Van Dyk, J. S., Sakka, M., Sakka, K. and Pletschke, B. I. (2010). Identification of endoglucanases, xylanases, pectinases and mannanases in the multi-enzyme complex of *Bacillus licheniformis* SVD1. *Enzyme and Microbial Technology*. 47, 112-118.
- Veerabhadrapa, M. B., Shivakumar, S. B. and Devappa, S. (2014). Solid-state fermentation of *Jatropha* seed cake for optimization of lipase, protease and detoxification of anti-nutrients in *Jatropha* seed cake using *Aspergillus versicolor* CJS-98. *Journal of Bioscience and Bioengineering*. 117, 208-214.
- Velu, C. (2011). Insilico screening and comparative study on the effectiveness of textile dye decolourization by crude laccase immobilised alginate encapsulated beads from *Pleurotus ostreatus*. *Journal of Bioprocessing and Biotechnique*. 1, 1-6.
- Verardi, A., Ricca, E., De Bari, I. and Calabrò, V. (2012). Hydrolysis of lignocellulosic biomass: Current status of processes and technologies and future perspectives. *INTECH Open Access Publisher*. 1-29.
- Verma, D. and Satyanarayana, T. (2012). Cloning, expression and applicability of thermo-alkali-stable xylanase of *Geobacillus thermoleovorans* in generating xylooligosaccharides from agro-residues. *Bioresource Technology*. 107, 333-338.
- Vijayaraghavan, P., Vincent, S. P. and Dhillon, G. (2016). Solid-substrate bioprocessing of cow dung for the production of carboxymethyl cellulase by *Bacillus halodurans* IND18. *Waste Management*. 48, 513-520.
- Vilanova, C., Marco, G., Domínguez-Escribà, L., Genovés, S., Sentandreu, V., Bataller, E., Ramón, D. and Porcar, M. (2012). Bacteria from acidic to strongly alkaline insect midguts: Potential sources of extreme cellulolytic enzymes. *Biomass and Bioenergy*. 45, 288-294.
- Viniegra-González, G., Favela-Torres, E., Aguilar, C. N., De Jesus Romero-Gomez, S., Diaz-Godínez, G. and Augur, C. (2003). Advantages of fungal enzyme production in solid state over liquid fermentation systems. *Biochemical Engineering Journal*. 13, 157-167.

- Virupakshi, S., Babu, K. G., Gaikwad, S. R. and Naik, G. (2005). Production of a xylanolytic enzyme by a thermoalkaliphilic *Bacillus* sp. JB-99 in solid state fermentation. *Process Biochemistry*. 40, 431-435.
- Vivekanand, V., Dwivedi, P., Pareek, N. and Singh, R. P. (2011). Banana peel: A potential substrate for laccase production by *Aspergillus fumigatus* VkJ2. 4.5 in solid-state fermentation. *Applied Biochemistry and Biotechnology*. 165, 204-220.
- Waeonukul, R., Kosugi, A., Prawitwong, P., Deng, L., Tachaapaikoon, C., Pason, P., Ratanakhanokchai, K., Saito, M. and Mori, Y. (2013). Novel cellulase recycling method using a combination of *Clostridium thermocellum* cellulosomes and *Thermoanaerobacter brockii* β -glucosidase. *Bioresource Technology*. 130, 424-430.
- Walia, A., Mehta, P., Chauhan, A. and Shirkot, C. K. (2013). Optimization of cellulase-free xylanase production by alkalophilic *Cellulosimicrobium* sp. CKMX1 in solid-state fermentation of apple pomace using central composite design and response surface methodology. *Annals of Microbiology*. 63, 187-198.
- Wan, C. and Li, Y. (2011). Effectiveness of microbial pretreatment by *Ceriporiopsis subvermispora* on different biomass feedstocks. *Bioresource Technology*. 102, 7507-7512.
- Wan, C. and Li, Y. (2012). Fungal pretreatment of lignocellulosic biomass. *Biotechnology Advances*. 30, 1447-1457.
- Wang, H. C., Chen, Y. C., Huang, C. T. and Hseu, R. S. (2013a). Cloning and characterization of a thermostable and pH-stable cellobiohydrolase from *Neocallimastix patriciarum* J11. *Protein Expression and Purification*. 90, 153-159.
- Wang, H., Peng, L., Ding, Z., Wu, J. and Shi, G. (2015a). Stimulated laccase production of *Pleurotus ferulae* JM301 fungus by *Rhodotorula mucilaginosa* yeast in co-culture. *Process Biochemistry*. 50, 901-905.
- Wang, J., Li, J., Liu, J., Hua, B., Wang, X., Lv, Y., Cao, Y. and Cui, Z. (2014). Influence of cell disruption and elution on cellulase release of *Clostridium straminisolvens* (CSK1). *Applied Biochemistry and Biotechnology*. 173, 510-521.

- Wang, X. J., Bai, J. G. and Liang, Y. X. (2006). Optimization of multienzyme production by two mixed strains in solid-state fermentation. *Applied Microbiology and Biotechnology*. 73, 533-540.
- Wang, Y., Feng, S., Zhan, T., Huang, Z., Wu, G. and Liu, Z. (2013b). Improving catalytic efficiency of endo- β -1, 4-xylanase from *Geobacillus stearothermophilus* by directed evolution and H179 saturation mutagenesis. *Journal of Biotechnology*. 168, 341-347.
- Wang, Y., Vazquez-Duhalt, R. and Pickard, M. A. (2001). Effect of growth conditions on the production of manganese peroxidase by three strains of *Bjerkandera adusta*. *Canadian Journal of Microbiology*. 47, 277-282.
- Wang, Z., Guo, H., Shen, F., Yang, G., Zhang, Y., Zeng, Y., Wang, L., Xiao, H. and Deng, S. (2015b). Biochar produced from oak sawdust by Lanthanum (La)-involved pyrolysis for adsorption of ammonium (NH^{+4}), nitrate (NO^{-3}), and phosphate (PO^{-34}). *Chemosphere*. 119, 646-653.
- Wati, L., Kumari, S. and Kundu, B. (2007). Paddy straw as substrate for ethanol production. *Indian Journal of Microbiology*. 47, 26-29.
- Wei, K. S. C., Teoh, T. C., Koshy, P., Salmah, I. and Zainudin, A. (2015). Cloning, expression and characterization of the endoglucanase gene from *Bacillus subtilis* UMC7 isolated from the gut of the indigenous termite *Macrotermes malaccensis* in *Escherichia coli*. *Electronic Journal of Biotechnology*. 18, 103-109.
- Weisburg, W. G., Barns, S. M., Pelletier, D. A. and Lane, D. J. (1991). 16S ribosomal DNA amplification for phylogenetic study. *Journal of Bacteriology*. 173, 697-703.
- Wen, X., Jia, Y. and Li, J. (2009). Degradation of tetracycline and oxytetracycline by crude lignin peroxidase prepared from *Phanerochaete chrysosporium*—a white rot fungus. *Chemosphere*. 75, 1003-1007.
- Wen, Z., Liao, W. and Chen, S. (2005). Production of cellulase/ β -glucosidase by the mixed fungi culture *Trichoderma reesei* and *Aspergillus phoenicis* on dairy manure. *Process Biochemistry*. 40, 3087-3094.
- Wenzel, M., Schöning, I., Berchtold, M., Kämpfer, P. and König, H. (2002). Aerobic and facultatively anaerobic cellulolytic bacteria from the gut of the termite *Zootermopsis angusticollis*. *Journal of Applied Microbiology*. 92, 32-40.

- Wesenberg, D., Kyriakides, I. and Agathos, S. N. (2003). White-rot fungi and their enzymes for the treatment of industrial dye effluents. *Biotechnology Advances*. 22, 161-187.
- Wilson, D. B. (2009). Cellulases and biofuels. *Current Opinion in Biotechnology*. 20, 295-299.
- Wong, D. W. (2009). Structure and action mechanism of ligninolytic enzymes. *Applied Biochemistry and Biotechnology*. 157, 174-209.
- Wong, W. Y. (2013). Isolation, Identification and Characterization of 2,2-dichloropropionate Utilizing Bacteria. MSc Thesis. Page 60, Universiti Teknologi Malaysia, Skudai.
- Woo, H. L., Hazen, T. C., Simmons, B. A. and DeAngelis, K. M. (2014). Enzyme activities of aerobic lignocellulolytic bacteria isolated from wet tropical forest soils. *Systematic and Applied Microbiology*. 37, 60-67.
- Workman, M., Andersen, M. R. and Thykaer, J. (2013). Integrated approaches for assessment of cellular performance in industrially relevant filamentous fungi. *Industrial Biotechnology*. 9, 337-344.
- Xiong, H., Von Weymarn, N., Leisola, M. and Turunen, O. (2004). Influence of pH on the production of xylanases by *Trichoderma reesei* Rut C-30. *Process Biochemistry*. 39, 731-736.
- Xu, G., Wang, L., Liu, J. and Wu, J. (2013). FTIR and XPS analysis of the changes in bamboo chemical structure decayed by white-rot and brown-rot fungi. *Applied Surface Science*. 280, 799-805.
- Yadav, S. and Chandra, R. (2015). Syntrophic co-culture of *Bacillus subtilis* and *Klebsiella pneumoniae* for degradation of kraft lignin discharged from rayon grade pulp industry. *Journal of Environmental Sciences*. 33, 229-238.
- Yaman, S. (2004). Pyrolysis of biomass to produce fuels and chemical feedstocks. *Energy Conversion and Management*. 45, 651-671.
- Yan, Y., Xu, J., Li, T. and Ren, Z. (1999). Liquefaction of sawdust for liquid fuel. *Fuel Processing Technology*. 60, 135-143.
- Yano, J. K. and Poulos, T. L. (2003). New understandings of thermostable and peizostable enzymes. *Current Opinion in Biotechnology*. 14, 360-365.
- Yanti, Y. (2015). Peroxidase enzyme activity of *Rhizobacteria*-introduced shallots bulbs to induce resistance of shallot towards bacterial leaf blight (*Xanthomonas axonopodis* pv *Allii*). *Procedia Chemistry*. 14, 501-507.

- Yasmeen, Q., Asgher, M., Sheikh, M. A. and Nawaz, H. (2013). Optimization of ligninolytic enzymes production through response surface methodology. *BioResources*. 8, 944-968.
- Yoon, L. W., Ang, T. N., Ngoh, G. C. and Chua, A. S. M. (2014). Fungal solid-state fermentation and various methods of enhancement in cellulase production. *Biomass and Bioenergy*. 67, 319-338.
- Yoon, L. W., Ngoh, G. C. and Chua, A. S. M. (2013). Simultaneous production of cellulase and reducing sugar through modification of compositional and structural characteristic of sugarcane bagasse. *Enzyme and Microbial Technology*. 53, 250-256.
- Yu, H., Guo, G., Zhang, X., Yan, K. and Xu, C. (2009). The effect of biological pretreatment with the selective white-rot fungus *Echinodontium taxodii* on enzymatic hydrolysis of softwoods and hardwoods. *Bioresource Technology*. 100, 5170-5175.
- Zadrazil, F. and Puniya, A. K. (1995). Studies on the effect of particle size on solid-state fermentation of sugarcane bagasse into animal feed using white-rot fungi. *Bioresource Technology*. 54, 85-87.
- Zenevich, M. C. P., Jacques, A., Furigo, A. F., Oliveira, J. V. and De Oliveira, D. (2016). Enzymatic hydrolysis of soybean and waste cooking oils under ultrasound system. *Industrial Crops and Products*. 80, 235-241.
- Zeng, G. M., Zhao, M. H., Huang, D. L., Lai, C., Huang, C., Wei, Z., Xu, P., Li, N. J., Zhang, C., Li, F. L. and Cheng, M. (2013). Purification and biochemical characterization of two extracellular peroxidases from *Phanerochaete chrysosporium* responsible for lignin biodegradation. *International Biodeterioration and Biodegradation*. 85, 166-172.
- Zhang, D., Ong, Y. L., Li, Z. and Wu, J. C. (2012a). Optimization of dilute acid-catalyzed hydrolysis of oil palm empty fruit bunch for high yield production of xylose. *Chemical Engineering Journal*. 181, 636-642.
- Zhang, H. and Sang, Q. (2015). Production and extraction optimization of xylanase and β -mannanase by *Penicillium chrysogenum* QML-2 and primary application in saccharification of corn cob. *Biochemical Engineering Journal*. 97, 101-110.

- Zhang, L., Liu, Y., Niu, X., Liu, Y. and Liao, W. (2012b). Effects of acid and alkali treated lignocellulosic materials on cellulase/xylanase production by *Trichoderma reesei* Rut C-30 and corresponding enzymatic hydrolysis. *Biomass and Bioenergy*. 37, 16-24.
- Zhang, Q., Zhang, W., Lin, C., Xu, X. and Shen, Z. (2012c). Expression of an *Acidothermus cellulolyticus* endoglucanase in transgenic rice seeds. *Protein Expression and Purification*. 82, 279-283.
- Zhang, S., Jiang, M., Zhou, Z., Zhao, M. and Li, Y. (2012d). Selective removal of lignin in steam-exploded rice straw by *Phanerochaete chrysosporium*. *International Biodeterioration and Biodegradation*. 75, 89-95.
- Zhang, X., Yu, H., Huang, H. and Liu, Y. (2007). Evaluation of biological pretreatment with white rot fungi for the enzymatic hydrolysis of bamboo culms. *International Biodeterioration and Biodegradation*. 60, 159-164.
- Zhang, Y. H. P. and Lynd, L. R. (2004). Toward an aggregated understanding of enzymatic hydrolysis of cellulose: Noncomplexed cellulase systems. *Biotechnology and Bioengineering*. 88, 797-824.
- Zheng, Y., Zhao, J., Xu, F. and Li, Y. (2014). Pretreatment of lignocellulosic biomass for enhanced biogas production. *Progress in Energy and Combustion Science*. 42, 35-53.
- Zheng, Z. and Shetty, K. (1998). Solid-state production of beneficial fungi on apple processing wastes using glucosamine as the indicator of growth. *Journal of Agricultural and Food Chemistry*. 46, 783-787.
- Zhou, Y., Selvam, A. and Wong, J. W. (2014). Evaluation of humic substances during co-composting of food waste, sawdust and Chinese medicinal herbal residues. *Bioresource Technology*. 168, 229-234.
- Zhu, Y., Zhang, H., Cao, M., Wei, Z., Huang, F. and Gao, P. (2011). Production of a thermostable metal-tolerant laccase from *Trametes versicolor* and its application in dye decolorization. *Biotechnology and Bioprocess Engineering*. 16, 1027-1035.
- Zhuang, J., Marchant, M., Nokes, S. and Strobel, H. (2007). Economic analysis of cellulase production methods for bio-ethanol. *Applied Engineering in Agriculture*. 23, 679-687.