ASSESSMENT OF DIATOM COMPOSITION AT SEVERAL LOCATIONS ALONG TEBRAU STRAITS FOR ASSIGNING SCENE OF DROWNING

NURUL AMIRA BINTI MOHAMED FARID

UNIVERSITI TEKNOLOGI MALAYSIA

ASSESSMENT OF DIATOM COMPOSITION AT SEVERAL LOCATIONS ALONG TEBRAU STRAITS FOR ASSIGNING SCENE OF DROWNING

NURUL AMIRA BINTI MOHAMED FARID

A dissertation submitted in partial fulfillment of the requirements for the award of degree of Master of Science

> Faculty of Science Universiti Teknologi Malaysia

> > FEBRUARY 2017

ACKNOWLEDGEMENT

In the name of Allah, the Most Gracious and the Most Merciful. I would like to express my gratitude to Allah S. W. T for the strength and His blessings for me in completing this dissertation with success.

I would like to take this chance to express my profound gratitude and deep regards toward my supervisor Dr Hasmerya Maarof ; my co-supervisor Dr Aznool Haidy and to my forensic lecturers Dr Naji A. Mahat and Assoc. Prof. Dr. Umi Kalthom for exemplary guidance, monitoring and constant encouragement during the completion of this dissertation.

I also take this opportunity to thank members of Forensic Lab, Ms Nurul Huda, Mr Mokhtar, Mr Khaidir and my classmates for all support, valuable information and guidance which helped me throughout my studies.

Finally, I want to thank my beloved family members especially my parents for constant encouragement and eternal love during the project, my friend for support and Faculty of Science at Universiti Teknologi Malaysia for the knowledge and experience given to me.

ABSTRACT

The use of diatoms for investigation of drowning death cases has been suggested in the literature. This study designates the assemblages of diatoms at different stations along the Tebrau Straits to establish diatomological mapping as a forensic measure to determine the site of drowning based on association of diatom assemblages recovered from the organ samples with that of the Straits. In each station, water samples from 3 substations (200 m from the shore), separated by a distance of 500 m apart were collected at 2 m depth using a Horizontal Van Dorn sampler. Replicating the same procedure, water samples from 5 different stations distanced by about 20 km from one another, along the Johor Straits were sampled during March-April 2015. To study the recoverability of diatoms from 13 real forensic cases, ranging between 9 months – 4 years, samples of organs (livers, lungs and kidneys) supplied by the pathologist were analyzed. The samples were processed using acid digestion technique and taxonomic identification of diatoms with their percentage assemblages were analyzed under light microscope. A total of 26 genera found from seawater samples were identified. The percentages assemblage of diatoms varied among the different sampling stations and hence, may be useful for distinguishing the stations. While diatoms remained recoverable in lungs and kidneys in majority of suspected drowning cases, the same was not observed in livers. The fact that diatoms were not observed in 3 later (11-12 months) cases, inferring the cause of death due to drowning may prove inconclusive.

ABSTRAK

Penggunaan diatom untuk penyiasatan kes mati lemas telah dicadangkan dalam ulasan tulisan. Kajian ini meninjau himpunan diatom pada stesen yang berbeza sepanjang Selat Tebrau untuk menghasilkan pemetaan diatomologikal sebagai satu langkah forensik bagi menentukan tapak mati lemas berdasarkan perkaitan diatom yang ditemui daripada sampel organ dan Selat Tebrau. Dalam setiap stesen, sampel air daripada 3 substesen (200 m dari pesisiran pantai), berasingan jarak 500 m telah dikumpul pada kedalaman 2 m menggunakan 'Horizontal Van Dorn sampler'. Sampel air daripada 5 stesen berbeza dijarakkan oleh 20 km dari satu sama lain, sepanjang Selat Tebrau telah disampel pada March-April 2015 dengan mengulangi langkah yang sama. Untuk mengkaji kebolehpulihan diatom dari 13 kes forensik sebenar, berjulat antara 9 bulan sehingga 4 tahun, sampel organ (hati, paru-paru dan buah pinggang) diperoleh daripada ahli patologi telah dianalisis. Sampel-sampel berikut diproses menggunakan teknik cernaan asid dan pengenalan taksonomi diatom serta peratusan himpunan diatom dianalisis menggunakan mikroskop cahaya. Sejumlah 26 genera daripada sampel air laut telah dikenalpasti. Peratus himpunan diatom yang berbeza di setiap stesen persampelan, membolehkan perbezaan stesen sepanjang Selat Tebrau dilakukan. Diatom yang terdapat dalam paru-paru dan buah pinggang dalam majoriti kes yang disyaki mati lemas tetap dibaikpulih, walaubagaimanapun, perkara yang sama tidak dapat diperhatikan pada diatom yang terdapat dalam hati. Diatom tidak diperhatikan dalam 3 kes (11-12 bulan) kemudian, membuktikan punca kematian adalah disebabkan oleh mati lemas mungkin tidak tepat.

TABLE OF CONTENTS

CHAPTER	TITLE	PAGE
	DECLARATION	ii
	ACKNOWLEDGEMENT	iii
	ABSTRACT	iv
	ABSTRAK	V
	TABLE OF CONTENTS	vi
	LIST OF TABLES	ix
	LIST OF FIGURES	Х
	LIST OF ABBREVIATIONS AND SYMBOLS	xi
1	INTRODUCTION	1
	1.1 Background of Study	1
	1.2 Problem Statement	2
	1.3 Objectives	3
	1.4 Scope of Study	3
	1.5 Significance of Study	4
2	LITERATURE REVIEW	5
	2.1 Biological Aspects of Diatom and the Importance of	5
	Diatom	
	2.1.1 Biological Aspects of Diatom	5
	2.1.2 The Importance of Diatom	9
	2.2 Drowning and Its Medico-legal Importance	11
	2.2.1 Pathophysiology of Drowning	11
	2.2.2 Macroscopical Findings of Drowning Victim	12

2.2.3 Postmortem Injuries of Drowning Victim	14
2.2.4 Postmortem Immersion of Drowning Victim	14
2.3 Diatom test	15
2.3.1 Methods of Recovery	15
2.3.1.1 Acid Digestion	16
2.3.1.2 Enzymatic Digestion	18
2.3.1.3 Soluene-350	18
2.3.2 Location of Diatom Recovery	19
2.3.3 Diatom Taphonomy	20
2.4 Method to diagnose drowning other than diatom tests	21
2.4.1 Microbiological tests	21
2.4.1.1 Marine Bacteria	22
2.4.1.2 Bioluminescent Bacteria	22
2.4.1.3 Fecal Bacteria	23
2.4.1.4 Metagenomic Analysis	24
2.4.1.5 Advantages and Disadvantages of	24
Microbiological Test	
2.4.2 Quantification of Drowning Markers	25
2.5 Tebrau Straits	26
METHODOLOGY	27
3.1 Materials	27
3.1.1 Instrument and Apparatus	27
3.1.2 Chemicals	28
3.2 Methods	28
3.2.1 Experimental design	28
3.2.2 Diatom Extraction from seawater and organ samples	31
3.2.3 Microscopic Observation and Taxonomic	32
Identification	

3

4	RESULTS AND DISCUSSION	
	4.1 Diatomological Mapping Along the Tebrau Straits	34
	4.1.1 Lebam River, Kota Tinggi, Johor (Station 1)	35
	4.1.2 Kabong Village, Pasir Gudang, Johor (Station 2)	50
	4.1.3 Johor Causeway, Johor Bharu, Johor (Station 3)	53
	4.1.4 Pendas, Gelang Patah, Johor (Station 4)	55
	4.1.5 Tanjung Piai, Pontian, Johor (Station 5)	56
	4.1.6 Diatom Distribution along the Tebrau Straits	58
	4.2 Recoverability of Diatom from Organ Samples of	59
	Suspected Drowning Cases	
	4.2.1 Recoverability of Diatom in Human Organs	60
5	CONCLUSION	63
	5.1 Summary of Findings	63
	5.2 Limitations and Recommendations	65

REFERENCES

66

LIST OF TABLES

TABLE NO.	TITLE	PAGE
3.1	List of sampling stations and substations with	30
	their geographic coordinates	
3.2	List of organ samples collected from bodies for	31
	diatom analysis	
4.1	The taxonomic traits used for identifying the	36
	different genera of diatoms recovered in this	
	study (Shamsudin, 1990, 1991; Spaulding et al.,	
	2010).	
4.2	The overall percentage assemblage of diatom at	50
	Lebam River (Station 1)	
4.3	The overall percentage assemblage of diatom at	52
	Kabong Village (Station 2)	
4.4	The overall percentage assemblage of diatom at	54
	Johor Causeway (Station 3)	
4.5	The overall percentage assemblage of diatom at	56
	Pendas (Station 4)	
4.6	The overall percentage assemblage of diatom at	57
	Tanjung Piai (Station 5)	
4.7	List of sampling stations with prevalent diatoms	59
	genera	
4.8	Quantity of diatoms recovered from organ	61
	samples of bodies	

LIST OF FIGURES

FIGURE NO.	TITLE	PAGE
2.1	Frustule of diatom	6
2.2	Sub-order of pennate diatoms	8
3.1	Five sampling stations along the Tebrau Straits and	
	Lebam River	
5.1	Prevalent diatoms at sampling stations along the	64
	Tebrau Straits	

LIST OF ABBREVIATION

CO_2	Carbon dioxide
Cu	Copper
DOE	Department of energy
Е	East
FAAS	Flameless atomic absorption spectrometry
FC	Fecal coliform
Fe	Iron
FS	Fecal streptococci
g	Gram
ICP-MS	Inductively coupled plasma mass spectrometry
km	Kilometre
L	Litre
LV	Left ventricle
m	Metre
min	Minute
mL	Millilitres
mm	Millimetres
Μ	Molar
Ν	North
NaCl	Sodium Chloride
pН	Potential hydrogen
Pb	Lead
RV	Right ventricle
rRNA	Ribosomal ribonucleic acid
rpm	Revolution per minute

Si(OH) ₄	Silicic acic
SDS	Sodium dodecyl sulfate
SEM	Scanning electron microscope
sp.	Species
Sr	Strontium
TH	Tyrosine hyroxylase
USA	United States of America
μm	Micrometer
μL	Microlitre
Zn	Zinc
SYMBOLS	
°C	Degree Celcius
%	Percent

CHAPTER 1

INTRODUCTION

1.1 Background of study

In medico-legal investigation of bodies recovered from water, the aim of the investigations often focuses on determination of the cause and manner of death along with the site of death. Deaths in water may be due to drowning, injuries, intoxications or natural causes and, in this context diatoms in airways as well as organs are widely studied (Lunetta and Modell, 2005). Diatom is a microscopic unicellular aquatic algae belonging to *Bacillariophyta* (Pollanen, 1997) and first described in drowning victim at the end of 19th century (Revenstof, 1904). Since then, diatom test has been used as forensic evidence in cases involving bodies recovered from water (Lunetta and Modell, 2005).

Diatom test is considered as the gold standard in diagnosing ante-mortem or post-mortem drownings and assigning drowning site especially when dealing with bodies recovered from water (Piette, 2006). In general, ante-mortem drowning caused diatoms to reach lungs *via* water aspiration, prior to being distributed to other organs through bloodstream. On the contrary, post-mortem drowning caused diatoms to passively enter airways and not travelling to the other organs due to lack of circulation system after death (Lunetta and Modell, 2005). Comparing the diatoms found in organ sample from drowned bodies with diatoms recovered from water samples can also suggest the drowning site (Vinayak *et al.*, 2013). The test required complete destruction of other particles than diatoms (such as organ and debris), excluding the diatom frustules (Lunetta and Modell, 2005). Ming *et al.* (2007) indicated the common extraction techniques applied in diatom test are digestion using (1) strong acid (hydrochloric acid, nitric acid), (2) using solubilizers (soluene-350) and (3) enzymatic digestion (Proteinase K).

1.2 Problem Statement

Diatoms are location specific, attributable to variations in salinity, temperature and presence of organic matters (Cameron, 2004), and hence, may prove useful in assigning the site of drowning. However, due to unavailability forensic baseline data within Johor marine ecology, investigation concerning site of drowning can be difficult. Hence, it is pertinent to conduct this study for exploring the diatom distributions and its mapping, specifically within the bustling waters of the Tebrau Straits, in view of its application in assigning drowning and suggesting the possible localities.

In Malaysia, forensic pathologists always rely on histopathological findings to draw conclusions pertaining to deaths due to drowning and, utilization of diatoms in this aspect remains unreported. This may be due to the lack of information and understanding on its recoverability in human organs, limiting its application in forensic cases. Considering such limitation, the specific attempts made in this present study to recover diatoms in such organs as lungs, kidneys and livers using the acidic digestion method, merits consideration.

1.3 Objectives

This study specifically designed:

- (a) To establish diatomological mapping related to the findings of diatoms assemblages found at several locations along the Tebrau Straits.
- (b) To study the recoverability of diatoms from organ samples collected from suspected drowning cases using acid-digestion method.

1.4 Scope of Study

This study involved the collection of seawater samples from Lebam River, Kabong Village, Johor Causeway, Pendas and Tanjung Piai of the Tebrau Straits. In addition, organ samples from bodies, presumably died because of drowning during December 2012 and April 2015, collected by a forensic pathologist at Hospital Sultanah Aminah Johor Bahru were also analyzed. Extraction of diatoms from water and organ samples was made using the acid digestion method suggested by the previous researchers (Yen and Jayaprakash, 2007; Malik and Kaidan, 2013). Upon identifying the genera of diatoms following the identification keys provided by previous researchers (Shamsudin, 1990, 1991; Spaulding *et al.*, 2010), qualitative and quantitative assessments were made.

1.5 Significance of Study

In addition to providing the diatomological mapping for the Tebrau Straits, the data reported here may be useful in real forensic cases for indicating the site of drowning, whenever specific assemblages of diatoms are recovered from the different organs of victims, purportedly drowned at locations along the Straits. Such aspect acquires forensic significance for suggesting the primary crime scenes, an important element for establishing the fact of the case as well as disputing an alibi.

REFERENCES

- Abdullah, M (1997). Hydrocarbon Pollution in the Sediment of Some Malaysian Coastal Areas. *Environmental Monitoring and Assessment*, 44(1-3), 443454.
- Ago, K., Ago, M., & Ogata, M. (2004). The Distribution of Diatoms in Yoronjima and Application of the Diatom Test for the Diagnosis of Death by Drowning in Open Sea Islands. *Medical Journal-Kagoshima University*, 56, 25-30.
- Ahmad, F., Azman, S., & Mohd Said, M. (2014). Distribution of Metal Contaminants in the Straits of Johor Due to Local Development. In International Conference on Chemical, Biological, and Environmental Sciences (ICCBES'14. Kuala Lumpur.
- Alcoverro, T., Conte, E., & Mazzella, L. (2000). Production of Mucilage by the Adriatic Epipelic Diatom Cylindrotheca Closterium (Bacillariophyceae) Under Nutrient Limitation. *Journal of Phycology*, *36*(6), 1087-1095.
- Amo, Y. D., & Brzezinski, M. A. (1999). The Chemical Form of Dissolved Si Taken Up by Marine Diatoms. *Journal of Phycology*, 35(6), 1162-1170.
- Auer, A. (1991). Qualitative Diatom Analysis as A Tool to Diagnose Drowning. *The American Journal Of Forensic Medicine and Pathology*, *12*(3), 213-218.
- Auer, A. & Mottonen, M. (1988). Diatoms and drowning. Zeitschrift For Rechtsmedizin, 101(2).
- Awang, N., Jusoh, W., & Hamid, M. (2014). Coastal Erosion at Tanjong Piai, Johor, Malaysia. *Journal of Coastal Research*, 71, 122-130.
- Azman, S., Chiang, B., Ismail, R., Jaafar, J., Said, M., & Salmiati, S. (2012). Effect of Land Use on Coastal Water and Perna viridis at Johor Straits, Malaysia. *International Journal of Environmental Science And Development*, 237-239.
- Azparren, J., Vallejo, G., Reyes, E., Herranz, A., & Sancho, M. (1998). Study of the Diagnostic Value of Strontium, Chloride, Haemoglobin and Diatoms in Immersion Cases. *Forensic Science International*, 91(2), 123-132.
- Barker, P., Fontes, J., & Gasse, F. (1994). Experimental Dissolution of Diatom Silica in Concentrated Salt Solutions and Implications for Paleoenvironmental Reconstruction. *Limnology and Oceanography*, 39(1), 99-110.

- Battarbee, R. W., Juggins, S., Gasse, F., Anderson, N. J., Bennion, H., Cameron, N. G. & Telford, R. (2001). An Information System for Palaeoenvironmental Reconstruction.
- Bere, T. & Tundisi, J. (2010). Epipsammic Diatoms in Streams Influenced by Urban Pollution, São Carlos, SP, Brazil. *Brazilian Journal of Biology*, 70(4), 920-930.
- Bidle, K. D., & Azam, F. (1999). Accelerated Dissolution of Diatom Silica by Marine Bacterial Assemblages. *Nature*, 397(6719), 508-512.
- Bucholtz, A. (2014). An Introduction to Forensic Pathology for the Nonscientist (1st ed., pp. 289-291). New York: Anderson Publishing.
- Buesseler, K. (1998). The Decoupling of Production and Particulate Export in the Surface Ocean. *Global Biogeochemical Cycles*, *12*(2), 297-310.
- Cameron, N. (2004). The Use of Diatom Analysis in Forensic Geoscience. Geological Society, London, Special Publications, 232(1), 277-280.
- Casamatta, D. & Verb, R. (2000). Algal Colonization of Submerged Carcasses in A Mid-Order Woodland Stream. *Journal of Forensic Sciences*, *45*(6), 14880J.
- Clavero, E., Hernandez-Marine, M., Grimalt, J., & Garcia-Pichel, F. (2000). Salinity Tolerance of Diatoms From Thalassic Hypersaline Environments. *Journal of Phycology*, 36(6), 1021-1034.
- Coelho, S., Ramos, P., Ribeiro, C., Marques, J., & Santos, A. (2016). Contribution to the Determination of the Place of Death by Drowning – A Study of Diatoms' Biodiversity In Douro River Estuary. *Journal Of Forensic And Legal Medicine*, 41, 58-64
- Department of Fisheries Malaysia Official Portal. (2013). www.dof.gov.my.
- Department of Statistics Malaysia Official Portal. (2016). Statistics.gov.my. Retrieved 10 December 2016.
- de Brouwer, J. F., & Stal, L. J. (2002). Daily Fluctuations of Exopolymers in Cultures of the Benthic Diatoms *Cylindrotheca closterium and Nitzschia* sp.
- de la Grandmaison, G., Leterreux, M., Lasseuguette, K., Alvarez, J., Mazancourt, P., & Durigon, M. (2006). Study of the Diagnostic Value of Iron in Freshwater Drowning. *Forensic Science International*, 157(2-3), 117-120.
- Díaz-Palma, P., Alucema, A., Hayashida, G., & Maidana, N. (2009). Development and Standardization of A Microalgae Test for Determining Deaths By Drowning. *Forensic Science International*, 184(1-3), 37-41.

- Dolinak, D., Matshes, E., & Lew, E. (2005). *Forensic Pathology* (1st ed.). Amsterdam: Elsevier Academic Press.
- Foged, N. (1983). Diatoms in Human Tissues, Greenland ab. 1460 AD-Funen 198182 AD. Nova Hedwigia: Zeitschrift fur Kryptogamenkunde.
- Frost, B. (1991). The Role of Grazing in Nutrient-Rich Areas of the Open Sea. *Limnology and Oceanography*, 36(8), 1616-1630.
- Fukui, Y., Hata, M., Takahashi, S., & Matsubara, K. (1980). A New Method for Detecting Diatoms in Human Organs. *Forensic Science International*, 16(1), 67-74.
- Gunn, A. (2006). Essential forensic biology (1st ed.). Chichester, England: J. Wiley.
- Hadibarata, T., Abdullah, F., Yusoff, A., Ismail, R., Azman, S., & Adnan, N. (2012).
 Correlation Study between Land Use, Water Quality, and Heavy Metals (Cd, Pb, and Zn) Content in Water and Green Lipped Mussels Perna viridis (Linnaeus.) at the Johor Strait. *Water, Air, & Soil Pollution, 223*(6), 3125-3136.
- Hasle, G. & Syvertsen, E. (1996). Marine Diatoms. In G. Hasle, E. Syvertsen, K. Steidinger, K. Tangen & C. Tomas, *Identifying Marine Diatoms and Dinoflagellates* (pp. 1-381). Academic Press.
- Hoagland, K., Rosowski, J., Gretz, M., & Roemer, S. (1993). Diatom Extracellular Polymeric Substances: Function, Fine Structure, Chemistry, And Physiology. *Journal Of Phycology*, 29(5), 537-566.
- Hoek, C., Mann, D., & Jahns, H. (1995). *Algae* (1st ed.). Cambridge: Cambridge University Press.
- Hildebrand, M. (2006). 10 Silicic Acid Transport and its Control During Cell Wall Silicification in Diatoms. *Biomineralization: Progress in Biology, Molecular Biology and Application*.
- Hussein, M. & Mohamed, N. (2007). Wetland Landscape Conservation and Ecotourism in Malaysia. *Pertanika Journal Social Science & Humanities*, 5(2), 159-168.
- Polis Diraja Malaysia,. Laporan Kes Lemas Dalam Tempoh 5 Tahun bagi Balai Polis Sentral.Unpublished data
- Peabody, A. (1977). Diatoms in Forensic Science. Journal of the Forensic Science Society, 17(2-3), 81-87

- Peabody, A. (1980). Diatoms and Drowning A review. *Medicine, Science and The Law*, 20(4), 254-261.
- Ismail, M. (2009). Application of A Hydrodynamic Water Quality Model in Sungai Johor Estuary (Ph.D). Universiti Teknologi Malaysia..
- Jamieson, A. & Moenssens, A. (2009). Wiley Encyclopedia of Forensic Science (1st ed.). Chichester, West Sussex, U.K.: John Wiley & Sons.
- Jusoff, K. & Bin Hj Taha, D. (2009). Managing Sustainable Mangrove Forests in Peninsular Malaysia. Journal of Sustainable Development, 1(1).
- Kaczmarska, I., Poulíčková, A., Sato, S., Edlund, M., Idei, M., Watanabe, T., &
 Mann, D. (2013). Proposals for A Terminology for Diatom Sexual Reproduction, Auxospores and Resting Stages. *Diatom Research*, 28(3), 263-294.
- Kakizaki, E., Takahama, K., Seo, Y., Kozawa, S., Sakai, M., & Yukawa, N. (2008). Marine Bacteria Comprise A Possible Indicator Of Drowning in Seawater. *Forensic Science International*, 176(2-3), 236-247.
- Kakizaki, E., Kozawa, S., Sakai, M., & Yukawa, N. (2009). Bioluminescent Bacteria Have Potential As A Marker Of Drowning in Seawater: Two Immersed Cadavers Retrieved Near Estuaries. *Legal Medicine*, 11(2), 91-96.
- Kakizaki, E., Kozawa, S., Imamura, N., Uchiyama, T., Nishida, S., Sakai, M., & Yukawa, N. (2011). Detection Of Marine and Freshwater Bacterioplankton in Immersed Victims: Post-Mortem Bacterial Invasion Does Not Readily Occur. *Forensic Science International*, 211(1-3), 9-18.
- Kakizaki, E., Ogura, Y., Kozawa, S., Nishida, S., Uchiyama, T., Hayashi, T., & Yukawa, N. (2012). Detection of Diverse Aquatic Microbes in Blood and Organs of Drowning Victims: First Metagenomic Approach Using High-Throughput 454-Pyrosequencing. *Forensic Science International*, 220(1-3), 135-146.
- Kazazi, M. (1996). *Burden of Proof and Related Issues* (1st ed.). The Hague [u.a.]: Kluwer Law International.
- Knight, B. & Saukko, P. (2015). *Knight's Forensic Pathology* (4th ed., pp. 399-414). CRC Press.
- Kobayashi, M., Yamada, Y., Zhang, W., Itakura, Y., Nagao, M., & Takatori, T. (1993). Novel Detection of Plankton from Lung Tissue by Enzymatic Digestion Method. *Forensic Science International*, 60(1-2), 81-90.

- Kohlhase, C. & Maxeiner, H. (2003). Morphometric Investigation of Emphysema Aquosum in the Elderly. *Forensic Science International*, *134*(2-3), 93-98.
- Krstic, S., Duma, A., Janevska, B., Levkov, Z., Nikolova, K., & Noveska, M. (2002). Diatoms in Forensic Expertise of Drowning—A Macedonian Experience. *Forensic Science International*, 127(3), 198-203.
- Kuczynska, P., Jemiola-Rzeminska, M., & Strzalka, K. (2015). Photosynthetic Pigments in Diatoms. *Marine Drugs*, *13*(9), 5847-5881.
- Latifah, A. & Met, L. (2014). An Ecological Evaluation Approach for Dam Project Development in Malaysia. *Life Science Journal*, 11(7), 225-237.
- Lucci, A. & Cirnelli, A. (2007). A Microbiological Test for the Diagnosis of Death by Drowning. *Forensic Science International*, *168*(1), 34-36.
- Lucci, A., Campobasso, C., Cirnelli, A., & Lorenzini, G. (2008). A Promising Microbiological Test for the Diagnosis of Drowning. *Forensic Science International*, 182(1-3), 20-26.
- Ludes, B., Quantin, S., Coste, M., & Mangin, P. (1994). Application of A Simple Enzymatic Digestion Method for Diatom Detection in the Diagnosis of Drowning in Putrefied Corpses by Diatom Analysis. *International Journal Of Legal Medicine*, 107(1), 37-41.
- Ludes, B., Coste, M., North, N., Doray, S., Tracqui, A., & Kintz, P. (1999). Diatom Analysis in Victim's Tissues as An Indicator of the Site of Drowning. *International Journal of Legal Medicine*, 112(3), 163-166.
- Lunetta, P. & Modell, J. (2005). Macroscopical, Microscopical, and Laboratory Findings in Drowning Victims. In *Forensic Pathology Reviews* (1st ed., pp. 3-77). Humana Press.
- Lunetta, P., Penttilä, A., & Hällfors, G. (1998). Scanning and Transmission Electron Microscopical Evidence of the Capacity Of Diatoms to Penetrate the Alveolo-Capillary Barrier in Drowning. *International Journal Of Legal Medicine*, 111(5), 229-237.
- Malik, M., Jakhar, P., & Kadian, A. (2013). Role of Diatoms in Forensic Investigation: Case Studies from Haryana. *International Journal Of Forensic Science & Pathology*, 11-12.
- Matsumoto, H. & Fukui, Y. (1993). A Simple Method for Diatom Detection in Drowning. *Forensic Science International*, 60(1-2), 91-95.

- Medlin, L., Kooista, W., & Anderson, R. (1997). Phylogenetic Relationships of the 'Golden Algae' (Haptophytes, Heterokont Chromophytes) and Their Plastids.
 In D. Bhattacharya, *Origins of Algae And Their Plastids* (1st ed., pp. 187-219). Vienna: Springer.
- Ming, M., Meng, X., & Wang, E. (2007). Evaluation of Four Digestive Methods for Extracting Diatoms. *Forensic Science International*, 170(1), 29-34.
- Mishulsky, A.M. (1990). Use of Bacteriologic Analysis of Blood in Diagnosis of Death from Drowning. Sud Med Ekspert, 33, 26–28 [in Russian with English abstract]
- Munn, C. (2011). Marine Microbiology (1st ed.). Garland Science.
- Nielsen, E. & Wium-Andersen, S. (1970). Copper Ions as Poison in the Sea and in Freshwater. *Marine Biology*, 6(2), 93-97.
- Piette, M. & De Letter, E. (2006). Drowning: Still a Difficult Autopsy Diagnosis. Forensic Science International, 163(1-2), 1-9.
- Pollanen, M. (1997). The Diagnostic Value of the Diatom Test for Drowning, II. Validity: Analysis of Diatoms in Bone Marrow and Drowning Medium. *Journal of Forensic Sciences*, 42(2), 14112J.
- Pollanen, M. (1998). Diatoms and Homicide. *Forensic Science International*, 91(1), 29-34.
- Pounder, D. (2005). Drowning. In R. Byard & J. Payne-James, *Encyclopedia of Forensic and Legal Medicine*, (1st ed., pp. 227-232). Academic Press.
- Revenstorf, V. (1904). Der Nachwies der aspirierten Ertrankungs flussigkeit als Kriterium des Todes, *Gerichtliche Medizin* **28**, 274–279.
- Rahmana, N. & Najibb, A. (2015). A Maritime Forecasting Method for Analysing the Total Cargo Handling at Johor Port Berhad from 2013 to 2020. *International Journal Of Business Management & Economic Research*, 6(3).
- Ryves, D., Juggins, S., Fritz, S., & Battarbee, R. (2001). Experimental Diatom Dissolution and the Quantification of Microfossil Preservation in Sediments. *Palaeogeography, Palaeoclimatology, Palaeoecology, 172*(1-2), 99-113.
- Saferstein, R. (2015). Forensic Science: From the Crime Scene to the Crime Lab (3rd ed.). Pearson.
- Santos, B., Martins, S., Almeida, A., Santos, A., & Magalhães, T. (2010). Diagnosing Death by Drowning Through the Analysis of Blood Markers by

Inductively Coupled Plasma-Mass Spectrometry. Acta Medicinae Legalis et Socialis.

- Sasidharan, A. & S, Resmi. (2013). Forensic Diatomology Review. *Health Sciences E-Journal Kerala University Of Health Sciences*, 3(1).
- Sarthou, G., Timmermans, K., Blain, S., & Tréguer, P. (2005). Growth Physiology and Fate of Diatoms in the Ocean: A Review. *Journal of Sea Research*, 53(1), 25-42.
- Shamsudin., L. (1990). *Diatom Marin Di Perairan Malaysia* (1st ed.). Kuala Lumpur: Dewan Bahasa dan Pustaka, Kementerian Pendidikan Malaysia.
- Shamsudin., L. (1991). *Diatom Air Tawar* (1st ed.). Kuala Lumpur: Dewan Bahasa dan Pustaka.
- Sidari, L., Di Nunno, N., Costantinides, F., & Melato, M. (1999). Diatom test with Soluene-350 to Diagnose Drowning in Seawater. *Forensic Science International*, 103(1), 61-65.
- Singh, R., Singh, R., & Thakar, M. (2006). Extraction methods of Diatoms-A Review. *Indian Internet Journal of Forensic Medicine & Toxicology*, 4(2).
- Sitthiwong, N., Ruangyuttikarn, W., Vongvivach, S., Peerapornpisal, Y., Charoonnate, N., & Narongchai, P. (2016). Detection and Identification of Diatoms in Tissue Samples of Drowning Victims. *Chiang Mai Journal Science*, 41(5.1), 1020-1031.
- Siver, P., Lord, W., & McCarthy, D. (1994). Forensic Limnology: The Use of Freshwater Algal Community Ecology to Link Suspects to an Aquatic Crime Scene in Southern New England. *Journal of Forensic Sciences*, 39(3), 13663J.
- Spaulding, S.A., Lubinski, D.J. and Potapova, M. (2010). Diatoms of the United States. http://westerndiatoms.colorado.edu
- Staats, N., De Winder, B., Stal, L., & Mur, L. (1999). Isolation and Characterization of Extracellular Polysaccharides from the Epipelic Diatoms Cylindrotheca Closterium and Navicula salinarum. *European Journal of Phycology*, 34(2), 161-169.
- Sterrenburg, F., Gordon, R., Tiffany, M., & Nagy, S. (2007). Diatoms. In J. Seckbach, Algae and Cyanobacteria in Extreme Environments (11th ed., pp. 141-172). Netherlands: Springer.

- Su, H., Liu, C., Hu, S., Liu, Y., Xu, X., Liu, H., & Zhao, J. (2012). Diatomological Distribution of Pearl River in Guangzhou and Yangtze River in Hubei Province and Its Application to the Diagnosis Of Drowning Cases. In International Conference on Biomedical Engineering and Biotechnology.
- Subramanian, A., Subramanian, B., & Venugopalan, V. (1980). Toxicity of Copper and Zinc on Cultures of Skeletonema costatum (Grev.) Cleve and Nitzschia longissima. *Current Science*, 49(7), 266-268.
- Tamaska, L. (1949). Diatom Content of Bone Marrow in Corpses in Water. Orv. Hetil, 16, 509-511.
- Thomas, F., Van Hecke, W., & Timperman, J. (1961). The Detection of Diatoms In the Bone Marrow as Evidence of Death by Drowning. *Journal of Forensic Medicine*, 8, 142-144.
- Timperman, J. (1972). The Diagnosis of Drowning. A review. *Forensic Science*, *1*(4), 397-409.
- University College London, (2016). *Pennate diatom*.Retrieved from http://www.ucl.ac.uk/GeolSci/micropal/diatom.html
- University College London,. (2016). *Frustule of diatom*.Retrieved from http://www.ucl.ac.uk/GeolSci/micropal/diatom.html
- Vinayak, V., Mishra, V., & MK, G. (2013). Diatom Fingerprinting to Ascertain Death in Drowning Cases. *Journal of Forensic Research*, 04(05).
- Wehr, J., Sheath, R., & Kociolek, J. (2014). Freshwater Algae of North America: Ecology and Classification (2nd ed.). Academic Press.
- Wetli, C., Mittleman, R., & Rao, V. (1999). *An Atlas of Forensic Pathology* (1st ed.). ASCP Press.
- Willers, V. (2008). Effect of Zinc Availability on the Morphology and Nutrient Physiology of A Coastal and An Oceanic Diatom (Ph.D). University of Victoria.
- Wilmans. (1896). Ueber den Tod durch Ertrinken. Vjschr Gerichtl Med, 12 (Suppl 12), 363–81.
- Wood, A., Ahmad, Z., Md. Shazili, N., Yaakob, R., & Carpenter, R. (1997). Geochemistry of Sediments in Johor Strait between Malaysia and Singapore. *Continental Shelf Research*, 17(10), 1207-1228.

- Yange, L., Chuanying, H., Chengxing, W., & Xu, W. (1999). Development of Can for Destruction of Organic Material in Use for Forensic Diatom Examination. *Forensic Science International*, 101(3), 163-166.
- Yap, C., Ismail, A., & Tan, S. (2004). Heavy metal (Cd, Cu, Pb and Zn) Concentrations in the Green-Lipped Mussel Perna viridis (Linnaeus) collected from Some Wild and Aquacultural Sites in the West Coast of Peninsular Malaysia. *Food Chemistry*, 84(4), 569-575.
- Yap, C., Ismail, A., Edward, F., Tan, S., & Siraj, S. (2006). Use of Different Soft Tissues of Perna viridis as Biomonitors of Bioavailability and Contamination by Heavy Metals (Cd, Cu, Fe, Pb, Ni, and Zn) in A Semi-Enclosed Intertidal Water, the Johore Straits. *Toxicological & Environmental Chemistry*, 88(4), 683-695.
- Yap, C., Shahbazi, A., & Zakaria, M. (2012). Concentrations of Heavy Metals (Cu, Cd, Zn and Ni) and PAHs in Perna viridis Collected from Seaport and Nonseaport Waters in the Straits of Johore. *Bulletin of Environmental Contamination and Toxicology*, 89(6), 1205-1210.
- Yen, L. & Jayaprakash, P. (2007). Prevalence of Diatom Frustules in Non-Vegetarian Foodstuffs and its Implications in Interpreting Identification of Diatom Frustules in Drowning Cases. *Forensic Science International*, 170(1), 1-7.
- Yukawa, N., Kakizaki, E., & Kozawa, S. (2013). Diatom and Laboratory Tests to Support A Conclusion of Death by Drowning. In G. Rutty, *Essentials of Autopsy Practice* (1st ed., pp. 1-36). London: Springer
- Zhuo, L., Lin, S., & Song, P. (2016). A Future Perspective in Forensic Medicine in China: Diatomological Mapping of Water Bodies. *Journal of Forensic Science and Medicine*, 2(2), 107.
- Zulkifli, S., Mohamat-Yusuff, F., Arai, T., Ismail, A., & Miyazaki, N. (2009). An Assessment of Selected Trace Elements in Intertidal Surface Sediments Collected from the Peninsular Malaysia. *Environmental Monitoring and* Assessment, 169(1-4), 457-472.
- Zulkifli, S., Mohamat-Yusuff, F., Ismail, A., Aziz, A., Azrin Sabuti, A., & Mohamed, C. (2015). Status of Heavy Metals in Surface Sediments of the Western Part of the Johor Straits Using a Sediment Quality Guideline. World Journal of Fish and Marine Sciences, 7(3), 214-222.