

SUPER ADSORBENT-BASED REMEDIATION AND ON-SITE FLOW
INJECTION ANALYSIS DETERMINATION OF ARSENIC IONS IN GOLD
MINING SITE

NABEL KALEL ASMEL

A thesis submitted in fulfilment of the
requirements for the award of the degree of
Doctor of Philosophy (Civil Engineering)

Faculty of Civil Engineering
Universiti Teknologi Malaysia

JULY 2017

SPECIAL DEDICATIONS TO:

My beloved parents, wife and children Yousif, Tiba, Mohammad, and Toqa, supportive siblings and to all my friends. May ALLAH always bless us now and forever.

My respected project supervisor

Prof. Dr. Abdull Rahim bin Mohd Yusoff

My dedicated co-supervisor

Assoc. Prof. Dr. Zaiton Abdul Majid

Dr. Salmiati

Thanks you for always helping and supporting me

ACKNOWLEDGEMENT

All praise is to Allah, the Lord of the universe, and thanks to Allah for His guidance, opportunity and for giving me the strength, patience and blessing to finish up this research with full of His helpfulness. I would like to express my utmost gratitude and appreciation to my sincere and supportive supervisors, Prof. Dr. Abdull Rahim Mohd Yusoff, Assoc. Prof. Dr. Zaiton Abdul Majid, and Dr. Salmiati for their determined guidance in completing this research project. With their great advice, valuable suggestions and monitors since from the beginning to finish up to this supreme level. Your critics, continuous support, encouragement, patience and inputs are really appreciated.

I would like to express my thanks to the staff of Centre for Environmental Sustainability and Water Security (IPASA). I am also very thankful to all my research associate especially Ahmed Rahomi, Faizuan Bin Abdullah, Syaiful Akmal, Jasmin Fathi Jasni, Azizul Bin Mohd Zahari, Mimi Suliza Binti Muhamad for sharing their knowledge, providing me assistance and regular discussions which really helped me in my lab work. I would also like to thank the technical staff from the Department of Environmental Engineering (Faculty of Civil Engineering) for their assistance during my lab work. I am grateful to the Ministry of Higher Education and scientific research, Northern Technical University, Iraq for providing scholarship throughout my study. This support is greatly appreciated. Last but not least, I would like to thank my parents, wife and children, for their patience, continuous support and encouragement in completing my PhD. May ALLAH bless all of us.

ABSTRACT

Arsenic is a mineral which is abundant in gold mining sites. It is a toxic substance which needs to be removed. Thus, this research investigated the problem of extremely high arsenic (As) concentration in effluent (water and slurry) from Selinsing gold mine site, Pahang, Malaysia and developed an effective operational remedial method using an adsorbent material (Ecomel). In this research, a new method for measuring As(III) and As(V), which exist in the contaminated soil and liquid effluents was developed using flow injection analysis (FIA) system, while the total As was analyzed using atomic fluorescence spectrometry (AFS) for the remediating samples. By using the FIA method, the As speciation detection limit of ($S/N = 3$) for both As(III) and As(V) were found to be $5 \mu\text{g/L}$ with standard deviation 2.2 ($n = 20$). It also showed a wide dynamic range coefficient of detection (R^2) of 0.999 for As(III) and 0.9989 for As(V). This portable analytical method was successfully applied for the determination of As speciation in the effluent collected from Selinsing gold mine site without further treatment. For the characterizations of slurry and adsorbent materials (Ecomel), X-ray diffraction (XRD), X-ray fluorescence (XRF), energy-dispersive X-ray spectroscopy (EDX), Field Emission Scanning Electron Microscopy (FESEM), Focus Ion Beam–Scanning Electron Microscope (FIB-SEM), Inductively Coupled Plasma Optical Emission Spectrophotometer (ICPOES), Elemental Analyzer (EA) and Brunauer–Emmett–Teller (BET) were used. In the case of liquid effluent from the tailing dam, the pH and cyanide concentrations were measured at 11.5 and 204 mg/L, respectively. However, the batch experiments clearly proved that As leaching from the slurry (contact time: 24 h, stirring speed: 200 rpm, and S:L ratio 1:5) was extremely higher at pH 11.5 compared to pH 2.5 and were measured to be 8,720 and 1,010 $\mu\text{g/L}$, respectively. The major contributors to the exceedingly elevated levels of As concentration in liquid effluents were attributed to alkaline pH, high cyanide and silicate concentration, as well as high oxidation environment. The batch experiments on Ecomel revealed that maximum adsorption capacity determined at initial pH 2.5 from the Langmuir-Freundlich isotherm model was found to be 704.7 and 122.7 mg/g for As(III) and As(V), respectively. These results indicated that Ecomel has high adsorbent efficiency, cost-effective and is suitable for in-situ and ex-situ remediation of highly concentrated As(III) and As(V) toxicants in aqueous solutions. For As standard solution with initial pH 2.5 and concentration of As at 1 mg/L, it was observed that 0.3125 g/L of Ecomel with 2 h contact time can adequately remove 97.0% of As(V) and 98.1% of As(III), respectively. As a conclusion, results from AFS showed that the treatment of As using Ecomel at initial pH 2.5 was the most proficient for remediation of liquid effluent and slurry.

ABSTRAK

Arsenik adalah mineral yang terkandung dengan banyak dalam lombong emas. Ia merupakan bahan toksik yang perlu disingkirkan. Justeru itu, kajian ini mengkaji tentang masalah arsenik yang tinggi di dalam kandungan efluen (air dan lumpur) daripada tapak lombong emas Selinsing, Pahang, Malaysia dan membangunkan satu kaedah pemulihan berkesan menggunakan bahan penjerap (Ecomel). Dalam kajian ini satu kaedah baru bagi pengukuran As(III) dan As(V) yang terkandung di dalam tanah yang tercemar dan cecair efluen juga telah dibangunkan menggunakan sistem analisis suntikan aliran (FIA), manakala jumlah As bagi sampel pemulihan telah dianalisis dengan menggunakan spektrometri pendarfluor atom (AFS). Dengan menggunakan kaedah FIA ini, had pengesanan ($S/N = 3$) penspesiesan As bagi kedua-dua As(III) dan As(V) masing-masing ialah $5 \mu\text{g/L}$ dan 2.2% ($n = 20$). Kaedah ini menunjukkan julat dinamik yang luas dengan pekali korelasi (R^2) 0.999 untuk As(III) dan 0.9989 untuk As(V). Analisis mudah alih ini telah berjaya digunakan untuk penspesiesan As di dalam efluen yang diambil dari lombong emas Selinsing tanpa memerlukan rawatan lanjut. Untuk pencirian buburan dan bahan penjerap (Ecomel), pembelauan sinar-x (XRD), pendarfluor sinar-x (XRF), tenaga serakan sinar-x spektroskopi (EDX), mikroskopi medan pemancaran pengimbasan elektron (FESEM), mikroskop fokus ion alur-pengimbasan elektron (FIB-SEM), spektrofotometer plasma gandingan aruhan pemancaran optik (ICPOES), penganalisa unsur (EA) dan Brunauer-Emmett-Teller (BET) telah digunakan. Dalam kes efluen cecair dari empangan tailing, ukuran pH dan kepekatan sianida masing-masing adalah 11.5 dan 204 mg/L . Walau bagaimanapun, eksperimen kelompok jelas membuktikan bahawa ketelaruhan lesap As dari buburan (masa sentuhan: 24 h , kelajuan kacau: 200 rpm , dan nisbah S:L $1:5$) adalah amat tinggi pada pH 11.5 berbanding pada pH 2.5 , masing-masing dengan ukuran $8,720$ dan $1,010 \mu\text{g/L}$. Penyumbang utama kepada tahap kepekatan As yang sangat tinggi dalam efluen cecair adalah disebabkan oleh pH beralkali, kepekatan sianida dan silikat yang tinggi, dan di tambah dengan keadaan pengoksidaan sekeliling yang tinggi. Eksperimen kelompok menggunakan Ecomel menunjukkan bahawa kapasiti penjerapan maksimum pada pH awal 2.5 dari model isoterma Langmuir-Freundlich masing-masing adalah 704.7 dan 122.7 mg/g untuk As(III) dan As(V). Dapatan ini menunjukkan bahawa Ecomel mempunyai kecekapan penjerapan yang tinggi, kos efektif dan sangat sesuai digunakan untuk pemulihan *in-situ* dan *ex-situ* bagi larutan yang mengandungi bahan toksik As(III) dan As(V) dalam kepekatan yang tinggi. Bagi larutan standard yang mengandungi As pada pH awal 2.5 dengan kepekatan 1 mg/L , didapati bahawa Ecomel dengan kepekatan 0.3125 g/L dan dengan masa sentuhan 2 h , dapat menyingkirkan 97.0% As(V) dan 98.1% As(III). Sebagai kesimpulan, dapatan daripada AFS menunjukkan rawatan As menggunakan Ecomel pada pH awal 2.5 adalah yang paling berkesan bagi pemulihan cecair efluen dan buburan.

TABLE OF CONTENTS

CHAPTER	TITLE	PAGE
	DECLARATION	ii
	DEDICATION	iii
	ACKNOWLEDGEMENT	iv
	ABSTRACT	v
	ABSTRAK	vi
	TABLE OF CONTENTS	vii
	LIST OF TABLES	xiv
	LIST OF FIGURES	xvi
	LIST OF ABBREVIATIONS	xxi
	LIST OF APPENDICES	xxviii
1	INTRODUCTION	1
	1.1 Background of the study	1
	1.2 Problem statement	5
	1.3 Research objective	7
	1.4 Scope of research	7
	1.5 Significance of research	11
	1.6 Limitation of research	12

2	LITERATURE REVIEW	14
2.1	Gold production in Malaysia	14
2.2	Sustainable mining	18
2.3	Arsenic speciation in environment	20
2.4	Arsenic toxicity and humans health effects	21
2.5	Gold mining processes	22
2.5.1	Crushing, grinding and sizing	24
2.5.2	Leaching, adsorption and filtration the metal from rock	24
2.5.3	Recovery of gold to generate dore bullion bars	25
2.6	Arsenopyrite and pyrite oxidation	26
2.7	Environmental impact of arsenopyrite and pyrite oxidation	27
2.8	Acid mine drainage	28
2.8.1	Sources and remediation methods of acid mine drainage	29
2.9	Critical review about the As contamination in Pahang-Malaysia	31
2.10	Gold mining site	33
2.11	Factors affecting of As leaching	34
2.11.1	pH	35
2.11.2	Oxidation-Reduction Potential	35
2.11.3	Silica	36
2.11.4	Oxidation processes	37
2.11.5	Cyanide	38
2.11.6	Organic matter	39

2.12	Arsenic remediation methods in effluent	41
2.13	Adsorption	43
2.13.1	Iron	43
2.13.2	Ferrihydrite	44
2.14	Factors affecting of As adsorption	45
2.14.1	Effect of pH	46
2.14.2	Effect of ORP	47
2.15	Method of As analysis	48
2.15.1	Atomic fluorescence spectrometry	50
2.15.2	Colorimetric techniques	50
2.15.2.1	Interference of Phosphorus and Silica	51
2.15.3	Flow Injection Analyses system for As speciation analysis using coloring reagent.	52
2.16	As(III) oxidation	53
2.17	As(V) reduction	54
2.18	Summary	54

3	A NEW APPROACH FOR ARSENIC SPECIATION IN ENVIRONMENTAL WATER COEXISTING WITH PHOSPHATE AND SILICATE VIA ON-SITE FLOW INJECTION REDOX ANALYSIS	56
3.1	Introduction	56
3.2	Materials and methods	59
3.2.1	Materials	59
3.2.2	Coloring reagent	60
3.2.3	Oxidant Reagent Selection	60

3.2.4	Reductant Reagent Selection	61
3.2.5	Water samples	61
3.2.6	Instrumentation	61
3.2.7	Preparation of calibration standard	62
3.2.8	Analytical Principle	63
3.2.9	Method validation	65
3.2.10	On-site system performance of FIA for As speciation determination in real sample	66
3.3	Results and Discussion	66
3.3.1	Oxidation	66
3.3.2	Reduction	71
3.3.3	Accuracy of FIA for measuring As based on comparison with AFS	73
3.3.4	Analytical determination	74
3.3.5	Effect of coil length on silicate	79
3.3.6	On-site system performance of FIA for As speciation determination in real sample	80
3.4	Summary	81
4	CASE STUDY OF SELINSING GOLD MINE, PAHANG - ASSESSMENT OF PROBLEMS	82
4.1	Introduction	82
4.2	Materials and methods	85
4.2.1	Sampling.	85
4.2.2	On-site water quality measurements	86
4.2.3	Analytical procedures	86

4.2.3.1	Soil characterization	86
4.2.3.2	Effluent characterization	88
4.2.4	Geotechnical Characteristics of As-bearing tailings	89
4.2.5	Soil deposit leaching tests	90
4.3	Investigation site	91
4.3.1	Study area	91
4.3.2	Sampling sites	92
4.3.3	Mining process	96
4.4	Results and Discussion	98
4.4.1	Characteristics of water from reservoir, tailing dam, and ponds.	98
4.4.1.1	Water from reservoir	98
4.4.1.2	Physical and chemical characteristics of As bearing tailing dam and ponds	101
4.4.2	Tailing dam characteristics	109
4.4.2.1	Mineralogy	110
4.4.2.2	BET and Soil characteristics	116
4.4.2.3	Particles size distribution	117
4.4.3	Leaching test	119
4.4.3.1	Effect of initial pH on As leaching	119
4.4.3.2	Effect of L/S ratio on As leaching	121
4.5	Summary	128

5 EFFICIENT REMOVAL OF ARSENIC FROM AQUEOUS SOLUTION AND SLURRY EFFLUENTS

USING NANO-IRON ION ENRICH MATERIAL (ECOMEL) SUPER ADSORBENT	130
5.1 Introduction	130
5.2 Materials and methods	133
5.2.1 Materials	133
5.2.2 Bench scale design tests	134
5.2.2.1 As(III) standard solution	134
5.2.2.2 As-bearing tailings	134
5.2.3 Adsorbents characterization	135
5.2.4 Adsorption experiments for standard As(III) and As(V)	136
5.2.5 Adsorption experiments for As-slurry	137
5.2.6 Ecomel dosage experiments	138
5.2.6.1 As(III) and As(V) remediation in effluent	138
5.2.6.2 Arsenic remediation in slurry	139
5.2.7 Analytical method	139
5.2.8 Statistical Analyses	140
5.3 Results and Discussion	140
5.3.1 Characterization of adsorbent (Ecomel)	140
5.3.2 Surface area, pore size and pore volume analysis	149
5.3.3 Effect of adsorbent dosage	156
5.3.4 Adsorption Isotherms	158
5.3.5 Effect of pH	164
5.3.6 Adsorbent Kinetics	167

		xiii
5.3.7	Mechanism of Arsenic removal	172
5.3.8	Cost analysis of Ecomel adsorbent	176
5.4	Remediation of As in slurry by Ecomel	178
5.4.1	Proposed sustainable remediation design for both As and CN^- ion	183
5.5	Summary	185
6	CONCLUSIONS AND RECOMMENDATIONS	187
6.1	Conclusions	187
6.2	Recommendations for future	191
	REFERENCES	192
	Appedices A-D	227-232

LIST OF TABLES

TABLE NO.	TITLE	PAGE
2.1	Malaysia's Production of Raw Gold 2006-2009 [52]	15
2.2	Typical As concentrations in rocks, sediments, soils in gold mining site and other surficial deposits	17
2.3	Typical As concentrations in aqueous effluent from mining site	18
2.4	Standard leaching test procedures for soil and / or waste in japan, EU, and USA [140]	40
3.1	Performance of FIA during the analysis of As	76
3.2	Detection of As speciation using FIA and comparing with AFS technique of the effluent of gold mine site–Malaysia	80
4.1	The location of sampling site	96
4.2	Measurement of in-situ parameters at five sampling sites	98
4.3	As concentration of effluent for five sampling sites using FIA and AFS systems	99
4.4	Analytical results of Arsenic and other pollutants of effluent in tailing dam at Selinsing gold mine site (average value for six samples)	104
4.5	Qualitative analysis results of the soil from tailing dam (Phase reg. detail: ICDD (PDF-2 Release 2015 RDB))	110
4.6	Analytical results of arsenic and other elements at mine tailings deposits at Selinsing gold mine site (average value for six samples) and comparing them with maximum allowable concentration in soil	111

4.7	Chemical composition of the soil from tailing dam used in the leaching experiments using XRF	112
4.8	Analytical results of some elements in effluent and As bearing tailing at Selinsing gold mine site	114
4.9	Soil characteristics	116
5.1	Chemical composition of Ecomel using XRF	141
5.2	Ecomel characteristics (BET and pH_{pzc}).	151
5.3	Estimated isotherm parameters for arsenic (As) speciation/adsorption	161
5.4	Comparison of As(III) or As(V) removal	163
5.5	Estimated kinetic model parameters for arsenic speciation adsorption*	170
5.6	Intra-particle diffusion model	171
5.7	Cost analysis for arsenic (As) removal	177
5.8	Efficiency of remediation of As-soil using Ecomel at 0.01 Ecomel/Soil, stirring speeds: 200 rpm; L/S ratio = 10 and initial pH 2.5	183

LIST OF FIGURES

FIGURE NO.	TITLE	PAGE
1.1	Experimental flow	10
3.1	Diagram of FIA for As(V) analysis. P1 & P2: double plunger pump, PW: pure water 0.6 mL/min, CR: coloring reagent 0.2 mL/min, AS: automated sampler, RC: reaction coil, I.D. 0.5 mm, 5.6 m at 70 ^o C, CC: cooling coil, I.D. 0.5 mm, 1.0 m at 5 ^o C, BPC: back pressure coil, I.D. 0.25 mm, 5.0 m at room temperature, CO: column oven at 70 ^o C, SPC: spectrophotometer at 700 nm, WW: wastewater	62
3.2	Schematic depiction for the analytical principle of As speciation by off-line oxidation and reduction process	64
3.3	Schematic depiction for the analytical principle to analyze dissolved inorganic As(III) and As(V) by differentiation of oxidized, reduced and untreated samples	65
3.4	Validation for off-line As(III) oxidation rate by a) KMnO ₄ and b) NaClO with As(V) control standard solution with 0.5 M-NO ₃ ⁻ ion	67
3.5	Validation for off-line As(III) oxidation rate by a) KMnO ₄ and b) NaClO with As(V) control standard solution (pure As(V) solution)	68

3.6	The relationship between peak area and As(V) concentration with and without NO_3^- ion at (a) 0-0.05 mM and (b) 0-0.5 mM of As(V)	70
3.7	The comparison between sample with NO_3^- ion and without NO_3^- ion	71
3.8	Validation of FIA for As(V) reduction rate: a) River water, b) UPW using $\text{Na}_2\text{S}_2\text{O}_3 \cdot 5\text{H}_2\text{O}$	72
3.9	Correlation between the total As [Oxidation As[III & V] and As(V)[Untreated] concentrations in river samples determined by the present As oxidation-reduction sequential analysis using FIA and AFS	74
3.10	Standard calibration curve for a) As(III), b) As(V) and c) PO_4^{3-} and SiO_3^{2-}	75
3.11	FIA signal peaks for As(V)	77
3.12	Repetitive FIA signal peaks of a) 10 $\mu\text{g/L}$ (0.13 μM) for each of As(III), As(V), PO_4^{3-} and SiO_3^{2-} , and b) 0.01 mM of As(V) with 0.3 mM of each of PO_4^{3-} and SiO_3^{2-}	78
3.13	FIA peak area for silicate with different length of the reaction coil	79
4.1	Map of (a) Selinsing gold mining site in Pahang, Malaysia and (b) focused sampling point	92
4.2	Sceneries of sampling sites. a: Reservoir, b: Tailing dam, c: 1 st tailing pond, d: 3 rd tailing pond and e: 4 th tailing pond	93
4.3	The schematic depiction for sampling sites	94
4.4	The whole landscape of tailing ponds and sampling sites 1: Reservoir, 2: Tailing dam, 3: 1 st tailing pond, 4: 3 rd tailing pond and 5: 4 th tailing pond	95
4.5	The schematic depiction of the mining process, and the treatment in Selinsing gold mining site	97

4.6	CN ⁻ ion concentration in effluent from tailing dam (dilution factor = 1000)	103
4.7	The concentration of As(III) and As(V) in reservoir, tailing dam and tailing ponds measured using FIA	106
4.8	The total As concentration in reservoir, tailing dam and tailing ponds measured using AFS	106
4.9	a) FE-SEM micrograph and b) EDX analyses of Gold mine tailing	115
4.10	Particle size distribution of soil tailing from Selensing gold mining	117
4.11	USDA soil textural triangle illustrates the classification of soil type (x, y, z): (sand 0.05-2.0 mm, clay <0.002 mm, silt 0.002-0.05 mm)	118
4.12	Change of total As and pH with time in the batch experiments with, solid:liquid ratio = 1:5 and initial pH a) 2.5, b) 3.7, and c) 11.5	121
4.13	Change of total As and pH with time in the batch experiments with, initial pH 2.5 and L/S ratio = a) 5, b) 10, and c) 20	124
4.14	Change of total As and pH with time in the batch experiments with, initial pH 3.7 and L/S ratio = a) 5, b) 10, and c) 20	126
4.15	Change of total As and pH with time in the batch experiments with, initial pH 11.5 and L/S ratio = a) 5, b) 10, and c) 20	128
5.1	TG and DTG of complex formed by Ecomel with a) As(III) and b) As(V)	143
5.2	FE-SEM micrographs of a) Ecomel b) complex generated from reaction between Ecomel and As(III) at initial pH2.5 c) complex generated from reaction between Ecomel and As(V) at initial pH2.5	144
5.3	EDX analyses of a) Ecomel b) complex generated from reaction between Ecomel and As(III) at initial pH 2.5 c) complex	

	generated from reaction between Ecomel and As(V) at initial pH 2.5	146
5.4	Ecomel with 100 mg/L of As(III)/As(V) before and after forming As(III)/As(V)- Ecomel complex at initial 1 mg Ecomel/L; initial pH 2.5; stirring speeds: 200 rpm; temperature $24 \pm 1^\circ\text{C}$; and contact time 24 h	147
5.5	X-ray Diffraction of Ecomel	148
5.6	X-ray Diffraction of a) Ecomel reaction with 100 mg-As(III)/L initial pH2 b) Ecomel reaction with 100 mg-As(V)/L at initial pH 2.5	150
5.7	a) N_2 adsorption/desorption isotherms b) Pore size distribution and pore volume of raw Ecomel	152
5.8	a) N_2 adsorption/desorption isotherms b) Pore size distribution and pore volume of Ecomel at initial pH 2.5 using HCl with 100 mg As(III) /L	154
5.9	a) N_2 adsorption/desorption isotherms b) Pore size distribution and pore volume of Ecomel at initial pH 2.5 using HCl with 100 mg As(V) /L	155
5.10	Ecomel Dosage effect on the removal of As(III) and As(V) at initial [As]:1 mg/L; initial pH 2.5; stirring speeds: 200 rpm; temperature: $24 \pm 1^\circ\text{C}$; equilibration time: 2 h	156
5.11	Bench scale design for the As(III) removal at initial [As(III)]: 100 mg/L; 1 g Ecomel/L; initial pH 2.5; stirring speeds: 200 rpm; temperature: $24 \pm 1^\circ\text{C}$; contact time: 24 h	157
5.12	Langmuir, Freundlich, Langmuir – Freundlich model isotherm plots for a) As(III) and b) As(V)	160
5.13	a) Effect of pH, b) Effect of initial pH on final pH, and c) pH_{pzc} of Ecomel	166
5.14	Pseudo first order, Pseudo second order, and Elovich model kinetics plots for a) As(III) and b) As(V)	169

- 5.15 Effect of Ecomel/Soil ratio on the removal of As from slurry. at initial : pH: 2.5; stirring speeds: 200 rpm; temperature: $24 \pm 1^{\circ}\text{C}$; L/S ratio: 10 and contact time: 24 h 179
- 5.16 Bench scale design of As-Soil leachout concentration with and without remediation using Ecomel a) reaction time b) settling time, under the following conditions, 0.01 Ecomel /Soil, stirring speeds: 200 rpm; L/S ratio = 10 and initial pH 2.5 181
- 5.17 Trend of As-Soil leachout concentration with and without remediation using Ecomel under the following conditions, 0.02 Ecomel/Soil, stirring speeds: 200 rpm; L/S ratio = 10 and initial pH 2.5 182
- 5.18 Schematic of proposed reactor for remediating As-slurry of Selinsing gold mining site in Pahang 184

LIST OF ABBREVIATIONS

AAS	-	Atomic absorption spectroscopy
ABS	-	Absorbance
AdSV	-	Adsorptive stripping voltammetry
AES	-	Atomic emission spectroscopy
AFS	-	Atomic fluorescence spectrometry
AFU	-	Arbitrary fluorescence unit
Al	-	Aluminum
AMD	-	Acid mine drainage
ARE	-	Average relative error
As	-	Arsenic
AS	-	Automated sampler
As(III)	-	Arsenite
As(T)	-	Total arsenic
As(V)	-	Arsenate
As ₂ S ₃	-	Orpiment
AsS	-	Realgar
Au	-	Gold
B	-	Boron
BET	-	Brunauer–Emmett–Teller (BET)
BOFS	-	Basic oxygen furnace slag
BPC	-	Back pressure coil

C	-	Carbon
Ca	-	Calcium
CC	-	Cooling coil
Cd	-	Cadmium
CE	-	Capillary electrophoresis
ClO ₂	-	Chlorine dioxide
CN ⁻	-	Cyanide
Co	-	Cobalt
CO	-	Column oven
COPD	-	Chronic Obstructive Pulmonary Disease
Cr	-	Chromium
CR	-	Coloring reagent
CS-P-NZVI	-	Nanoscale zero-valent iron on pumice
Cu	-	Copper
DO	-	Dissolved oxygen
DOE	-	Department of Environment
DTA	-	Differential thermal analysis
EA	-	Elemental Analyzer
EC	-	Electrical conductivity
EDX	-	Energy-dispersive X-ray spectroscopy
Eh	-	Redox potential
e-MWCNT/Fe ²⁺	-	Iron(II) oxide coated ethylenediamine functionalized multiwall carbon nanotubes
e-MWCNT/Fe ³⁺	-	Iron(III) oxide coated ethylenediamine functionalized multiwall carbon nanotubes
ER/DETA/FO/FD	-	Cross-linked macroporous polymer impregnated with hydrous iron oxide:
ET-AAS	-	Electrothermal atomic absorption spectrometry
Fe	-	Iron

Fe ⁰	-	Zero Iron
Fe ²⁺	-	Ferrous ions
Fe ³⁺	-	Ferric ions
FeAsS	-	Arsenopyrite
FeS	-	Pyrite
FESEM	-	Field Emission Scanning Electron Microscopy
FIA	-	Flow injection analysis
FIB-SEM	-	Focus Ion Beam –Scanning Electron Microscope
GC-MS	-	Gas chromatography-mass spectrometry
GFAA	-	Graphite furnace atomic absorption
H ₂ O ₂	-	Hydrogen peroxide
H ₄ SiO ₄	-	Silicic acid
HCN	-	Hydrogen cyanide
HG-AAS	-	Hydride generation coupled with atomic absorption spectrometry
HG-AFS	-	Hydride generation coupled with atomic fluorescence spectrometry
HIO-alginate beads	-	Hydrous iron oxide-impregnated alginate beads
HPLC-AFS	-	High performance liquid chromatography with atomic fluorescence spectrometry
HPLC-ICP-MS	-	High performance liquid chromatography-inductively coupled plasma mass spectrometry
IARC	-	International Agency for Research on Cancer
ICP-AES	-	Inductively coupled plasma emission spectrophotometry
ICP-MS	-	Inductively coupled plasma mass spectroscopy
ICPOES	-	Inductively Coupled plasma optical Emission spectrophotometer
KIO ₃	-	Potassium iodate
KMnO ₄	-	Potassium permanganate

L/S	-	Liquid to solid ratio
LC-MS-MS	-	High performance liquid chromatography-tandem mass spectrometry
LDL	-	lower detection limit
LOI	-	Loss on ignition
Mn	-	Manganese
Mo	-	Molybdenum ion
N	-	Nitrogen
N**	-	Not measured
Na ₂ S ₂ O ₃	-	Sodium thiosulphate
NAA	-	Electrochemical methods, neutron activation analysis
NaClO	-	Sodium hypochlorite
NH ₂ Cl	-	Monochloramine
NIIEM	-	Nano-iron ion Enrich materials
-NO ₃ ⁻	-	Without NO ₃ ⁻
NSD	-	Normalized standard deviation
NST	-	New straits time
NZVI	-	Nano-zero-valent iron
nZVI in nZVI/BC composite	-	Nano-zero-valent iron in biochar supported zero-valent iron nanocomposite
nZVI/BC	-	Biochar supported zero-valent iron nanocomposite
NZVI-AC	-	Nano zero-valent iron on activated carbon
OM	-	Organic matter
ORP	-	Oxidation-Reduction Potential
PO ₄ ³⁻	-	phosphate
P1 & P2	-	Double plunger pump
pb	-	Lead
pH _f	-	Final pH value
pH _i	-	Initial pH value

pH _{pzc}	-	Point of zero charge
PHTEs	-	Potential hazardous toxic elements
PP	-	polypropylene
PW	-	Pure water 0.6 mL/min
PZD	-	Particle size distribution
R ²	-	Coefficient of determination
RC	-	Reaction coil
Redox	-	Reduction/Oxidation
RMSE	-	Residual root mean square error
RSD	-	Relative standard deviation
S	-	Sulfur
S/N	-	Signal-to-noise ratio
SDS	-	Sodium dodecyl sulfate
SE	-	Standard error
SiO ₃ ²⁻	-	Silicate
SPC	-	Spectrophotometer
STD	-	Standard deviation
Temp.	-	Temperature
TG	-	Thermogravimetry
TGA	-	Thermo-gravimetry analysis
Unimpregnated HIOs	-	Unimpregnated hydrous iron oxide
UPW	-	Ultra-pure water
WHO	-	World health organization
WTRs	-	Water treatment residuals
WW	-	Wastewater
XRD	-	X-ray diffraction
XRF	-	X-ray fluorescence
Zn	-	Zinc
ZVI	-	Zero-valent iron

LIST OF SYMBOLS

C_0	-	Liquid phase concentrations of As(III)/As(V) at initial time zero
C_e	-	Equilibrium concentration in the solution (mg/L)
C_t	-	Liquid phase concentrations of As(III)/As(V) at time t
E%	-	Percentage removal of arsenic
K_1	-	Pseudo-first order sorption rate constant (h^{-1})
K_2	-	Pseudo-second order constant (g/mg h)
K_f	-	Freundlich constant (L/g)
K_{id}	-	Intra-particle diffusion rate constant ($\text{mg/g h}^{0.5}$)
K_{L1}	-	Langmuir constant (mg/g)
K_{L2}	-	Langmuir-Freundlich constant ($\text{L/mg}^{1/n}$)
n	-	Freundlich adsorption equilibrium constant (dimensionless)
q	-	Quantity of adsorbed As(III)/As(V) at equilibrium settings (mg/g)
q_e	-	Quantity of As(III)/As(V) adsorbed on the surface of Ecomel at equilibrium (mg/g)
q_m	-	Maximum adsorption capacity (mg/g)
q_t	-	Amount of As(III)/As(V) adsorbed by Ecomel, at time, t (mg/g)
R_{L1}	-	Dimensionless equilibrium constant for Langmuir model
R_{L2}	-	Dimensionless equilibrium constant for Langmuir-Freundlich models
t	-	Time (h)
$t_{1/2}$	-	Time for half equilibrium (min)
V	-	Volume of the As-contaminated solution (L)

W	-	Mass of Ecomel used for As(III)/As(V) adsorption (mg)
α	-	Parameters of the Elovich rate equation (mg/g)
β	-	Parameters of the Elovich rate equation (g/mg)
χ^2	-	Chi-square test
n_1		Langmuir-Freundlich constant (dimensionless)

LIST OF APPENDICES

APPENDIX	TITLE	PAGE
A1	Calibration curve for As standard with different concentration detected using FIA system a) high concentration, b) medium concentration, c) low concentration	227-228
B1	Calibration Curve For As Standard With Different Concentrations Detected Using AFS Gain 1	229
B2	Calibration Curve For As Standard With Different Concentrations Detected Using AFS Gain10	230
C	Ecomel Manufacture	231
D	Publication	232

CHAPTER 1

INTRODUCTION

1.1 Background of the study

Gold mining sites are mostly associated with positive economic development of any country; however extraction of gold could also generate negative impacts on the ecosystem and human health. Mining and ore-processing activities generate huge quantities of tailing that could have severe and widespread impacts on ecosystem over a long period of time [1]. Gold mining is considered as a significant source of environmental pollution by trace elements especially high arsenic (As) concentration arising from ore excavation, transportation, mineral processing, and disposal of huge quantities of waste around the mining site [2,3]. Recently, ecosystem deterioration is regarded as an important problem, and both active and expected mining exploitations are legally compelled to minimize the contamination impact and to restore the exploited sites after their abandonment. The serious environmental problem and public-health concerns associated with gold mining activities is As, which is released from the oxidation of sulfide minerals such as pyrite (FeS_2) and arsenopyrite (FeAsS) [4].

Gold is often occurring as encapsulated in other minerals such as FeS_2 , quartz, FeAsS , etc. The particles of gold are generally extracted via ore leaching using cyanide (CN^-) ion solution. Throughout the cyanidation process, the metal ore

reacts with CN^- ion solution under highly alkaline conditions ($\text{pH} > 11$) for duration of up to 36 h, thus, releasing high As concentration. The As-bearing tailings from sulfide ore mines are usually chemically reactive because of their content of Fe-sulfide minerals, like as FeS_2 and FeAsS . The weathering of Fe-sulfide minerals is a strong acidifying process, and regarded to be the major reason for acid mine drainage (AMD) [5,6]. Arsenic can't form stable complexes together with CN^- ion such as gold (Au), however, it is well-known that As are very soluble at the highly alkaline conditions observed for the cyanidation process. Thus, the water used in this process is highly contaminated with dissolved As and CN^- ion.

Consequently, the huge amount of water used in this process must be remediated to remove dissolved As prior to its discharge into tailings dam. The management of As-bearing tailings generated through gold extraction processes usually concentrates on the stability of secondary As-phases mainly because As-sulfides such as, realgar (AsS), orpiment (As_2S_3), FeAsS are assumed to be oxidized during gold extraction. However, As could be existing in two major redox states, namely; arsenite As(III) and arsenate As(V) in aqueous effluent, both having differences in solubility and mobility according to the oxidation-reduction potential (ORP) and pH conditions in the ecosystem [7]. The major secondary As minerals existing in tailing dam after cyanidation process are supposed to be As(V), due to the addition of high hydrogen peroxide concentration during the gold extraction process. At oxidizing conditions, As(V) generally forms complexes with metals such as Fe, Al and Ca [7,8]. However, the stability and formation of these As(V)-metal complexes is based on several important factors, such as, pH, Fe/As or Al/As ratio, redox potential conditions, availability of O_2 and CO_2 as well as the presence of anions, such as SO_4^{2-} ion, phosphorus (PO_4^{3-}) and silicate (SiO_3^{2-}) that compete with As(V) for sorption surfaces [9,10]. Fe and Al-As(V) are the dominating phases in acidic condition during gold mining and are less soluble than Ca-As(V), such as schwertmannite or ferrihydrite. Gold mine tailings slurries after gold extraction containing As-enriched sulfide ores are usually remediated by lime and $\text{Fe}_2(\text{SO})_3$ to

enhance the formation of stable Fe-As-precipitates and increase the solution to prevent the formation of acid mine drainage. Furthermore, As-bearing tailings slurries are usually pre-remediated with H_2O_2 and O_2 in order to enhance the oxidation of As(III) complexes into less mobile As(V) species [6,11].

However, in recent years the world have witnessed the emergent cases of As contamination in water and aqueous effluent as a big global environmental problem, associated the rapid growth of modern industries and mining process [12]. In many cases, the As concentration in aqueous samples from gold mining site are determined within the range of 10-100 mg/L at acidic settings, especially in FeAsS-rich areas [4,13]. Whilst in soil, As was detected ranging from 2,000-390,970 mg/kg in acid mine drainages [14,15]. These highly elevated concentration of As should be effectively remediated using highly efficient adsorbent to reduce its environmental impact.

Due to the high toxicity of As, the world health organization (WHO) announced the maximum permissible limit of As in drinking water to be 10 $\mu\text{g/L}$ [16]. And also U.S. Environmental Protection Agency (EPA) adopted a lower standard limit from 50 to 10 $\mu\text{g/L}$ of As concentration in drinking water [17]. Therefore, there is great concern for As contaminations for human health, plants, soil, and water to prevent the ecosystem from deteriorations caused by the high toxicity of As presence.

These increasing trend of growth of interest in As remediation is in agreement with the result of the study performed in 2007 where an estimate of more than 137 million people in over 70 countries, including Argentina, Bangladesh, China, Hungary, India, Japan, Malaysia, Mexico, Romania, United States and Vietnam are exposed As toxicity via water sources [18–20]. Thus, the great harm to human health for As was affirmed and for this reason, As was categorized as class

one carcinogen by the International Agency for Research on Cancer (IARC) [21]. Additionally, more than 200,000 people suffered death due to As poisoning every year after being subject to a variety of cancerous growths such liver, lungs, bladder, kidney and prostate cancers [22,23]. Similarly, As has also been related to non-cancerous multi-systemic health problems such as renal failure, diabetes mellitus, dermatological diseases, respiratory disorder, cardiovascular disease, hypertension and so forth [18,24]. Unfortunately, As contamination removal is not easy as well, as it can easily pass through food chains and drinking water into the body. Moreover, As exhibits a strong tendency for accumulation in the human body and hence threatens prolong health risks [12].

Due to gold mining processes, high As concentrations are generated for both tailing and aqueous effluents. These highly elevated As must be remediated before discharge into the water bodies. The remediation of As-bearing tailings or As contaminated area is a considerable challenge to the scientific community, especially at mining sites that have dramatically impacted the water resources and soils in the environment [25]. In other hand, As removal from aqueous effluent can be a very costly process. The costs for As removal could exceed the benefited, needing very careful selection of the technology to be used. First of all, the selection must consider the water characteristics to be remediated [26]. To date, there are various technologies applied to remediate As from natural waters and industrial effluents including coagulation-precipitation, oxidation, coagulation, ion-exchange, membrane separation, adsorption, reverse osmosis and electro-kinetic methods. However, these technologies have several drawbacks such as high waste remediation and operating costs, high consumption of reagents as well as large amounts of toxic sludge formation. Among these processes, adsorption technique is considered as one of the most promising technologies for mitigating As in aqueous effluent scenario, due to its low cost, operational simplicity, high concentration efficiency, and environmental friendliness [27,28]. Consequently, it is vital to properly remediate this aqueous

effluent utilizing techniques that involve, for example, adsorption in a complementary way or as alternatives for conventional techniques.

Nowadays, As remediation from aqueous effluent has become a major active research focus. Numerous approaches and studies are applied to obtain abundant, porous structure and chemically-stable inexpensive adsorbent with high surface area [18]. These includes adsorbents such as iron oxide, iron oxyhydroxides, aluminum oxide/oxyhydroxides, activated carbon, activated alumina, rare earth oxides, fly ash, manganese green sand, iron-containing mesoporous carbon, natural material, both Fe_2O_3 and Fe_3O_4 nanomaterials, red mud, zeolite, zero valent iron nanoparticles, granulated ferric oxide, manganese oxides, transition metal sulfides, magnetic wheat straw, titanium dioxide, iron-oxide@carbon and cellulose@ Fe_2O_3 composites, etc. The effectiveness of adsorption technique selection is mostly dependent on the adsorbent's removal efficiency and their economic value. Due to the tremendous adsorption potential, magnetic properties, and strong affinity of iron toward As, iron oxide compounds or embedded iron have high sorption affinity toward As species, therefore, iron based materials are commonly used in various methods for the remediation of As contaminated water and soil [17,20,28].

1.2 Problem statement

Nowadays, many instruments used for measuring As includes hydride generation coupled with atomic fluorescence spectrometry (HG-AFS), hydride generation coupled with atomic absorption spectrometry (HG-AAS), electrothermal absorption spectrometry (ET-AAS), inductively coupled plasma with atomic emission spectrometry/inductively coupled plasma with mass spectrometry (ICP-AES; ICP-MS), high performance liquid chromatography with atomic fluorescence

spectrometry (HPLC-AFS), high performance liquid chromatography with inductively coupled plasma-mass spectrometry (HPLC-ICP-MS) etc. [29–31]. These analytical techniques cannot be considered as portable instruments for on-site applications as they required complicated sample pre-treatment process, and skilled personnel [32]. As(III) dominates the gold mining site due to FeS_2 or FeAsS oxidation. As(III) can commonly get oxidized and transformed to As(V) during the sampling to laboratory. Therefore, the result of the test is often biased and does not express the exact As speciation of the samples [33]. Consequently, there is a need to develop a simple, sensitive and accurate method for on-site analysis of As speciation in environmental water sources.

In the gold mining site, great amount of water is used for gold extraction. For example, in Selinsing gold mining site, about $6000 \text{ m}^3/\text{d}$ of effluent are discharged into the tailing dam. This aqueous effluent contained extremely high concentration of As and CN^- ion as well as high alkaline pH and oxygen concentration. Consequently, the water resources near the gold mine are contaminated. Thus, exposing the inhabitants to severe health risks such as arsenicosis, cancers, diabetes, hearing loss, hematological disorders, keratosis, neurobehavioral disorders, and pulmonary fibrosis [30,34,35].

In our visit to Selinsing gold mine site, the site engineer informed us that they had problem in the management of the aqueous effluent (and not the sediment which according to him is stable). We were informed that the As concentration in first tailing pond is generally between 10-17 mg/L which was extremely high in comparison with the effluent standard for As in Malaysia (0.1 mg/L).

1.3 Research objective

The main objectives of this research are as follows:

- a) To develop a new technical method for measuring As speciation.
- b) To determine the As concentration in soil and aqueous effluent from Selinsing gold mining site (Malaysia), and identify the problems associated with the elevated As concentration in its discharge effluent.
- c) To investigate the applicability of Ecomel (Iron ion enrich materials) for removal of As(III) and As(V) in standard aqueous solutions and treating the slurry from gold mine site using batch method.

1.4 Scope of research

For measuring As speciation, redox–FIA system was developed. The remediation of As in liquid effluents was carried out using Ecomel under acidic condition without the addition of oxidants. The experimental flow of the research is shown in Figure 1.1. Thus, this research thesis was divided into three phases as follows:

Phase I: Developing new technical method for As speciation measurement in water samples co-existing with phosphorus (PO_4^{3-}) and silicate (SiO_3^{2-}) using FIA.

The first phase focused on developing new method for measuring As speciation and includes:

- a) Investigation of the simple As speciation analysis for As (III), As (V) and in PO_4^{3-} and SiO_3^{2-} contaminated environmental sample.
- b) Optimization of the oxidant concentration that can complete oxidation of As(III).
- c) Optimization of the reductant concentration that can complete reduction of As(V).
- d) Investigating techniques to eliminate the effect of PO_4^{3-} and SiO_3^{2-} as interference to As analysis by molybdenum blue.
- e) Testing the accuracy of Flow Injection Analysis (FIA) for measuring As based on comparison with atomic fluorescence spectrometry (AFS).
- f) The system performance of FIA for As speciation determination in real sample.

Phase II: Sampling field works at Selinsing gold mining site in Pahang Malaysia

The second phase focused on obtaining the environmental assessment for the mining site as the background of the study. Activities done during this phase includes;

- a) Case study of Selinsing gold mining site problems in aqueous effluent and tailings.

- b) All aqueous effluent and As-bearing tailings were collected from Selinsing gold mining site at 17-7-2015 and 3-1-2016.
- c) Characterizations of soil from tailing dam using XRD, XRF, pH, pH_{pzc} , FESEM-EDX, nitrogen-adsorption analysis, ICPOES, AFS.
- d) Onsite water quality measurements such as, pH, DO, ORP, EC, and temperature with ICPOES, AFS, and As speciation by FIA.
- e) Studying the characterizations of soil such as pH, particle size distribution, classification of soil and Loss on ignition (LOI).
- f) Studying the effect of pH and L/S on As leaching.

Phase III: Batch experimental set using Ecomel adsorbent, a new application for As remediation in aqueous solution and slurry waste.

This phase focused on the preparations, testing and characterizations of Ecomel before and after adsorption. The processes include:

- a) Characterizations using XRD, FESEM-EDX, pH, pH_{pzc} , TGA-DTA, nitrogen-adsorption analysis.
- b) Adsorbent kinetics and isotherms.
- c) Study the mechanism of As removal.
- d) Cost analysis study for Ecomel.
- e) Studying the optimum dosage of Ecomel for remediation As-bearing

tailings.

- f) Studying the optimum dosage of Ecomel for remediation As(III) and As(V) from water.
- g) Studying the optimum dosage of Ecomel for remediation As from slurry.
- h) Model suggestion for slurry remediation.




Figure 1.1 Experimental flow

1.5 Significance of research

The significance of this research is to resolve an actual problem in Malaysia and many other developing and developed countries resulting from high As concentration. This explores the viability of a proposed model for As remediation in aqueous effluent in gold mining site.

The present research focuses on Selinsing gold mining site, Pahang–Malaysia. Field studies were performed to identify the problem of high As leaching from tailing. Adsorption under acidic conditions in laboratory batch scale was used for the remediation of queues effluent. Generally, the present study provides a complete view on the cause of As leaching in aqueous solution and suggest the reliable system to remediate As and overcome any potential problems associated with As leaching.

Even though the aqueous effluents were remediated by adding hydrogen peroxide (H_2O_2) for the oxidation of As(III) to As(V) and as well integrating high concentration of iron impregnated sand to adsorb As(V), As concentration in treated effluents were still extremely high and does not meet Malaysia's As effluent discharge standards (0.1 mg/L). Thus, this study introduces a new application of super-adsorbent namely Ecomel with low-cost and high adsorption capacity to adsorb As especially As(III) without any further pre-remediation. Ecomel can remediate high As concentration in both aqueous effluent and slurry.

The success of this research will consider Malaysia as a contributing nation for sustainable mining policy and as well create higher reputation to the Malaysian's research activity and applied novelty of low-cost technology for on-site remediation in As-bearing tailings. This enables Malaysia to have within their domain, a recent

technology for remediating As from gold mining sediment and protect its water resources from As contaminations. Therefore, this study provides a promising alternative technique for the remediation of As-bearing tailings in view of its low-cost and long term application as a sustainable mitigation technology.

Similarly, a simple, selective and stable on-site Flow Injection Analysis (FIA) redox analytical method was developed to determine As speciation in water samples constituted with phosphate (P) and silicate (Si). The developed method senses the As(V), P and Si by reacting with ammonium molybdenum, which forms a molybdenum blue complex under acidic condition. This study was carried out to define an alternative method to remediate As contamination in aqueous effluent. As mobility in As-bearing tailings can be controlled by two important processes: (i) adsorption and desorption processes and (ii) solid-phase precipitation and dissolution.

1.6 Limitation of research

Briefly, the limitation of this research can be summarized by the following:

- a) Difficulty to get information about the industrial process, concentration on the influent/effluent and adsorbent material that was used for As remediation in aqueous effluent in Selinsing gold mining site.
- b) Limitation of sampling frequency due to stringent regulation from the management and the location of sampling site is far from UTM.

- c) Difficulty to apply the research on-site for remediating both aqueous and slurry effluents.

REFERENCES

- [1] M. M. Chico, G. Damian, D. Stumbea, N. Buzgar, T. Ungureanu, V. Nica, and G. Iepure, "Mineralogy and geochemistry of the tailings pond from straja valley (suceava county, romania). factors affecting the mobility of the elements on the surface of the waste deposit," *Carpathian Journal of Earth and Environmental Sciences*, vol. 11, no. 1, pp. 265–280, 2016.
- [2] B. J. B. Nyarko, Y. Serfor-Armah, J. Holzbecher, E. K. Osae, E. H. K. Akaho, D. Carboo, and A. Chatt, "Epithermal instrumental neutron activation analysis for the determination of gold and arsenic in Ghanaian gold tailings using conventional and anti-coincidence counting," *Journal of Radioanalytical and Nuclear Chemistry*, vol. 256, no. 2, pp. 253–257, 2003.
- [3] F. B. Ono, L. R. G. Guilherme, L. A. Mendes, and G. S. Carvalho, "Replication of an ivg protocol to estimate bioaccessible arsenic in materials from a gold mining area in Brazil," *Revista Brasileira de Ciência do Solo*, vol. 36, no. 4, pp. 1355–1360, 2012.
- [4] M. A. Acheampong, K. Paksirajan, and P. N. L. Lens, "Assessment of the effluent quality from a gold mining industry in Ghana," *Environmental Science and Pollution Research*, vol. 20, no. 6, pp. 3799–3811, 2013.
- [5] N. E. Nieva, L. Borgnino, F. Locati, and M. G. García, "Mineralogical control on arsenic release during sediment-water interaction in abandoned mine wastes from the Argentina Puna," *Science of the Total Environment*, vol. 550, pp. 1141–1151, 2016.
- [6] A. Suttigarn, Y. Wang, and M. Asce, "Arsenite Oxidation by *Alcaligenes faecalis* Strain O1201," *Journal of environmental engineering*, vol. 131, no. 9, pp. 1293–1301, 2006.

- [7] C. Roussel, C. Néel, and H. Bril, "Minerals controlling arsenic and lead solubility in an abandoned gold mine tailings," *Science of the Total Environment*, vol. 263, no. 1–3, pp. 209–219, 2000.
- [8] S. Kim, W. Kim, and K. Kim, "Evaluation of electrokinetic remediation of arsenic-contaminated soils," *Environmental Geochemistry and Health*, vol. 27, pp. 443–453, 2005.
- [9] S. Hyun, J. Kim, D. Y. Kim, and D. H. Moon, "Effect of seepage conditions on chemical attenuation of arsenic by soils across an abandoned mine site," *Chemosphere*, vol. 87, no. 6, pp. 602–607, 2012.
- [10] G. Dermont, M. Bergeron, G. Mercier, and M. Richer-Lafèche, "Metal-Contaminated Soils: Remediation Practices and Treatment Technologies," *Practice Periodical of Hazardous, Toxic, and Radioactive Waste Management*, vol. 12, no. 3, pp. 188–209, 2008.
- [11] R. Hamberg, G. Bark, C. Maurice, and L. Alakangas, "Release of arsenic from cyanidation tailings," *Minerals Engineering*, vol. 93, pp. 57–64, 2016.
- [12] S. Lin, D. Lu, and Z. Liu, "Removal of arsenic contaminants with magnetic γ -Fe₂O₃ nanoparticles," *Chemical Engineering Journal*, vol. 211–212, pp. 46–52, 2012.
- [13] F. Bodéan, P. Baranger, P. Piantone, A. Lassin, M. Azaroual, E. Gaucher, and G. Braibant, "Arsenic behaviour in gold-ore mill tailings, Massif Central, France: Hydrogeochemical study and investigation of in situ redox signatures," *Applied Geochemistry*, vol. 19, no. 11, pp. 1785–1800, 2004.
- [14] C. Roussel, H. Brilatb, and A. Fernandez, "Evolution of sulphides-rich mine tailings immobilization of As and Fe," *Comptes Rendus de l'Académie des Sciences-Series IIA-Earth and Planetary Science*, vol. 329, no. 11, pp. 787–794, 1999.
- [15] D. Mains and D. Craw, "Composition and mineralogy of historic gold processing residues, east Otago, New Zealand," *New Zealand Journal of Geology and Geophysics*, vol. 48, no. 4, pp. 641–647, 2005.

- [16] T. Basu, K. Gupta, and U. C. Ghosh, "Performances of As(V) adsorption of calcined (250 °C) synthetic iron(III)–aluminum(III) mixed oxide in the presence of some groundwater occurring ions," *Chemical Engineering Journal*, vol. 183, pp. 303–314, 2012.
- [17] J. Cantu, L. E. Gonzalez, J. Goodship, M. Contreras, M. Joseph, C. Garza, T. M. Eubanks, and J. G. Parsons, "Removal of arsenic from water using synthetic Fe₇S₈ nanoparticles," *Chemical Engineering Journal*, vol. 290, pp. 428–437, 2016.
- [18] M. Marinovic, K. Taleb, J. Markovski, M. Milosavljevic, M. Ristic, and A. Marinkovic, "Efficient arsenic removal by cross-linked macroporous polymer impregnated with hydrous iron oxide: Material performance," *Chemical Engineering Journal*, vol. 279, pp. 66–78, 2015.
- [19] L. Hao, T. Zheng, J. Jiang, G. Zhang, and P. Wang, "Removal of As(III) and As(V) from water using iron doped amino functionalized sawdust: Characterization, adsorptive performance and UF membrane separation," *Chemical Engineering Journal*, vol. 292, pp. 163–173, 2016.
- [20] T. Mishra and D. K. Mahato, "A comparative study on enhanced arsenic (V) and arsenic (III) removal by iron oxide and manganese oxide pillared clays from ground water," *Journal of Environmental Chemical Engineering*, vol. 4, no. 1, pp. 1224–1230, 2016.
- [21] Z. Zhou, Y. Liu, S. Liu, H. Liu, G. Zeng, X. Tan, C. Yang, Y. Ding, Z. Yan, and X. Cai, "Sorption performance and mechanisms of arsenic(V) removal by magnetic gelatin-modified biochar," *Chemical Engineering Journal*, vol. 314, pp. 223–231, 2017.
- [22] K. R. Kim, B. T. Lee, and K. W. Kim, "Arsenic stabilization in mine tailings using nano-sized magnetite and zero valent iron with the enhancement of mobility by surface coating," *Journal of Geochemical Exploration*, vol. 113, pp. 124–129, 2012.
- [23] T. Weerasiri, W. Wirojanagud, and T. Srisatit, "Localized Profile of Arsenic

- in Soil and Water in the Area Around Gold Mine,” *Current World Environment*, vol. 8, no. 2, pp. 231–240, 2013.
- [24] S. Lin, D. Lu, and Z. Liu, “Removal of arsenic contaminants with magnetic γ -Fe₂O₃ nanoparticles,” *Chemical Engineering Journal*, vol. 211–212, pp. 46–52, 2012.
- [25] J. V. Dos Santos, W. De Melo Rangel, A. Azarias Guimarães, P. M. Duque Jaramillo, M. Rufini, L. M. Marra, M. Varón López, M. A. Pereira Da Silva, C. R. Fonsêca Sousa Soares, and F. M. De Souza Moreira, “Soil biological attributes in arsenic-contaminated gold mining sites after revegetation,” *Ecotoxicology*, vol. 22, no. 10, pp. 1526–1537, 2013.
- [26] C. N. Kesavachandran, “Reviews of Environmental Contamination and Toxicology Vol 200,” *Toxicology*, vol. 200, no. 80, pp. 87–103, 2010.
- [27] J. Wang, W. Xu, L. Chen, X. Huang, and J. Liu, “Preparation and evaluation of magnetic nanoparticles impregnated chitosan beads for arsenic removal from water,” *Chemical Engineering Journal*, vol. 251, pp. 25–34, 2014.
- [28] U. K. Sahu, M. K. Sahu, S. S. Mohapatra, and R. K. Patel, “Removal of As(V) from aqueous solution by Ce-Fe bimetal mixed oxide,” *Journal of Environmental Chemical Engineering*, vol. 4, no. 3, pp. 2892–2899, 2016.
- [29] C. Núñez, V. Arancibia, and M. Gómez, “Determination of arsenic in the presence of copper by adsorptive stripping voltammetry using pyrrolidine dithiocarbamate or diethyl dithiophosphate as chelating-adsorbent agents. Effect of CPB on the sensitivity of the method,” *Microchemical Journal*, vol. 126, pp. 70–75, 2016.
- [30] M. M. Bradley, L. M. Siperko, and M. D. Porter, “Colorimetric-solid phase extraction method for trace level determination of arsenite in water,” *Talanta*, vol. 86, pp. 64–70, 2011.
- [31] J. Ma, M. K. Sengupta, D. Yuan, and P. K. Dasgupta, “Speciation and detection of arsenic in aqueous samples : A review of recent progress in non-atomic spectrometric methods,” *Analytica Chimica Acta*, vol. 831, pp. 1–23, 2014.

- [32] K. Morita and E. Kaneko, "Spectrophotometric determination of arsenic in water samples based on micro particle formation of ethyl violet-molybdoarsenate.," *Japan Society for Analytical Chemistry*, vol. 22, no. 8, pp. 1085–1089, 2006.
- [33] S. Hu, J. Lu, and C. Jing, "A novel colorimetric method for field arsenic speciation analysis," *Journal of Environmental Sciences*, vol. 24, no. 7, pp. 1341–1346, 2012.
- [34] C. Baltazar, T. Igarashi, and T. Yoneda, "Applied Geochemistry Mobilization and speciation of arsenic from hydrothermally altered rock containing calcite and pyrite under anoxic conditions," *Applied Geochemistry*, vol. 27, no. 12, pp. 2300–2314, 2012.
- [35] C. B. Tabelin and T. Igarashi, "Mechanisms of arsenic and lead release from hydrothermally altered rock," *Journal of Hazardous Materials*, vol. 169, pp. 980–990, 2009.
- [36] A. Romero, I. González, and E. Gala, "Estimation of potential pollution of waste mining dumps at Peña del Hierro (Pyrite Belt , SW Spain) as a base for future mitigation actions," *Applied Geochemistry*, vol. 21, pp. 1093–1108, 2006.
- [37] P. Tschakert and K. Singha, "Contaminated identities: Mercury and marginalization in Ghana's artisanal mining sector," *Geoforum*, vol. 38, no. 6, pp. 1304–1321, 2007.
- [38] G. M. Ritcey, "Tailings management in gold plants," *Hydrometallurgy*, vol. 78, no. 1–2 SPEC. ISS., pp. 3–20, 2005.
- [39] D. Kossoff, W. E. Dubbin, M. Alfredsson, S. J. Edwards, M. G. Macklin, and K. A. Hudson-Edwards, "Mine tailings dams: Characteristics, failure, environmental impacts, and remediation," *Applied Geochemistry*, vol. 51, pp. 229–245, 2014.
- [40] M. B. J. Lindsay, M. C. Moncur, J. G. Bain, J. L. Jambor, C. J. Ptacek, and D. W. Blowes, "Geochemical and mineralogical aspects of sulfide mine tailings," *Applied Geochemistry*, vol. 57, pp. 157–177, 2015.

- [41] E. Remon, J. L. Bouchardon, B. Cornier, B. Guy, J. C. Leclerc, and O. Faure, "Soil characteristics, heavy metal availability and vegetation recovery at a former metallurgical landfill: Implications in risk assessment and site restoration," *Environmental Pollution*, vol. 137, no. 2, pp. 316–323, 2005.
- [42] C. Néel, H. Bril, A. Courtin-Nomade, and J. P. Dutreuil, "Factors affecting natural development of soil on 35-year-old sulphide-rich mine tailings," *Geoderma*, vol. 111, no. 1–2, pp. 1–20, 2003.
- [43] G. J. Zagury, K. Oudjehani, and L. Deschênes, "Characterization and availability of cyanide in solid mine tailings from gold extraction plants," *Science of the Total Environment*, vol. 320, no. 2–3, pp. 211–224, 2004.
- [44] M. M. Chico, G. Damian, D. Stumbea, N. Buzgar, T. Ungureanu, V. Nica, and G. Iepure, "Mineralogy and geochemistry of the tailings pond from Straja Valley (Suceava County, Romania). Factors affecting the mobility of the elements on the surface of the waste deposit," *Carpathian Journal of Earth and Environmental Sciences*, vol. 11, no. 1, 2016.
- [45] S. Abdul-Wahab and F. Marikar, "The environmental impact of gold mines: pollution by heavy metals," *Open Engineering*, vol. 2, no. 2, pp. 304–313, 2012.
- [46] E. R. Harish and M. David, "Assessment of Potentially Toxic Cyanide from the Gold and Copper Mine Ore Tailings of Karnataka , India," *The International Journal Of Science & Technoledge*, vol. 3, no. 7, pp. 171–178, 2015.
- [47] E. R. Harish and M. David, "Assessment of Potentially Toxic Cyanide from the Gold and Copper Mine Ore Tailings of Karnataka , India Abstract :," *The International Journal Of Science & Technoledge*, vol. 3, no. 7, pp. 171–178, 2015.
- [48] G. M. Mudd, "Global trends in gold mining: Towards quantifying environmental and resource sustainability," *Resources Policy*, vol. 32, no. 1–2, pp. 42–56, 2007.
- [49] R. Adler Miserendino, B. A. Bergquist, S. E. Adler, J. R. D. Guimarães, P. S.

- J. Lees, W. Niquen, P. C. Velasquez-López, and M. M. Veiga, “Challenges to measuring, monitoring, and addressing the cumulative impacts of artisanal and small-scale gold mining in Ecuador,” *Resources Policy*, vol. 38, no. 4, pp. 713–722, 2013.
- [50] J. Bosse Jønsson, E. Charles, and P. Kalvig, “Toxic mercury versus appropriate technology: Artisanal gold miners’ retort aversion,” *Resources Policy*, vol. 38, no. 1, pp. 60–67, 2013.
- [51] A. Kumah, “Sustainability and gold mining in the developing world,” *Journal of Cleaner Production*, vol. 14, no. 3–4, pp. 315–323, 2006.
- [52] M. Williams, “Arsenic in mine waters: International study,” *Environmental Geology*, vol. 40, no. 3, pp. 267–278, 2001.
- [53] L. Meunier, I. Koch, and K. J. Reimer, “Effects of dissolution kinetics on bioaccessible arsenic from tailings and soils,” *Chemosphere*, vol. 84, no. 10, pp. 1378–1385, 2011.
- [54] W. X. Zhang, “Nanoscale iron particles for environmental remediation: An overview,” *Journal of Nanoparticle Research*, vol. 5, no. 3–4, pp. 323–332, 2003.
- [55] S. A. Welch, A. G. Christy, L. Isaacson, and D. Kirste, “Mineralogical control of rare earth elements in acid sulfate soils,” *Geochimica et Cosmochimica Acta*, vol. 73, no. 1, pp. 44–64, 2009.
- [56] A. Khamar Shah, *Mesothermal lode gold deposit Central Belt Peninsular Malaysia*. INTECH Open Access Publisher, 2012.
- [57] N. Breward, M. Williams, C. Stanley, and R. Bowell, “Preliminary appraisal of the environmental geochemistry of the Bau mining area, Sarawak, Malaysia,” United, 1994.
- [58] A. El Azhari, A. Rhoujjati, and M. L. EL Hachimi, “Assessment of heavy metals and arsenic contamination in the sediments of the Moulouya River and the Hassan II Dam downstream of the abandoned mine Zeïda (High Moulouya, Morocco),” *Journal of African Earth Sciences*, vol. 119, pp. 279–288, 2016.

- [59] P. L. Smedley and D. G. Kinniburgh, "A review of the source, behaviour and distribution of arsenic in natural waters," *Applied Geochemistry*, vol. 17, no. 5, pp. 517–568, 2002.
- [60] E. Deschamps, V. S. T. Ciminelli, and W. H. Höll, "Removal of As(III) and As(V) from water using a natural Fe and Mn enriched sample," *Water Research*, vol. 39, no. 20, pp. 5212–5220, 2005.
- [61] P. Drahotka, M. Filippi, V. Ettler, J. Rohovec, M. Mihaljevič, and O. Šebek, "Natural attenuation of arsenic in soils near a highly contaminated historical mine waste dump," *Science of the Total Environment*, vol. 414, pp. 546–555, 2012.
- [62] E. Deschamps, V. S. T. Ciminelli, F. T. Lange, J. Matschullat, B. Raue, and H. Schmidt, "Soil and Sediment Geochemistry of the Iron Quadrangle, Brazil," *Journal of Soils and Sediments*, vol. 4, no. 3, pp. 169–169, 2002.
- [63] P. J. Kavanagh, M. E. Farago, I. Thornton, and R. S. Braman, "Bioavailability of arsenic in soil and mine wastes of the Tamar valley, SW England," *Chemical Speciation and Bioavailability*, vol. 9, no. 3, pp. 77–81, 1997.
- [64] J. R. Hale, A. Foos, J. S. Zubrow, and J. Cook, "Better characterization of arsenic and chromium in soils: a field-scale example.," *J. Soil Contam.*, vol. 6, no. December, pp. 371–389, 1997.
- [65] J. M. Azcue, A. Mudroch, F. Rosa, G. E. M. Hall, T. A. Jackson, and T. Reynoldson, "Trace elements in water, sediments, porewater, and biota polluted by tailings from an abandoned gold mine in British Columbia, Canada," *Journal of Geochemical Exploration*, vol. 52, no. 1–2, pp. 25–34, 1995.
- [66] H. W. Yaacob, W.Z.W., Pauzi, N.S.M., Mutalib, "Acid mine drainage and heavy metals contamination at abandoned and active mine sites in Pahang," *Bulletin of the Geological Society of Malaysia*, vol. 55, pp. 15–20, 2009.
- [67] P. L. Smedley and D. G. Kinniburgh, "A review on the sources, behavior and distribution of arsenic in natural waters," *Applied Geochemistry*, vol. 17, pp. 517–568, 2002.

- [68] D. K. Nordstrom, "Worldwide Occurrences of Arsenic in Ground Water," *Science*, vol. 296, no. June, pp. 64–65, 2002.
- [69] G. S. Plumlee, K. S. Smith, M. R. Montour, W. H. Flicklin, and E. L. Mosier, "Geologic controls on the composition of natural waters and mine waters draining diverse mineral-deposit types," *The environmental geochemistry of mineral deposits. Part B: case studies and research topics*, vol. 6, pp. 373–432, 1999.
- [70] H. Mccreadie, D. W. Blowes, C. J. Ptacek, and J. L. Jambor, "Influence of reduction reactions and solid-phase composition on porewater concentrations of arsenic," *Environmental Science and Technology*, vol. 34, no. 15, pp. 3159–3166, 2000.
- [71] F. H. Wilson and D. B. Hawkins, "Arsenic in streams, stream sediments, and ground water, fairbanks area, alaska," *Environmental Geology*, vol. 2, no. 4, pp. 195–202, 1978.
- [72] M. F. Lengke and R. N. Tempel, "Geochemical modeling of arsenic sulfide oxidation kinetics in a mining environment," *Geochimica et Cosmochimica Acta*, vol. 69, no. 2, pp. 341–356, 2005.
- [73] A. alshaebi Fares Yahya, Y. Wan Zuhairi Wan, S. Abdul Rahim, and A. Esmail, "Risk Assessment at Abandoned Tin Mine," *Electronic Journal of Geotechnical Engineering*, vol. 14, pp. 1–9, 2009.
- [74] N. Suppen, M. Carranza, M. Huerta, and M. A. Hernández, "Environmental management and life cycle approaches in the Mexican mining industry," *Journal of Cleaner Production*, vol. 14, no. 12–13 SPEC. ISS., pp. 1101–1115, 2006.
- [75] G. Hilson and B. Murck, "Progress toward pollution prevention and waste minimization in the North American gold mining industry," *Journal of Cleaner Production*, vol. 9, pp. 405–415, 2001.
- [76] M. M. Veiga, G. Angeloci, M. Hitch, and P. Colon Velasquez-Lopez, "Processing centres in artisanal gold mining," *Journal of Cleaner Production*, vol. 64, pp. 535–544, 2014.

- [77] G. M. Mudd, "Gold mining in Australia: linking historical trends and environmental and resource sustainability," *Environmental Science and Policy*, vol. 10, no. 7–8, pp. 629–644, 2007.
- [78] A. Whitmore, "The emperors new clothes: Sustainable mining?," *Journal of Cleaner Production*, vol. 14, no. 3–4, pp. 309–314, 2006.
- [79] J. E. S. Quarrie, "Earth Summit'92. The UN Conference on Environment and Development," in *United Nations Sustainable development*, 1992, p. 351.
- [80] "Environmental Quality Act (EQA), 1974. Laws of Malaysia. Act 127. International Law Book Services."
- [81] *Mineral Development Act, 1994. Laws of Malaysia. Act 525. Sengupta, M., 2013. Environmental Impacts of Mining. Lewis Publishers. .*
- [82] Department of Environment, "Environmental Requirements: A Guide For Investors. Department of Environment Ministry of Natural Resources and Environmen Wisma Sumber Asli, Precinct 4 Federal Government Administrative Centre 62574 PUTRAJAYA Eleventh Edition October 2010," pp. 1–78.
- [83] D. Mohan and C. U. Pittman, "Arsenic removal from water/wastewater using adsorbents-A critical review," *Journal of Hazardous Materials*, vol. 142, no. 1–2, pp. 1–53, 2007.
- [84] S. K. Maji, Y. Kao, C. Wang, G. Lu, J. Wu, and C. Liu, "Fixed bed adsorption of As (III) on iron-oxide-coated natural rock (IOCNR) and application to real arsenic-bearing groundwater," *Chemical Engineering Journal*, vol. 203, pp. 285–293, 2012.
- [85] S. Hokkanen, E. Repo, S. Lou, and M. Sillanpää, "Removal of arsenic(V) by magnetic nanoparticle activated microfibrillated cellulose," *Chemical Engineering Journal*, vol. 260, pp. 886–894, 2015.
- [86] D. Budimirović, Z. S. Veličković, V. R. Djokić, M. Milosavljević, J. Markovski, S. Lević, and A. D. Marinković, "Efficient As(V) removal by α -FeOOH and α -FeOOH/ α -MnO₂ embedded PEG-6-arm functionalized multiwall carbon nanotubes," *Chemical Engineering Research and Design*,

- vol. 119, no. V, pp. 75–86, 2017.
- [87] D. Ociński, I. Jacukowicz-Sobala, P. Mazur, J. Raczyk, and E. Kociołek-Balawejder, “Water treatment residuals containing iron and manganese oxides for arsenic removal from water - Characterization of physicochemical properties and adsorption studies,” *Chemical Engineering Journal*, vol. 294, pp. 210–221, 2016.
- [88] A. Sigdel, J. Park, H. Kwak, and P. K. Park, “Arsenic removal from aqueous solutions by adsorption onto hydrous iron oxide-impregnated alginate beads,” *Journal of Industrial and Engineering Chemistry*, vol. 35, pp. 277–286, 2015.
- [89] D. Shaw, “Mobility of arsenic in saturated, laboratory test sediments under varying pH conditions,” *Engineering Geology*, vol. 85, no. 1–2, pp. 158–164, 2006.
- [90] J. N. Geroni, R. J. Howell, M. Dey, D. J. Sapsford, and K. P. Williams, “Removal of Arsenic (III) from Contaminated Waters using Iron (II) and Citrate,” *In Proceedings of the 8th International Conference on Acid Rock Drainage (ICARD) and securing the future: mining, metals & the environment in a sustainable society*, pp. 22–26, 2009.
- [91] R. Li, Q. Li, S. Gao, and J. K. Shang, “Exceptional arsenic adsorption performance of hydrous cerium oxide nanoparticles: Part A. Adsorption capacity and mechanism,” *Chemical Engineering Journal*, vol. 185–186, pp. 127–135, 2012.
- [92] A. Santos, F. W. F. de Oliveira, F. H. A. Silva, D. A. Maria, J. D. Ardisson, W. A. de A. Macêdo, H. E. L. Palmieri, and M. B. Franco, “Synthesis and characterization of iron-PVA hydrogel microspheres and their use in the arsenic (V) removal from aqueous solution,” *Chemical Engineering Journal*, vol. 210, pp. 432–443, 2012.
- [93] M. A. Malana, R. B. Qureshi, and M. N. Ashiq, “Adsorption studies of arsenic on nano aluminium doped manganese copper ferrite polymer (MA, VA, AA) composite: Kinetics and mechanism,” *Chemical Engineering Journal*, vol. 172, no. 2–3, pp. 721–727, 2011.

- [94] B. Pan, Z. Li, Y. Zhang, J. Xu, L. Chen, H. Dong, and W. Zhang, "Acid and organic resistant nano-hydrated zirconium oxide (HZO)/poly-styrene hybrid adsorbent for arsenic removal from water," *Chemical Engineering Journal*, vol. 248, pp. 290–296, 2014.
- [95] Y. Wang, S. Sikora, H. Kim, B. Dubey, and T. Townsend, "Mobilization of iron and arsenic from soil by construction and demolition debris landfill leachate," *Waste Management*, vol. 32, no. 5, pp. 925–932, 2012.
- [96] Wikipedia, "Gold cyanidation." [Online]. Available: https://en.wikipedia.org/wiki/Gold_cyanidation. [Accessed: 20-Feb-2017].
- [97] M. F. Hiji and S. P. Maganga, "Application of Response Surface Methodology for Optimization of Vat Leaching Parameters in Small Scale Mines: Case Study of Tanzania," *American Journal of Engineering, Technology and Society*, vol. 2, no. 2, pp. 21–25, 2015.
- [98] S. Ravengai, D. Love, M. Mabvira-Meck, K. Musiwa, and W. Moyce, "Water quality in an abandoned gold mining belt, Beatrice, Sanyati Valley, Zimbabwe," *Physics and Chemistry of the Earth*, vol. 30, no. 11–16 SPEC. ISS., pp. 826–831, 2005.
- [99] A. Courtin-Nomade, H. Bril, C. Neel, and J. F. Lenain, "Arsenic in iron cements developed within tailings of a former metalliferous mine - Enguiales, Aveyron, France," *Applied Geochemistry*, vol. 18, no. 3, pp. 395–408, 2003.
- [100] P. C. S. Carvalho, A. M. R. Neiva, M. M. V. G. Silva, and E. A. F. da Silva, "Geochemical comparison of waters and stream sediments close to abandoned Sb-Au and As-Au mining areas, northern Portugal," *Chemie der Erde - Geochemistry*, vol. 74, no. 2, pp. 267–283, 2014.
- [101] D. Craw, D. Chappell, A. Reay, and D. Walls, "New Zealand Journal of Geology and Geophysics Mobilisation and attenuation of arsenic around gold mines, east Otago, New Zealand Mobilisation and attenuation of arsenic around gold mines, east Otago, New Zealand," *New Zealand Journal of Geology and Geophysics New Zealand Journal of Geology & Geophysics*, vol. 43, no. 43, pp. 373–383, 2000.

- [102] L. Haffert and D. Craw, "Geochemical processes influencing arsenic mobility at Bullendale historic gold mine, Otago, New Zealand," *New Zealand Journal of Geology and Geophysics*, vol. 53, no. 2–3, pp. 129–142, 2010.
- [103] L. Haffert, D. Craw, and J. Pope, "Climatic and compositional controls on secondary arsenic mineral formation in high-arsenic mine wastes, South Island, New Zealand," *New Zealand Journal of Geology and Geophysics*, vol. 53, no. 2–3, pp. 91–101, 2010.
- [104] J. Wurl, L. Mendez-Rodriguez, and B. Acosta-Vargas, "Arsenic content in groundwater from the southern part of the San Antonio-El Triunfo mining district, Baja California Sur, Mexico," *Journal of Hydrology*, vol. 518, no. PC, pp. 447–459, 2014.
- [105] C. Makoundi, K. Zaw, R. R. Large, S. Meffre, C. K. Lai, and T. G. Hoe, "Geology, geochemistry and metallogenesis of the Selinsing gold deposit, central Malaysia," *Gondwana Research*, vol. 26, no. 1, pp. 241–261, 2014.
- [106] B. Palumbo-Roe, J. Wragg, and M. Cave, "Linking selective chemical extraction of iron oxyhydroxides to arsenic bioaccessibility in soil," *Environmental Pollution*, vol. 207, pp. 256–265, 2015.
- [107] A. M. Sánchez de la Campa, J. D. de la Rosa, J. C. Fernández-Caliani, and Y. González-Castanedo, "Impact of abandoned mine waste on atmospheric respirable particulate matter in the historic mining district of Rio Tinto (Iberian Pyrite Belt)," *Environmental Research*, vol. 111, no. 8, pp. 1018–1023, 2011.
- [108] A. Courtin-Nomade, C. Grosbois, H. Bril, and C. Roussel, "Spatial variability of arsenic in some iron-rich deposits generated by acid mine drainage," *Applied Geochemistry*, vol. 20, no. 2, pp. 383–396, 2005.
- [109] J. H. Park, Y.-S. Han, and J. S. Ahn, "Comparison of arsenic co-precipitation and adsorption by iron minerals and the mechanism of arsenic natural attenuation in a mine stream," *Water Research*, vol. 106, pp. 295–303, 2016.
- [110] L. M. Jackson and A. Parbhakar-Fox, "Mineralogical and geochemical characterization of the Old Tailings Dam, Australia: Evaluating the

- effectiveness of a water cover for long-term AMD control,” *Applied Geochemistry*, vol. 68, pp. 64–78, 2016.
- [111] S. Tamoto, C. B. Tabelin, T. Igarashi, M. Ito, and N. Hiroyoshi, “Short and long term release mechanisms of arsenic, selenium and boron from a tunnel-excavated sedimentary rock under in situ conditions,” *Journal of Contaminant Hydrology*, vol. 175–176, pp. 60–71, 2015.
- [112] A. Bhattacharya, J. Routh, G. Jacks, P. Bhattacharya, and M. Mörth, “Environmental assessment of abandoned mine tailings in Adak, Västerbotten district (northern Sweden),” *Applied Geochemistry*, vol. 21, no. 10, pp. 1760–1780, 2006.
- [113] M. A. Sobahan, S. I. Mir, I. Zakaria, M. A. Hossain, and A. S. Area, “Surface Water Contamination Due To Industrial Activities in Gebeng Area , Kuantan , Malaysia,” *International Conference on Civil and Architecture Engineering*, 2013.
- [114] N. H. Ab Razak, S. M. Praveena, A. Z. Aris, and Z. Hashim, “Quality of Kelantan drinking water and knowledge, attitude and practice among the population of Pasir Mas, Malaysia,” *Public Health*, vol. 131, pp. 103–111, 2016.
- [115] N. H. Ab Razak, S. M. Praveena, A. Z. Aris, and Z. Hashim, “Drinking water studies: A review on heavy metal, application of biomarker and health risk assessment (a special focus in Malaysia),” *Journal of Epidemiology and Global Health*, vol. 5, no. 4, pp. 297–310, 2015.
- [116] V. D. Martinez, E. A. Vucic, D. D. Becker-santos, L. Gil, and W. L. Lam, “Arsenic Exposure and the Induction of Human Cancers,” *Journal of Toxicology*, vol. 2011, p. 13, 2011.
- [117] F. Hossen, S. Hamdan, and R. Rahman, “Review on the Risk Assessment of Heavy Metals in Malaysian Clams,” *The Scientific World Journal*, vol. 2015, p. 7, 2015.
- [118] K. Lias, T. Jamil, and S. N. Aliaa, “A Preliminary Study on Heavy Metal Concentration in the Marine Bivalves *Marcia Marmorata* Species and

- Sediments Collected From the Coastal Area of Kuala Perlis, North Of Malaysia,” *Journal of Applied Chemistry*, vol. 4, no. 1, pp. 48–54, 2013.
- [119] A. Taweel, M. Shuhaimi-othman, and A. K. Ahmad, “Evaluation of copper, lead and arsenic level in tilapia fish in Cempaka Lake (Bangi , Malaysia) and human daily / weekly intake,” *VERSITA*, vol. 68, no. 5, p. 983—991, 2013.
- [120] A. SHAH, “Water , fish contain high level of arsenic,” *New Straits Time Online*, 2015. [Online]. Available: <http://www.nst.com.my/news/2015/09/water-fish-contain-high-level-arsenic>.
- [121] D. D’Ippoliti, E. Santelli, M. De Sario, M. Scortichini, M. Davoli, and P. Michelozzi, “Arsenic in drinking water and mortality for cancer and chronic diseases in Central Italy, 1990-2010,” *PLOS ONE*, vol. 10, no. 9, pp. 1–19, 2015.
- [122] M. Hutton and U. Kingdom, *Human Health Concerns of Lead , Mercury , Cadmium and Arsenic*. John Wiley & Sons Ltd, 1987.
- [123] EPA, *Abandoned mine site characterization and cleanup handbook (EPA 910-B-00-001)*. Washington, D.C.: EPA, 2000.
- [124] C. B. Tabelin, A. Hashimoto, T. Igarashi, and T. Yoneda, “Leaching of boron, arsenic and selenium from sedimentary rocks: II. pH dependence, speciation and mechanisms of release,” *Science of the Total Environment*, vol. 473–474, pp. 244–253, 2014.
- [125] C. B. Tabelin, T. Igarashi, S. Tamoto, and R. Takahashi, “The roles of pyrite and calcite in the mobilization of arsenic and lead from hydrothermally altered rocks excavated in Hokkaido, Japan,” *Journal of Geochemical Exploration*, vol. 119–120, pp. 17–31, 2012.
- [126] V. Chatain, F. Sanchez, R. Bayard, P. Moszkowicz, and R. Gourdon, “Effect of experimentally induced reducing conditions on the mobility of arsenic from a mining soil,” *Journal of Hazardous Materials*, vol. 122, no. 1–2, pp. 119–128, 2005.
- [127] R. Bayard, V. Chatain, C. Gachet, A. Troadec, and R. Gourdon, “Mobilisation of arsenic from a mining soil in batch slurry experiments under bio-oxidative

- conditions,” *Water Research*, vol. 40, no. 6, pp. 1240–1248, 2006.
- [128] C. B. Tabelin, T. Igarashi, T. Arima, D. Sato, T. Tatsuhara, and S. Tamoto, “Characterization and evaluation of arsenic and boron adsorption onto natural geologic materials, and their application in the disposal of excavated altered rock,” *Geoderma*, vol. 213, pp. 163–172, 2014.
- [129] A. Maiti, J. K. Basu, and S. De, “Experimental and kinetic modeling of As(V) and As(III) adsorption on treated laterite using synthetic and contaminated groundwater: Effects of phosphate, silicate and carbonate ions,” *Chemical Engineering Journal*, vol. 191, pp. 1–12, 2012.
- [130] T. Tatsuhara, T. Arima, T. Igarashi, and C. B. Tabelin, “Combined neutralization-adsorption system for the disposal of hydrothermally altered excavated rock producing acidic leachate with hazardous elements,” *Engineering Geology*, vol. 139–140, pp. 76–84, 2012.
- [131] S. A. Akoitai, “Arsenic in Leachates From Mine Waste Rocks,” 2000.
- [132] M. A. Acheampong, J. Adiyiah, E. David, and O. Ansa, “Physico-chemical Characteristics of a Gold Mining Tailings Dam Wastewater,” *Journal of Environmental Science and Engineering A*, vol. 2, pp. 469–475, 2013.
- [133] D. B. Donato, O. Nichols, H. Possingham, M. Moore, P. F. Ricci, and B. N. Noller, “A critical review of the effects of gold cyanide-bearing tailings solutions on wildlife,” *Environment International*, vol. 33, no. 7, pp. 974–984, 2007.
- [134] N. Kuyucak and A. Akcil, “Cyanide and removal options from effluents in gold mining and metallurgical processes,” *Minerals Engineering*, vol. 50–51, pp. 13–29, 2013.
- [135] a G. M. Zabawi, S. M. Esa, and C. P. Leong, “Effects of simulated acid rain on germination and growth of rice plant,” *Journal of Tropical Agriculture and Food Science*, vol. 36, no. 2, pp. 1–6, 2008.
- [136] W. N. A. Sulaiman, H. A., and M. H. RosliA., “Identification of Flood Source Areas in Pahang River Basin, Peninsular Malaysia,” *EnvironmentAsia*, vol. 3, no. special Issue, pp. 73–78, 2010.

- [137] J. Searles, "Extraction and analysis of cyanide in soil and sédiment samples," p. 7 pages, 2014.
- [138] EPA, "Treatment of Cyanide Heap Leaches and Tailings," Washington, 1994.
- [139] A. D. Redman, D. Macalady, and D. Ahmann, "Natural organic matter affects arsenic speciation and sorption onto hematite RN - Environ. Sci. Technol., vol. 36, pp. 2889-2896," *Environmental Science and Technology*, vol. 36, no. 13, pp. 2889–2896, 2002.
- [140] K. P. Nguyen, R. Itoi, and R. Yamashiro, "Influence of Redox Potential on Arsenic Release from Soil in the Presence of Iron Oxyhydroxide," *Memoirs of the Faculty of Engineering, Kyushu University*, vol. 69, no. 2, 2009.
- [141] F. Martin, M. Simon, E. Arco, A. Romero, and C. Dorronsoro, "Arsenic behaviour in polluted soils after remediation activities," in *Soil Health and Land Use Management*, M. C. H. Soriano, Ed. Shanghai,China: InTech Published, 2012, pp. 201–2017.
- [142] N. Mladenov, Y. Zheng, M. P. Miller, D. R. Nemerlut, T. Legg, B. Simone, C. Hageman, M. M. Rahman, K. M. Ahmed, and D. M. Mcknight, "Dissolved organic matter sources and consequences for iron and arsenic mobilization in Bangladesh aquifers," *Environmental Science and Technology*, vol. 44, no. 1, pp. 123–128, 2010.
- [143] S. Wang and C. N. Mulligan, "Natural attenuation processes for remediation of arsenic contaminated soils and groundwater," *Journal of Hazardous Materials*, vol. 138, no. 3, pp. 459–470, 2006.
- [144] D. Mohapatra, D. Mishra, M. Rout, and G. R. Chaudhury, "Adsorption kinetics of natural dissolved organic matter and its impact on arsenic(V) leachability from arsenic-loaded ferrihydrite and Al-ferrihydrite," *Journal of environmental science and health. Part A*, vol. 42, no. 1, pp. 81–88, 2007.
- [145] D. E. Giles, M. Mohapatra, T. B. Issa, S. Anand, and P. Singh, "Iron and aluminium based adsorption strategies for removing arsenic from water," *Journal of Environmental Management*, vol. 92, no. 12, pp. 3011–3022, 2011.
- [146] C. B. Tabelin, A. Hashimoto, T. Igarashi, and T. Yoneda, "Leaching of boron,

- arsenic and selenium from sedimentary rocks: I. Effects of contact time, mixing speed and liquid-to-solid ratio,” *Science of the Total Environment*, vol. 472, pp. 620–629, 2014.
- [147] A. J. Bora, S. Gogoi, G. Baruah, and R. K. Dutta, “Utilization of co-existing iron in arsenic removal from groundwater by oxidation-coagulation at optimized pH,” *Journal of Environmental Chemical Engineering*, vol. 4, no. 3, pp. 2683–2691, 2016.
- [148] B. F. Urbano, B. L. Rivas, F. Martinez, and S. D. Alexandratos, “Water-insoluble polymer-clay nanocomposite ion exchange resin based on N-methyl-d-glucamine ligand groups for arsenic removal,” *Reactive and Functional Polymers*, vol. 72, no. 9, pp. 642–649, 2012.
- [149] S. A. Schmidt, E. Gukelberger, M. Hermann, F. Fiedler, B. Großmann, J. Hoinkis, A. Ghosh, D. Chatterjee, and J. Bundschuh, “Pilot study on arsenic removal from groundwater using a small-scale reverse osmosis system—Towards sustainable drinking water production,” *Journal of Hazardous Materials*, vol. 318, pp. 671–678, 2016.
- [150] E. Fogarassy, I. Galambos, E. Bekassy-Molnar, and G. Vatai, “Treatment of high arsenic content wastewater by membrane filtration,” *Desalination*, vol. 240, no. 1–3, pp. 270–273, 2009.
- [151] V. Bhadja, J. S. Trivedi, and U. Chatterjee, “Efficacy of polyethylene Interpolymer membranes for fluoride and arsenic ion removal during desalination of water via electrodialysis,” *RSC Advances*, vol. 6, no. 71, pp. 67118–67126, 2016.
- [152] D. Cortés-Arriagada and A. Toro-Labbé, “Aluminum and iron doped graphene for adsorption of methylated arsenic pollutants,” *Applied Surface Science*, vol. 386, pp. 84–95, 2016.
- [153] M. Bhaumik, C. Noubactep, V. Kumar, R. I. Mccrindle, and A. Maity, “Polyaniline / Fe 0 composite nanofibers: An excellent adsorbent for the removal of arsenic from aqueous solutions,” *Chemical Engineering Journal*, vol. 271, pp. 135–146, 2015.

- [154] T. Basu, D. Nandi, P. Sen, and U. C. Ghosh, "Equilibrium modeling of As(III,V) sorption in the absence/presence of some groundwater occurring ions by iron(III)-cerium(IV) oxide nanoparticle agglomerates: A mechanistic approach of surface interaction," *Chemical Engineering Journal*, vol. 228, pp. 665–678, 2013.
- [155] J. M. Calo, L. Madhavan, J. Kirchner, and E. J. Bain, "Arsenic removal via ZVI in a hybrid spouted vessel/fixed bed filter system," *Chemical Engineering Journal*, vol. 189–190, pp. 237–243, 2012.
- [156] S. Bhowmick, S. Chakraborty, P. Mondal, and W. Van Renterghem, "Montmorillonite-supported nanoscale zero-valent iron for removal of arsenic from aqueous solution: Kinetics and mechanism," *Chemical Engineering Journal*, vol. 243, pp. 14–23, 2014.
- [157] R. A. Crane, D. E. Sinnett, P. J. Cleall, and D. J. Sapsford, "Physicochemical composition of wastes and co-located environmental designations at legacy mine sites in the south west of England and Wales: Implications for their resource potential," *Resources, Conservation & Recycling*, 2016.
- [158] C. B. Tabelin, T. Igarashi, and S. Tamoto, "Factors affecting arsenic mobility from hydrothermally altered rock in impoundment-type in situ experiments," *Minerals Engineering*, vol. 23, no. 3, pp. 238–248, 2010.
- [159] Z. M. Migaszewski, A. Gałuszka, S. Hałas, S. Dołęgowska, J. Dabek, and E. Starnawska, "Geochemistry and stable sulfur and oxygen isotope ratios of the Podwiśniówka pit pond water generated by acid mine drainage (Holy Cross Mountains, south-central Poland)," *Applied Geochemistry*, vol. 23, no. 12, pp. 3620–3634, 2008.
- [160] K. Rout, M. Mohapatra, and S. Anand, "2-line ferrihydrite: synthesis, characterization and its adsorption behaviour for removal of Pb(II), Cd(II), Cu(II) and Zn(II) from aqueous solutions.," *Dalton Transactions*, vol. 41, no. 11, pp. 3302–12, 2012.
- [161] L. Wang and D. E. Giammar, "Effects of pH, dissolved oxygen, and aqueous ferrous iron on the adsorption of arsenic to lepidocrocite," *Journal of Colloid*

- and Interface Science*, vol. 448, pp. 331–338, 2015.
- [162] F. M. Romero, C. Canet, P. Alfonso, R. N. Zambrana, and N. Soto, “The role of cassiterite controlling arsenic mobility in an abandoned stanniferous tailings impoundment at Llallagua, Bolivia,” *Science of the Total Environment*, vol. 481, no. 1, pp. 100–107, 2014.
- [163] N. Bektas, S. Aydin, and M. S. Oncel, “The Adsorption of Arsenic Ions Using Beidellite, Zeolite, and Sepiolite Clays: A Study of Kinetic, Equilibrium and Thermodynamics,” *Separation Science and Technology*, vol. 46, no. 6, pp. 1005–1016, 2011.
- [164] R. Huang, Y. Liu, Z. Chen, D. Pan, Z. Li, M. Wu, C. H. Shek, C. M. L. Wu, and J. K. L. Lai, “Fe-species-loaded mesoporous MnO₂ superstructural requirements for enhanced catalysis,” *ACS Applied Materials and Interfaces*, vol. 7, no. 7, pp. 3949–3959, 2015.
- [165] E. Jansen, A. Kyek, W. Schäfer, and U. Schwertmann, “The structure of six-line ferrihydrite,” *Applied Physics A: Materials Science and Processing*, vol. 74, no. SUPPL.II, pp. 1004–1006, 2002.
- [166] C. W. Childs, N. Matsue, and N. Yoshinaga, “Soil Science and Plant Nutrition Ferrihydrite in volcanic ash soils of Japan Ferrihydrite in Volcanic Ash Soils of Japan,” *Soil Science and Plant Nutrition*, vol. 37, no. 2, pp. 299–311, 1990.
- [167] M. Mohapatra, L. Mohapatra, D. Hariprasad, S. Anand, B. K. Mishra, L. Mohapatra, D. Hariprasad, S. Anand, and B. K. Mishra, “Nano-structured Mg-doped Fe₂O₃ – ferrihydrite powder – a new adsorbent for cation removal from aqueous solutions,” *Environmental Technology*, vol. 33, no. 15, pp. 1717–1726, 2012.
- [168] C. A. J. Appelo, M. J. J. Van Der Weiden, C. Tournassat, and L. Charlet, “Surface complexation of ferrous iron and carbonate on ferrihydrite and the mobilization of arsenic,” *Environmental Science and Technology*, vol. 36, no. 14, pp. 3096–3103, 2002.
- [169] M. Mohapatra and S. Anand, “Synthesis and applications of nano-structured iron oxides / hydroxides – a review,” *International Journal of Engineering*,

- Science and Technology*, vol. 2, no. 8, pp. 127–146, 2010.
- [170] M. Jang, J. S. Hwang, and S. Il Choi, “Sequential soil washing techniques using hydrochloric acid and sodium hydroxide for remediating arsenic-contaminated soils in abandoned iron-ore mines,” *Chemosphere*, vol. 66, no. 1, pp. 8–17, 2007.
- [171] G. . A. Waychunas, B. . A. Rea, C. . C. Fuller, and J. . A. Davis, “Surface chemistry of ferrihydrite: Part 1. EXAFS studies on geometry of coprecipitated and adsorbed arsenate,” *Geochimica et Cosmochimica Acta*, vol. 57, no. 10, pp. 2251–2269, 1993.
- [172] R. K. Kukkadapu, J. M. Zachara, J. K. Fredrickson, S. C. Smith, and A. C. Dohnalkova, “Transformation of 2-line ferrihydrite to 6-line ferrihydrite under oxic and anoxic conditions,” *American Mineralogist*, vol. 88, pp. 1903–1914, 2003.
- [173] Y. Cudennec and A. Lecerf, “The transformation of ferrihydrite into goethite or hematite, revisited,” *Journal of Solid State Chemistry*, vol. 179, no. 3, pp. 716–722, 2006.
- [174] S. G. Johnston, A. F. Keene, E. D. Burton, R. T. Bush, and L. A. Sullivan, “Iron and arsenic cycling in intertidal surface sediments during wetland remediation,” *Environmental Science and Technology*, vol. 45, no. 6, pp. 2179–2185, 2011.
- [175] A. Jain, K. P. Raven, and R. H. Loeppert, “Arsenite and arsenate adsorption on ferrihydrite: Surface charge reduction and net OH⁻ release stoichiometry,” *Environmental Science and Technology*, vol. 33, no. 8, pp. 1179–1184, 1999.
- [176] C. Grosbois, A. Courtin-Nomade, E. Robin, H. Bril, N. Tamura, J. Sch??fer, and G. Blanc, “Fate of arsenic-bearing phases during the suspended transport in a gold mining district (Isle river Basin, France),” *Science of the Total Environment*, vol. 409, no. 23, pp. 4986–4999, 2011.
- [177] C. Casiot, S. Lebrun, G. Morin, O. Bruneel, J. C. Personné, and F. Elbaz-Poulichet, “Sorption and redox processes controlling arsenic fate and transport in a stream impacted by acid mine drainage,” *Science of the Total*

- Environment*, vol. 347, no. 1–3, pp. 122–130, 2005.
- [178] J. Qiao, Z. Jiang, B. Sun, Y. Sun, Q. Wang, and X. Guan, “Arsenate and arsenite removal by FeCl₃: Effects of pH, As/Fe ratio, initial As concentration and co-existing solutes,” *Separation and Purification Technology*, vol. 92, pp. 106–114, 2012.
- [179] M. Gräfe, R. V. Tappero, M. A. Marcus, and D. L. Sparks, “Arsenic speciation in multiple metal environments. II. Micro-spectroscopic investigation of a CCA contaminated soil,” *Journal of Colloid and Interface Science*, vol. 321, no. 1, pp. 1–20, 2008.
- [180] S. K. Maji, Y. H. Kao, C. J. Wang, G. S. Lu, J. J. Wu, and C. W. Liu, “Fixed bed adsorption of As(III) on iron-oxide-coated natural rock (IOCNR) and application to real arsenic-bearing groundwater,” *Chemical Engineering Journal*, vol. 203, pp. 285–293, 2012.
- [181] S. Wang, B. Gao, Y. Li, A. E. Creamer, and F. He, “Adsorptive removal of arsenate from aqueous solutions by biochar supported zero-valent iron nanocomposite: Batch and continuous flow tests,” *Journal of Hazardous Materials*, vol. 322, pp. 172–181, 2015.
- [182] H. Zhu, Y. Jia, X. Wu, and H. Wang, “Removal of arsenic from water by supported nano zero-valent iron on activated carbon,” *Journal of Hazardous Materials*, vol. 172, no. 2–3, pp. 1591–1596, 2009.
- [183] Y. Mamindy-Pajany, C. Hurel, N. Marmier, and M. Roméo, “Arsenic (V) adsorption from aqueous solution onto goethite, hematite, magnetite and zero-valent iron: Effects of pH, concentration and reversibility,” *Desalination*, vol. 281, no. 1, pp. 93–99, 2011.
- [184] A. J. Bednar, J. R. Garbarino, J. F. Ranville, and T. R. Wildeman, “Effects of iron on arsenic speciation and redox chemistry in acid mine water,” *Journal of Geochemical Exploration*, vol. 85, pp. 55–62, 2005.
- [185] D. L. Corwin, A. David, and S. Goldberg, “Mobility of arsenic in soil from the Rocky Mountain Arsenal area,” *Journal of Contaminant Hydrology*, vol. 39, no. 1–2, pp. 35–58, 1999.

- [186] K. P. Nguyen, R. Itoi, and R. Yamashiro, "Influence of redox potential on arsenic release from soil in the presence of iron oxyhydroxide," *Memoirs of the Faculty of Engineering, Kyushu University*, vol. 68, no. 2, pp. 129–140, 2008.
- [187] X. Jia, D. Gong, J. Wang, F. Huang, T. Duan, and X. Zhang, "Arsenic speciation in environmental waters by a new specific phosphine modified polymer microsphere preconcentration and HPLC–ICP-MS determination," *Talanta*, vol. 160, pp. 437–443, 2016.
- [188] W. Siangproh, O. Chailapakul, and K. Songsrirote, "Simple and fast colorimetric detection of inorganic arsenic selectively adsorbed onto ferrihydrite-coated silica gel using silver nanoplates," *Talanta*, vol. 153, pp. 197–202, 2016.
- [189] A. A. Ensa, N. Kazemifard, and B. Rezaei, "Biosensors and Bioelectronics A simple and sensitive fluorimetric aptasensor for the ultrasensitive detection of arsenic (III) based on cysteamine stabilized CdTe / ZnS quantum dots aggregation," *Biosensors and Bioelectronics*, vol. 77, pp. 499–504, 2016.
- [190] S. K. Toor, P. Devi, and B. K. S. Bansod, "Electrochemical Detection of Trace Amount of Arsenic (III) at Glassy Carbon Electrode Modified with Au/Fe₃O₄ Nanocomposites," *Aquatic Procedia*, vol. 4, no. Icwrcoc, pp. 1107–1113, 2015.
- [191] K. Shrivastava, R. Shankar, and K. Dewangan, "Gold nanoparticles as a localized surface plasmon resonance based chemical sensor for on-site colorimetric detection of arsenic in water samples," *Sensors and Actuators B*, vol. 220, pp. 1376–1383, 2015.
- [192] S. Tsang, F. Phu, M. M. Baum, and G. A. Poskrebyshev, "Determination of phosphate / arsenate by a modified molybdenum blue method and reduction of arsenate by S₂O₄²⁻," *Talanta*, vol. 71, pp. 1560–1568, 2007.
- [193] F. Juillot, P. Ildefonse, G. Morin, G. Calas, A. M. De Kersabiec, and M. Benedetti, "Remobilization of arsenic from buried wastes at an industrial site: Mineralogical and geochemical control," *Applied Geochemistry*, vol. 14, no. 8,

- pp. 1031–1048, 1999.
- [194] A. B. Mukherjee and P. Bhattacharya, “Arsenic in groundwater in the Bengal Delta Plain: slow poisoning in Bangladesh,” *Environmental Reviews*, vol. 9, no. 3, pp. 189–220, 2001.
- [195] M. Yang, Z. Guo, L. N. Li, Y. Y. Huang, J. H. Liu, Q. Zhou, X. Chen, and X. J. Huang, “Electrochemical determination of arsenic(III) with ultra-high anti-interference performance using Au-Cu bimetallic nanoparticles,” *Sensors and Actuators, B: Chemical*, vol. 231, pp. 70–78, 2016.
- [196] Y. Kiso, S. Asaoka, Y. Kamimoto, S. Tanimoto, and K. Yokota, “Detection tube method for trace level arsenic,” *Journal of Environmental Chemical Engineering*, vol. 3, no. 1, pp. 40–45, 2015.
- [197] H. Y. Eom, D. H. Yang, J. H. Suh, U. Kim, J. Kim, H. D. Cho, and S. B. Han, “Determination of residual arsenic compounds in chicken muscle by ultra-performance liquid chromatography coupled with ultraviolet detection after pre-column derivatization with toluene-3,4-dithiol,” *Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences*, vol. 1006, pp. 151–157, 2015.
- [198] É. R. Pereira, T. S. De Almeida, D. L. G. Borges, E. Carasek, B. Welz, J. Feldmann, and J. Del Campo Menoyo, “Investigation of chemical modifiers for the direct determination of arsenic in fish oil using high-resolution continuum source graphite furnace atomic absorption spectrometry,” *Talanta*, vol. 150, pp. 142–147, 2016.
- [199] J. Hao, M. J. Han, S. Han, X. Meng, T. L. Su, and Q. K. Wang, “SERS detection of arsenic in water: A review,” *Journal of Environmental Sciences*, vol. 36, no. V, pp. 152–162, 2015.
- [200] S. Li, M. Wang, Y. Zhong, Z. Zhang, and B. Yang, “Cloud point extraction for trace inorganic arsenic speciation analysis in water samples by hydride generation atomic fluorescence spectrometry,” *Spectrochimica Acta - Part B Atomic Spectroscopy*, vol. 111, pp. 74–79, 2015.
- [201] C. L. T. Correia, R. A. Goncalves, M. S. Azevedo, M. A. Vieira, and R. C.

- Campos, "Determination of total arsenic in seawater by hydride generation atomic fluorescence spectrometry," *Microchemical Journal*, vol. 96, no. January, pp. 157–160, 2010.
- [202] R. K. Dhar, Y. Zheng, J. Rubenstone, and A. Van Geen, "A rapid colorimetric method for measuring arsenic concentrations in groundwater," *Analytica Chimica Acta*, vol. 526, pp. 203–209, 2004.
- [203] E. C. Chemical, E. D. Series, and T. Roorkee, "Batch and Column Study : Adsorption of Arsenate Using Untreated Laterite as Adsorbent," no. MARCH, pp. 1620–1629, 2008.
- [204] P. Jones, R. Stanley, and N. Barnett, "Determination of arsenate, germanate, phosphate and silicate by ion chromatography using a post-column reaction (molybdenum blue) detector," *Analytica Chimica Acta*, vol. 249, pp. 539–544, 1991.
- [205] R. E. Stauffer, "Molybdenum blue applied to arsenic and phosphorus determinations in fluoride- and silica-rich geothermal waters.," *Environmental science & technology*, vol. 14, no. 12, pp. 1475–81, 1980.
- [206] A. C. Heimann and R. Jakobsen, "Filtration through nylon membranes negatively affects analysis of arsenic and phosphate by the molybdenum blue method," *Talanta*, vol. 72, no. 2, pp. 839–841, 2007.
- [207] K. Higuchi, H. Tamanouchi, and S. Motomizu, "On-Line photo-Oxidative Decomposition of Phosphorus Compounds to Orthophosphate and Its Application to Flow Injection Spectrophotometric Determinations of Total Phosphorus in River and Waste Waters.," *Analytical Sciences*, vol. 14, no. 5, pp. 941–946, 1998.
- [208] D. Nyamah and J. O. Torgbor, "Colorimetric method for the determination of arsenic in potable water," *Water Research*, vol. 20, no. 11, pp. 1341–1344, 1986.
- [209] K. Morita and E. Kaneko, "Spectrophotometric Determination of Arsenic in Water Samples Based on Micro Particle Formation of Ethyl Violet –," *Japan Society for Analytical Chemistry*, vol. 22, no. August, pp. 1085–1089, 2006.

- [210] G. R. Bartlett, "Phosphorus assay in column chromatography," *Journal of Biological Chemistry*, vol. 234, no. 3, pp. 466–468, 1959.
- [211] A. Hashem, M. Takaki, T. Jodai, and K. Toda, "Measurements of arsenite and arsenate contained in mining river waters and leached from contaminated sediments by sequential hydride generation flow injection analysis," *Talanta*, vol. 84, no. 5, pp. 1336–1341, 2011.
- [212] K. Toda, T. Ohba, and M. Takaki, "Speciation-Capable Field Instrument for the Measurement of Arsenite and Arsenate in Water," *Analytical Chemistry*, vol. 77, no. 15, pp. 4765–4773, 2005.
- [213] S. Motomizu, M. Oshima, and A. Hirashima, "Spectrophotometric determination of phosphorus in river water based on the reaction of vanadomolybdophosphate with malachite green," *Analytica Chimica Acta*, vol. 211, pp. 119–127, 1988.
- [214] D. Melamed, "Monitoring arsenic in the environment: A review of science and technologies with the potential for field measurements," *Analytica Chimica Acta*, vol. 532, no. 1, pp. 1–13, 2005.
- [215] S. Sorlini and F. Gialdini, "Conventional oxidation treatments for the removal of arsenic with chlorine dioxide, hypochlorite, potassium permanganate and monochloramine," *Water Research*, vol. 44, no. 19, pp. 5653–5659, 2010.
- [216] N. Li, M. Fan, J. Van Leeuwen, B. Saha, H. Yang, and C. P. Huang, "Oxidation of As(III) by potassium permanganate," *Journal of Environmental Sciences*, vol. 19, no. 7, pp. 783–786, 2007.
- [217] G. Lee, K. Song, and J. Bae, "Permanganate oxidation of arsenic (III): Reaction stoichiometry and the characterization of solid product," *Geochimica et Cosmochimica Acta*, vol. 75, no. 17, pp. 4713–4727, 2011.
- [218] B. Serpaud and J. Bollinger, "Arsenite oxidation and arsenate determination by the molybdene blue method," *Talanta*, vol. 61, pp. 267–276, 2003.
- [219] K. L. Linge and C. E. Oldham, "Interference from arsenate when determining phosphate by the malachite green spectrophotometric method," *Analytica Chimica Acta*, vol. 450, no. 1–2, pp. 247–252, 2001.

- [220] D. L. Johnson, "Simultaneous determination of arsenate and phosphate in natural waters," *Environmental science & technology*, vol. 5, no. 5, pp. 411–414, 1971.
- [221] B. Chang, Y. Chen, and H. Yak, "Determination of Arsenic Species by Ion Chromatography with Chemiluminescence Detection," *International Journal of Chemical and Environmental Engineering*, vol. 3, no. 6, pp. 3–5, 2012.
- [222] S. Saha and P. Sarkar, "Differential pulse anodic stripping voltammetry for detection of As (III) by Chitosan-Fe(OH)₃ modified glassy carbon electrode: A new approach towards speciation of arsenic," *Talanta*, vol. 158, pp. 235–245, 2016.
- [223] D. K. Nordstrom, "Worldwide occurrences of arsenic in groundwater," *Science*, vol. 296, pp. 2143–2145, 2002.
- [224] C. Cerveira, D. Pozebon, D. P. de Moraes, and J. C. Silva de Fraga, "Speciation of inorganic arsenic in rice using hydride generation atomic absorption spectrometry (HG-AAS)," *Analytical Methods*, vol. 7, no. 11, pp. 4528–4534, 2015.
- [225] L. K. Tan and J. E. Dutrizac, "Determination of arsenic(III) and arsenic(V) in ferric chloride-hydrochloric acid leaching media by ion chromatography," *Analytical Chemistry*, vol. 57, no. 6, pp. 1027–1032, 1985.
- [226] A. B. Mukherjee and P. Bhattacharya, "Arsenic in groundwater in the Bengal Delta Plain: slow poisoning in Bangladesh," *Environmental Reviews*, vol. 220, pp. 189–220, 2001.
- [227] H. Kitami and Y. Ishihara, "Simultaneous Determination of Total Nitrogen and Total Phosphorus in Sewage Water by Flow Injection Analysis after Continuous Flow Sample Degradation Using Heating Aluminum Block," *Bunseki Kagaku*, vol. 58, no. 9, p. P 839-844, 2009.
- [228] V. Lenoble, V. Deluchat, B. Serpaud, and J. C. Bollinger, "Arsenite oxidation and arsenate determination by the molybdene blue method," *Talanta*, vol. 61, no. 3, pp. 267–276, 2003.

- [229] P. Mondal, S. Bhowmick, D. Chatterjee, A. Figoli, and B. Van der Bruggen, "Remediation of inorganic arsenic in groundwater for safe water supply: A critical assessment of technological solutions," *Chemosphere*, vol. 92, no. 2, pp. 157–170, 2013.
- [230] P. K. Dasgupta, H. Huang, and G. Zhang, "Photometric Measurement of Trace As (III) and As (V) in Drinking Water," *Talanta*, vol. 58, no. III, pp. 153–164, 2002.
- [231] S. Vasudevan, S. Mohan, and G. Sozhan, "Studies on the Oxidation of As (III) to As (V) by In-Situ-Generated Hypochlorite," *Industrial & engineering chemistry research*, vol. 45, pp. 7729–7732, 2006.
- [232] M. Zaw and M. T. Emmett, "Arsenic removal from water using advanced oxidation processes," *Toxicology Letters*, vol. 133, pp. 113–118, 2002.
- [233] L. I. Na, F. A. N. Maohong, V. L. Johannes, S. Basudeb, Y. Hongqun, and C. P. Huang, "Oxidation of As (III) by potassium permanganate," *Journal of Environmental Sciences*, vol. 19, pp. 783–786, 2007.
- [234] G. Lee, K. Song, and J. Bae, "Permanganate oxidation of arsenic(III): Reaction stoichiometry and the characterization of solid product," *Geochimica et Cosmochimica Acta*, vol. 75, no. 17, pp. 4713–4727, 2011.
- [235] M. I. Ikedo, M. M. Mori, K. K. Kurachi, H. U. Wenzhi, and K. T. Tanaka, "Selective and Simultaneous Determination of Phosphate and Silicate Ions in Leaching Process Waters for Ceramics Glaze Raw Materials of Natural Origin by Ion-Exclusion Chromatography Coupled with UV-detection after Postcolumn Derivatization," *The International Journal of the Japan Society for Analytical Chemistry*, vol. 22, no. January, pp. 117–121, 2006.
- [236] W. Z. W. Yaacob, nur S. M. Pauzi, and hazwani Abdul Mutalib, "Acid mine drainage and heavy metals contamination at abandoned and active mine sites in Pahang," *Bulletin of the Geological Society of Malaysia*, vol. 55, pp. 15–20, 2009.
- [237] V. K. Gupta, M. Gupta, and S. Sharma, "Process development for the removal

- of lead and chromium from aqueous solutions using red mud - an aluminium industry waste,” *Water Research*, vol. 35, no. 5, pp. 1125–1134, 2001.
- [238] E. E. Adomako, C. Deacon, and A. a. Meharg, “Variations in Concentrations of Arsenic and Other Potentially Toxic Elements in Mine and Paddy Soils and Irrigation Waters from Southern Ghana,” *Water Quality, Exposure and Health*, vol. 2, pp. 115–124, 2010.
- [239] R. K. Rowe and M. S. Hosney, “Laboratory investigation of GCL performance for covering arsenic contaminated mine wastes,” *Geotextiles and Geomembranes*, vol. 39, pp. 63–77, 2013.
- [240] R. V. Kumar, A. K. Basumatary, A. K. Ghoshal, and G. Pugazhenthii, “Performance assessment of an analcime-C zeolite–ceramic composite membrane by removal of Cr(vi) from aqueous solution,” *The Royal Society of Chemistry*, vol. 5, pp. 6246–6254, 2015.
- [241] P. Manual, *DR5000 Spectrophotometer Procedures Manuel*. 2005.
- [242] British Standard Methods, *Methods of test for Soils for civil engineering purposes*, no. 1. London: British Standard Institution, 1990.
- [243] M. S. Ko, J. Y. Kim, H. S. Park, and K. W. Kim, “Field assessment of arsenic immobilization in soil amended with iron rich acid mine drainage sludge,” *Journal of Cleaner Production*, vol. 108, pp. 1–8, 2015.
- [244] G. Cavey and D. R. Gunning, “Summary Report on the Buffalo Reef pProject Pahang State, Malaysia for Monument Mining Limited,” 2007.
- [245] J. Li, B. Dabrowski, J. D. Miller, S. Acar, M. Dietrich, K. M. LeVier, and R. Y. Wan, “The influence of pyrite pre-oxidation on gold recovery by cyanidation,” *Minerals Engineering*, vol. 19, no. 9, pp. 883–895, 2006.
- [246] D. Paktunc, A. Foster, S. Heald, and G. Laflamme, “Speciation and characterization of arsenic in gold ores and cyanidation tailings using X-ray absorption spectroscopy,” *Geochimica et Cosmochimica Acta*, vol. 68, no. 5, pp. 969–983, 2004.
- [247] V. Kyoseva, E. Todorova, and I. Dombalov, “Comparative Assessment of the Methods for Destruction of Cyanides Used in Gold Mining Industry,” *Journal*

- of the university of chemical technology and metallurgy*, vol. 44, no. 4, pp. 403–408, 2010.
- [248] W. Douglas Gould, M. King, B. R. Mohapatra, R. A. Cameron, A. Kapoor, and D. W. Koren, “A critical review on destruction of thiocyanate in mining effluents,” *Minerals Engineering*, vol. 34, pp. 38–47, 2012.
- [249] C. B. Tabelin, A. Hashimoto, T. Igarashi, and T. Yoneda, “Leaching of boron , arsenic and selenium from sedimentary rocks : II . pH dependence , speciation and mechanisms of release,” *Science of the Total Environment*, vol. 473–474, pp. 244–253, 2014.
- [250] S. M. Rao and B. V. V. Reddy, “Characterization of Kolar gold field mine tailings for cyanide and acid drainage,” *Geotechnical and Geological Engineering*, vol. 24, no. 6, pp. 1545–1559, 2006.
- [251] C. B. Tabelin, A. Hashimoto, T. Igarashi, and T. Yoneda, “Leaching of boron, arsenic and selenium from sedimentary rocks: II. pH dependence, speciation and mechanisms of release,” *Science of the Total Environment*, vol. 473–474, pp. 244–253, 2014.
- [252] “Dutch list The Ministry of Housing, Spatial Planning and Environment, Department of Soil Protection.” [Online]. Available: <http://www.contaminatedland.co.uk/std-guid/dutch-l.htm>.
- [253] I. Kelly, “Guidelines for Contaminated Soils specifically developed for gasworks sites in London.” Society of the Chemical Industry, London, 1980.
- [254] CCME, “Canadian Soil Quality Guidelines for the Protection of Environmental and Human Health,” *Canadian Council of Ministers of the Environment*, pp. 1–6, 2007.
- [255] ICRCL, “Guidance on the Assessment and Redevelopment of Contaminated Land,” Guidance Note 59/83 2nd ed., 1987.
- [256] T. Igarashi, H. Imagawa, H. Uchiyama, and K. Asakura, “Leaching behavior of arsenic from various rocks by controlling geochemical conditions,” *Minerals Engineering*, vol. 21, no. 3, pp. 191–199, 2008.

- [257] L. Wang and D. E. Giammar, "Effects of pH, dissolved oxygen, and aqueous ferrous iron on the adsorption of arsenic to lepidocrocite," *Journal of Colloid and Interface Science*, vol. 448, pp. 331–338, 2015.
- [258] C. Roussel, H. Bril, and A. Fernandez, "Evolution of sulphides-rich mine tailings immobilization of As and Fe," *Comptes Rendus de l'Académie Des Sciences-Series IIA-Earth and Planetary Science*, vol. 329, no. 11, pp. 787–794, 1999.
- [259] N. Singh, S. Gupta, N. Marwa, V. Pandey, P. C. Verma, S. Rathaur, and N. Singh, "Arsenic mediated modifications in *Bacillus aryabhatai* and their biotechnological applications for arsenic bioremediation," *Chemosphere*, vol. 164, pp. 524–534, 2016.
- [260] L. Cumbal and A. K. Sengupta, "Arsenic removal using polymer-supported hydrated iron(III) oxide nanoparticles: Role of Donnan membrane effect," *Environmental Science and Technology*, vol. 39, no. 17, pp. 6508–6515, 2005.
- [261] L. Zeng, "Arsenic adsorption from aqueous solutions on an Fe (III)-Si binary oxide adsorbent," *Water quality research journal of canada*, vol. 39, no. 3, pp. 267–275, 2004.
- [262] J. H. Park, Y. Han, and J. S. Ahn, "Comparison of arsenic co-precipitation and adsorption by iron minerals and the mechanism of arsenic natural attenuation in a mine stream," *Water Research*, vol. 106, pp. 295–303, 2016.
- [263] S. Tresintsi, K. Simeonidis, I. Georgiou, G. Vourlias, G. Stavropoulos, M. Mitrakas, and I. W. A. S. Conferences, "Effect of precipitation pH on arsenic removal by single-Fe and binary-Fe / Mn oxy-hydroxides," pp. 1–9, 2011.
- [264] S. Bakardjieva, V. Štengl, J. Šubrt, and E. Večerníková, "Characteristic of hydrous iron (III) oxides prepared by homogeneous precipitation of iron (III) sulphate with urea," *Solid State Sciences*, vol. 7, no. 4, pp. 367–374, 2005.
- [265] D. Pokhrel and T. Viraraghavan, "Arsenic removal from an aqueous solution by modified *A. niger* biomass: Batch kinetic and isotherm studies," *Journal of Hazardous Materials*, vol. 150, no. 3, pp. 818–825, 2008.
- [266] B. Chen, Z. Zhu, Y. Guo, Y. Qiu, and J. Zhao, "Facile synthesis of

- mesoporous Ce-Fe bimetal oxide and its enhanced adsorption of arsenate from aqueous solutions,” *Journal of Colloid and Interface Science*, vol. 398, pp. 142–151, 2013.
- [267] S. Kong, Y. Wang, H. Zhan, S. Yuan, and Q. Hu, “Adsorption Mechanism of Humic Acid on Cu/Fe Bimetallic Particles and Its Influence on the Reduction of Nitrobenzene in Groundwater,” *Water, Air, Soil Pollution*, vol. 225, no. 11, p. 1985, 2014.
- [268] M. Gui and D. Bhattacharyya, “Iron and Iron Oxide Functionalized Membranes With Applications To Selected Chloro-Organic and Metal Removal From Water,” University of Kentucky, 2014.
- [269] A. Tiraferri, K. L. Chen, R. Sethi, and M. Elimelech, “Reduced aggregation and sedimentation of zero-valent iron nanoparticles in the presence of guar gum,” *Journal of Colloid and Interface Science*, vol. 324, no. 1–2, pp. 71–79, 2008.
- [270] E. M. Hotze, T. Phenrat, and G. V. Lowry, “Nanoparticle aggregation: challenges to understanding transport and reactivity in the environment,” *Journal of Environmental Quality*, vol. 39, pp. 1909–1924, 2010.
- [271] G. Cherr, B. J. Cardinale, R. Miller, and Z. Ji, “Stability and Aggregation of Metal Oxide Nanoparticles in Natural Aqueous Matrices,” *Environmental science & technology*, vol. 44, no. 6, pp. 1962–1967, 2010.
- [272] J. Yuan, T. Zhou, and H. Pu, “Nano-sized silica hollow spheres: Preparation, mechanism analysis and its water retention property,” *Journal of Physics and Chemistry of Solids*, vol. 71, no. 7, pp. 1013–1019, 2010.
- [273] F. Rouquerol, J. Rouquerol, K. S. W. Sing, P. Llewellyn, and G. Maurin, *Adsorption by powders and porous solids Principles, Methodology and Applications*, Second edi. Oxford, UK: Elsevier, 2014.
- [274] M. Thommes, K. Kaneko, A. V. Neimark, J. P. Olivier, F. Rodriguez-Reinoso, J. Rouquerol, and K. S. W. Sing, “Physisorption of gases, with special reference to the evaluation of surface area and pore size distribution (IUPAC Technical Report),” *Pure and Applied Chemistry*, vol. 87, no. 9–10, pp. 1051–

1069, 2015.

- [275] T. Liu, Y. Yang, Z.-L. Wang, and Y. Sun, "Remediation of arsenic(III) from aqueous solutions using improved nanoscale zero-valent iron on pumice," *Chemical Engineering Journal*, vol. 288, pp. 739–744, 2016.
- [276] M. M. Vuruna and N. B. E. N. Issa, "Equilibrium, kinetic and thermodynamic studies on removal of Cd (II), Pb (II) and As (V) from wastewater using carp (cyprinus carpio) Scales," *Digest Journal of Nanomaterials and Biostructures*, vol. 8, no. 4, pp. 1581–1590, 2013.
- [277] S. G. Johnston, E. D. Burton, A. F. Keene, B. Planer-friedrich, A. Voegelin, M. G. Blackford, and G. R. Lumpkin, "Arsenic mobilization and iron transformations during sulfidization of As (V)-bearing jarosite," *Chemical Geology*, vol. 334, pp. 9–24, 2012.
- [278] G. P. Jeppu and T. P. Clement, "A modified Langmuir-Freundlich isotherm model for simulating pH-dependent adsorption effects," *Journal of Contaminant Hydrology*, vol. 129–130, pp. 46–53, 2012.
- [279] S. Bagherifam, S. Komarneni, A. Lakzian, A. Fotovat, and R. Khorasani, "Applied Clay Science Evaluation of Zn – Al – SO₄ layered double hydroxide for the removal of arsenite and arsenate from a simulated soil solution: Isotherms and kinetics," *Applied Clay Science*, vol. 95, pp. 119–125, 2014.
- [280] D. Tchoubar, J. Y. Bottero, P. Quienne, and M. Arnaud, "Partial hydrolysis of ferric chloride salt. Structural investigation by photon-correlation spectroscopy and small-angle x-ray scattering," *Langmuir*, vol. 7, no. 2, pp. 398–402, 1991.
- [281] C. C. Fuller, J. A. Dadis, and G. A. Waychunas, "Surface chemistry of ferrihydrite: Part 2. Kinetics of arsenate adsorption and coprecipitation," *Geochimica et Cosmochimica Acta*, vol. 57, no. 10, pp. 2271–2282, 1993.
- [282] G. A. Waychunas, C. C. Fuller, B. A. Rea, and J. A. Davis, "Wide angle X-ray scattering (WAXS) study of 'two-line' ferrihydrite structure: Effect of arsenate sorption and counterion variation and comparison with EXAFS results," *Geochimica et Cosmochimica Acta*, vol. 60, no. 10, pp. 1765–1781,

1996.

- [283] M. Mosaferi, S. Nemati, A. Khataee, S. Nasser, and A. A. Hashemi, "Removal of arsenic (III , V) from aqueous solution by nanoscale zero-valent iron stabilized with Starch and Carboxymethyl cellulose," *Journal of Environmental Health Science and Engineering* , vol. 12, no. 1, pp. 1–11, 2014.
- [284] S. R. Kanel, D. Ph, A. M. Asce, and H. Choi, "Removal of Arsenic from Groundwater by Industrial Byproducts and Its Comparison with Zero-Valent Iron Removal of Arsenic from Groundwater by Industrial Byproducts and Its Comparison with Zero-Valent Iron," *Journal of Hazardous, Toxic, and Radioactive Waste*, no. November, p. 4016028, 2016.
- [285] Z. Veličković, G. D. Vuković, A. D. Marinković, M. S. Moldovan, A. A. Perić-Grujić, P. S. Uskoković, and M. D. Ristić, "Adsorption of arsenate on iron(III) oxide coated ethylenediamine functionalized multiwall carbon nanotubes," *Chemical Engineering Journal*, vol. 181–182, pp. 174–181, 2012.
- [286] D. Kołodyńska, M. Kowalczyk, and M. Gęca, "Hybrid sorbents in metal ions removal from water and wastewater," *CHEMIK*, vol. 66, no. 11, pp. 1249–1258, 2012.
- [287] C. B. Tabelin, T. Igarashi, and R. Takahashi, "Mobilization and speciation of arsenic from hydrothermally altered rock in laboratory column experiments under ambient conditions," *Applied Geochemistry*, vol. 27, no. 1, pp. 326–342, 2012.
- [288] S. M. Nam, M. Kim, S. Hyun, and S. H. Lee, "Chemical attenuation of arsenic by soils across two abandoned mine sites in Korea," *Chemosphere*, vol. 81, no. 9, pp. 1124–1130, 2010.
- [289] R. Larios, R. Fernández-Martínez, R. Álvarez, and I. Rucandio, "Arsenic pollution and fractionation in sediments and mine waste samples from different mine sites," *Science of the Total Environment*, vol. 431, pp. 426–435, 2012.

- [290] K. Fukushi, M. Sasaki, T. Sato, N. Yanase, H. Amano, and H. Ikeda, "A natural attenuation of arsenic in drainage from an abandoned arsenic mine dump," *Applied Geochemistry*, vol. 18, no. 8, pp. 1267–1278, 2003.
- [291] S. Wang and C. N. Mulligan, "Natural attenuation processes for remediation of arsenic contaminated soils and groundwater," *Journal of Hazardous Materials*, vol. 138, pp. 459–470, 2006.
- [292] C. Baltazar, T. Igarashi, T. Yoneda, and S. Tamamura, "Utilization of natural and artificial adsorbents in the mitigation of arsenic leached from hydrothermally altered rock," *Engineering Geology*, vol. 156, pp. 58–67, 2013.
- [293] A. Rubo, R. Kellens, J. Reddy, N. Steier, and W. Hasenpusch, "Alkali Metal Cyanides," in *Ullmann's Encyclopedia of Industrial Chemistry*,. Wiley-VCH Verlag GmbH & Co. KGaA, 2000.
- [294] E. C. F. Ecotoxicology and T. of Chemicals, *Cyanides of Hydrogen, Sodium and Potassium, and Acetone Cyanohydrin (CAS No. 74-90-8, 143-33-9, 151-50-8 and 75-86-5)*, no. 74. ECETOC, 2007.
- [295] K. IJIMA, E. YOSHIKAWA, and S. FURUTA, "Adsorptive Properties of 'ECOMEL TM ' 53NJ for Heavy Metal Compounds," 2015.
- [296] "Iron Powder ' Ecomel ' for Purification of Heavy Metal-contaminated Soil and Groundwater," *Kobe Steel, Ltd.* [Online]. Available: www.kankeiren.or.jp/kankyou/en/pdf/en137.