

SPATIOTEMPORAL MODEL FOR LANDSCAPE ECOLOGICAL
ASSESSMENT IN LANDSCAPE PLANNING

WAN YUSRYZAL BIN WAN IBRAHIM

A thesis submitted in fulfilment of the
requirements for the award of the degree of
Doctor of Philosophy (Urban and Regional Planning)

Faculty of Built Environment
Universiti Teknologi Malaysia

JANUARY 2018

Specially dedicated to all.

Al-Fatihah

ACKNOWLEDGEMENT

Alhamdulillah, I am grateful to the Allah S.W.T for the good health and wellbeing that were necessary to complete this PhD study. Peace and blessing of Allah be upon the prophet Muhammad (S.A.W), his family, companions and all those who follow his footsteps, Ameen.

I wish to express my sincere thanks to all that supporting me during my study. Firstly, I would like to express my sincere gratitude to my supervisor Prof. Dato' Dr. Ahmad Nazri bin Muhamad Ludin for the continuous support of my Ph.D study for his patience, motivation, and immense knowledge. I thank him for sharing his valuable time and for giving helpful information to complete this PhD program. His guidance helped me in all the time. Similarly, I am grateful to my family members, especially my parent for their upbringing, prayers and encouragement. I have to thank my wife and my children for their patience, understanding and support during this program. My appreciation also goes to my family members, colleagues and friends for their support and encouragement during the program. I thank you very much. May the almighty reward you all with prosperity in this life and endless bliss hereafter.

ABSTRACT

Ecological landscape area is rich with biodiversity and ecosystem are two important factors that balance the serenity of the environment through its ecological function and services. However, landscape change especially rapid urbanization has led to extensive land use and land cover (LULC) transformation that degrades the ecological landscape area and ecosystem services. The limitation of integration analysis in LULC change with ecological interaction has caused detrimental impact on natural landscape area and environmental quality. Analysing the spatiotemporal characteristics of landscape changes and ecological response in a multidisciplinary research is necessary to extend the understanding of spatial change behaviour and ecological consequences. Thus, the aim of this research is to study the integration of spatiotemporal dimension of landscape change with ecological landscape sensitivity consideration in Iskandar Malaysia region (Johor Bahru). The spatiotemporal dimension of historical and future LULC change is analysed to identify the direction and characteristics of the landscape structure and function change. Logistic regression model, analytical hierarchical process, markov chain model and cellular automata were used to identify the spatiotemporal LULC change in the study area. A series of landscape matrices in landscape index at class and landscape levels were used to analyse the spatiotemporal dimension of the landscape change pattern. It includes measurement of the ecological integrity and function responses towards spatiotemporal landscape change by using Core Area Model. Satellite images of 1994, 2000, 2007 and 2013 were used to understand the historical landscape changes and as a basis for future projection. Geographic Information System and Remote Sensing were utilized to evaluate the temporal landscape characteristics and spatial pattern changes. The results indicate that rapid urbanization of Iskandar Malaysia region from 2007 to 2013 has substantially changed the structure and function of the ecological area. The urban area significantly increased from 8,031.6 hectares (3.84%) in 1994 to 42,972.94 (20.1%) in 2013, and expected to increase to 112,224.6 hectares (53.59%) in 2030. As a consequence, the natural ecological areas reduced from 55,201.77 hectares (26.37%) in 1994 to 19,011.5 hectares (9.08%) in 2013. Due to the landscape mosaic change, the core ecological areas are affected from 21,465.9 hectares (38%) reduced to 9,317.61 hectares (49%) and expected to further reduce at 8,416.71 hectares (41%) in 1994, 2013 and 2030, respectively. It shows the response of ecological condition in natural landscape areas towards the landscape changes which subsequently disturb the ecological values and services. As a conclusion, the findings of this research could provide decision makers with better understanding on the environmental consequences of the landscape changes. In addition, it contributes to enhancement of methods in multidisciplinary research and finally increases the capability of the process in adaptive management for the spatiotemporal landscape change.

ABSTRAK

Kawasan landskap ekologi yang kaya dengan kepelbagaian biologi dan ekosistem adalah dua faktor penting untuk mengimbangi kedamaian alam sekitar melalui fungsi dan perkhidmatan ekologi. Walau bagaimanapun, perubahan landskap terutama proses pembandaran yang pesat telah membawa kepada perubahan penggunaan dan liputan tanah (*LULC*) yang besar dan merosakkan kawasan landskap ekologi dan perkhidmatan ekosistem. Keterbatasan analisis dalam mengintegrasikan analisis *LULC* dengan interaksi ekologi telah menyebabkan kesan buruk kepada kawasan landskap semulajadi dan kualiti alam sekitar. Analisis ciri-ciri perubahan ruwang (*spatiotemporal*) landskap dan tindak balas ekologi dalam penyelidikan pelbagai disiplin adalah penting untuk memahami secara mendalam tentang corak perubahan ruwang dan kesan kepada ekologi. Dengan itu, tujuan kajian ini adalah untuk mengintegrasikan perubahan dimensi ruwang landskap dengan pertimbangan sensitiviti landskap ekologi di dalam wilayah Iskandar Malaysia (Johor Bahru). Dimensi ruwang terdahulu dan perubahan *LULC* masa depan dianalisis untuk mengenalpasti arah perubahan ciri-ciri struktur landskap dan perubahan fungsinya. *Logistic regression model*, *analytical heirarchical process (AHP)*, *markov chain (MC)* dan *cellular automata (CA)* telah digunakan untuk mengenal pasti perubahan *LULC* di kawasan kajian. Satu siri landskap matrik di dalam indeks landskap di peringkat kelas dan landskap telah digunakan untuk menganalisis dimensi ruwang bagi corak perubahan landskap tersebut. Ia termasuk pengukuran tindak balas integriti dan fungsi ekologi terhadap perubahan landskap ruwang dengan menggunakan *Core Area Model*. Imej satelit pada tahun 1994, 2000, 2007 dan 2013 telah digunakan untuk memahami perubahan landskap terdahulu dan sebagai asas untuk unjuran perubahan pada masa depan. Sistem maklumat geografi (*GIS*) dan penderiaan jauh (*Remote Sensing*) telah digunakan untuk menilai perubahan ciri-ciri landskap dan corak perubahan ruwang. Hasil kajian menunjukkan bahawa proses pembandaran pesat wilayah Iskandar Malaysia dari 2007 hingga 2013 telah mengubah struktur dan fungsi kawasan ekologi dengan ketara. Keluasan kawasan bandar meningkat dengan ketara daripada 8,031.6 hektar (3.84%) pada tahun 1994 kepada 42,972.94 (20.1%) pada tahun 2013 dan dijangka mencapai 112,224.6 hektar (53.59%) pada tahun 2030. Akibatnya kawasan landskap ekologi semulajadi telah berkurang daripada 55,201.77 (26.37%) pada tahun 1994 kepada 19,011.5 (9.08%) pada tahun 2013. Disebabkan oleh perubahan mozek landskap, kawasan teras ekologi semulajadi (*core area*) telah terjejas daripada 21,465.9 hektar (38%) berkurangan kepada 9,317.61 hektar (49%), dan akan terus berkurang kepada 8,416.71 hektar (41%) pada 1994, 2013 dan 2030 tersebut. Ia menunjukkan tindak balas keadaan ekologi bagi kawasan-kawasan landskap semulajadi terhadap perubahan persekitaran yang seterusnya mengganggu nilai-nilai dan perkhidmatan ekologi. Kesimpulannya, hasil kajian ini dapat menyediakan pemahaman yang lebih baik kepada pembuat keputusan mengenai kesan alam sekitar daripada perubahan landskap. Di samping itu, ia menyumbang kepada memperkukuhkan kaedah-kaedah penyelidikan yang melibatkan kepelbagaian disiplin dan akhirnya meningkatkan keupayaan proses penyesuaian pengurusan untuk perubahan landskap ruwang.

TABLE OF CONTENTS

CHAPTER	TITLE	PAGE
	DECLARATION	ii
	DEDICATION	iii
	ACKNOWLEDGEMENTS	iv
	ABSTRACT	v
	ABSTRAK	vi
	TABLE OF CONTENTS	vii
	LIST OF TABLES	xiv
	LIST OF FIGURES	xvii
	LIST OF APPENDICES	xxiv
	LIST OF ACRONYMS	xxv
1	INTRODUCTION	1
	1.0 Introduction	1
	1.1 Research Backgroud	4
	1.2 Problem Statement	7
	1.3 Research Questions	9
	1.4 Research Aim and Objectives	10
	1.5 Significance of Research	11
	1.6 Scope of Research	11
	1.7 Research Outline	13
	1.8 Thesis Organization	18
2	ECOLOGICAL IMPORTANCE AND SPATIOTEMPORAL LANDSCAPE CHANGE	21

2.0	Introduction	21
2.1	Landscape Ecological Resource: Theories, Function and Application In Landscape Planning	22
2.1.1	Landscape Ecological Resource Definition and Ecological Interaction	22
2.1.2	The Significance of Landscape Ecological Function	25
2.1.3	The Role of Landscape Structure For Ecological Function Integrity	27
2.1.4	Ecological Landscape Consideration In Landscape Planning Theory	32
2.2	Landscape Change and Ecological Consequence	36
2.2.1	Landscape Change; Urbanization and Agriculture	37
2.2.2	Ecological Impact On Landscape Change	39
2.2.2.1	Fragmentation and Homogenisation of Natural Landscape Areas	40
2.2.2.2	Natural Landscape Area Change in Malaysia	44
2.2.2.3	Edge Effect; Reduction of Core Natural Area Quantity, Quality and Habitat Suitability	48
2.3	Summary	55
3	SPATIAL DYNAMIC MODELLING AND APPLICATION IN LANDSCAPE CHANGE	57
3.0	Introduction	57
3.1	Ecological Analysis in Spatiotemporal Landscape Change	58
3.2	Spatial Dynamic Modelling in Landscape Ecological Consideration	62
3.2.1	Ecological Modelling In Landscape Planning	64

3.2.2	Spatial Landscape Ecology Modelling and GIS Application	70
3.2.3	Spatial Dynamic Modelling In Landscape Ecology Application	74
3.2.3.1	Overview of Spatial Models	76
3.2.3.2	Markov Chain Model	77
3.2.3.3	Multi-criteria Evaluation	78
3.2.3.4	CA_Markov Model	79
3.3	Incorporating of Ecological Assessment Tool in Spatiotemporal Landscape Change	80
3.3.1	Landscape Index	80
3.3.2	Data Acquisition in Ecological Landscape	82
3.3.3	Integrated Consideration on Landscape Ecology in Landscape Change	83
3.4	Summary	84
4	METHODOLOGY AND SPATIAL MODEL DEVELOPMENT	85
4.0	Introduction	85
4.1	Model Design and Development	86
4.2	Landscape Change Identification	91
4.3	Prediction of Landscape Change Probabilities	92
4.3.1	Land Use and Land Cover Change Spatial Relationship	93
4.3.2	Future Landscape Change Probability	96
4.3.3	Identification of Rules and Probability of Area Changes	99
4.4	Spatiotemporal Landscape Change Simulation	101
4.4.1	Cellular Automata Based Simulation	101
4.4.2	Validation of Rules and Simulation	103
4.5	Landscape Ecological Assessment	104
4.5.1	Configuration of Ecological Landscape	106

	Analysis	
4.5.2	Composition Landscape Ecological Analysis	108
4.5.3	Core Ecological Area Analysis	109
4.6	Selection of Study Area	111
4.6.1	Population, Density and Ethnicity	113
4.6.2	Urbanization and LULC in Iskandar Malaysia	116
4.6.3	Ecological Planning Framework in Development of Iskandar Malaysia	123
4.7	Database Design and Data Preparation	128
4.7.1	Database design	128
4.7.1.1	Land Use and Land Cover	130
4.7.1.2	Physical characteristics	130
4.7.1.3	Proximity to Land Use, Infrastructure and Facilities	131
4.7.1.4	Strategy and policy dimension	132
4.7.2	Data collection, Database development and Data Processing	132
4.7.2.1	Land Use and Land Cover Map Preparation (Satellite Image Processing)	134
4.7.2.2	Geophysical Database Development	138
4.7.2.3	Urban and Natural Landscape Change	139
4.7.2.4	Proximity of Variables	140
4.7.2.5	Delineation of Strategy and Policy	141
4.8	Summary	142
5	SPATIOTEMPORAL LANDSCAPE CHANGE IN ISKANDAR MALAYSIA	144
5.0	Introduction	144
5.1	Spatiotemporal Historical Land Use and Land Cover Change	145

5.1.1	Performance of LULC Classification and LULC Composition	145
5.1.2	LULC Change from 1994 to 2000	147
5.1.3	LULC Change from 2000 to 2007	151
5.1.4	LULC Change from 2007 to 2013	154
5.2	LULC Change Behaviour and Characteristics	156
5.2.1	Scenario of Characteristics in LULC Change	156
5.2.2	Landscape Change Behaviour and Magnitude	163
5.2.2.1	Urban Landscape Change Correlation	163
5.2.2.2	Natural Landscape Change Correlation	167
5.3	Projection of LULC Change	169
5.3.1	Transitional of LULC in Future Change	170
5.3.2	Suitability Area Identification for Future Landscape Change	174
5.4	Spatiotemporal of Future Landscape Change	177
5.4.1	Future LULC between 2013 and 2020	178
5.4.2	Future LULC between 2020 and 2025	182
5.4.3	Future LULC between 2025 and 2030	185
5.4.4	Overall Spatiotemporal of future landscape change in Iskandar Malaysia from 2013 to 2030	188
5.5	Summary	190
6	ECOLOGICAL LANDSCAPE CONSEQUENCE IN SPATIOTEMPORAL LULC CHANGE	192
6.0	Introduction	191
6.1	Spatiotemporal Analysis of Landscape Change Pattern	193
6.1.1	LULC Change Analysis	193
6.1.2	Urban Growth Pattern Analysis	195
6.1.3	Urban Growth Characteristics in Different Development Zones	200

6.2	Spatiotemporal of Ecological Landscape Structure Change	206
6.2.1	Configuration of Ecological Landscape Change	210
6.2.1.1	Intertidal forest areas landscape structure change between 1994 and 2030	215
6.2.1.2	Inland forest landscape change between 1994 and 2030	224
6.3	Edge Effect and Ecological Function Evaluation	227
6.3.1	Depth of Influence and Ecological Edge Response	228
6.3.2	Core Ecological Area Degradation	233
6.3.2.1	Intertidal Forest Area Implication	238
6.3.2.2	Inland Forest Area Implication	242
6.4	Summary	243
7	DISCUSSION AND CONCLUSION	245
7.0	Introduction	245
7.1	The Implication of Development Plan Dimension Towards Spatiotemporal Landscape Change	246
7.2	Ecological Structure and Function Change	250
7.2.1	Ecological Value Destruction	250
7.2.2	Ecological Service Degradation	253
7.2.3	Edge Effect and Core Ecological Area	255
7.3	Achievement and Contribution of the Study	257
7.3.1	Integration of Spatiotemporal LULC in Landscape Ecology	258
7.3.2	Enhancement of GIS Model Capability in Landscape Change Analysis	261
7.3.3	Statistical Method in Quantifying Landscape Change and Ecological Assessment	263
7.3.4	Strengthening Multidisciplinary Approach in	265

	Adaptive Management	
7.4	Limitation of the Study and Future Research	269
7.5	Improvement and Future Research	271
7.6	Conclusion	272
	REFERENCES	274
	Appendices A – F	295-323

LIST OF TABLES

TABLE NO.	TITLE	PAGE
2.1	Examples of selected studies on spatiotemporal landscape change implication due to urban expansion and agriculture activity at the different countries	42
2.2	Selected studies of land use and land cover change and natural landscape degradation at regional and local levels in Malaysia	46
3.1	Basic type of models	66
3.2	Ecological models and description of functions	67
3.3	Selected studies on spatial dynamic modelling in landscape planning with ecological consideration	75
3.4	Matrix table of criterion-alternative	78
4.1	List of variables included in the logistic regression model	95
4.2	Ratio scale for PCM	98
4.3	The indexes used in quantifying natural landscape configuration	107
4.4	The indexes used in quantifying landscape composition	108
4.5	The indexes used in quantifying core area landscape characteristics	110
4.6	Total of jurisdiction areas under different local authorities with sub-districts	113

4.7	Percentage of built up areas in all sub-districts under different local authorities	118
4.8	Comparison of urban land uses composition of Iskandar Malaysia between 2008 and 2025.	120
4.9	Type of data and the sources of data	133
5.1	The performance of accuracy assessment based classification on LULC between 1994 and 2013	146
5.2	Land covers composition in Iskandar Malaysia between 1994 and 2013	147
5.3	Historical transition of land use change from 1994 to 2013	158
5.4	Coefficient table for independent variables in relationship with the urban land use change from 2007 to 2013	164
5.5	Coefficient table for independent variables in relationship with natural landscape change from 2007 to 2013	168
5.6a	Transition probability of LULC change in 2020	171
5.6b	Projection transition of LULC change in 2020 (hectares)	171
5.7a	Transition probability of LULC change in 2025	172
5.7b	Projection transition of LULC change in 2025 (hectares)	172
5.8a	Transition probability of LULC change in 2030	173
5.8b	Projection transition of LULC change in 2030 (hectares)	173
5.9	Spatiotemporal future landscape change in Iskandar Malaysia region from 2015 to 2030 (hectare)	178

5.10	Transition of spatiotemporal landscape change between 2013 and 2030.	181
6.1	Results of spatial matrix calculation on urban pattern in Iskandar Malaysia region between 1994 and 2030	197
6.2	Development zones boundary and sub-districts	201
6.3	Natural landscape composition areas in the study area between 1994 and 2013	207
6.4	Natural ecological landscape zones in the study area	211
6.5	The distance (meters) of influence for the land covers towards natural landscape area (forest).	213

LIST OF FIGURES

FIGURE NO.	TITLE	PAGE
1.1	The study process	13
2.1	The resource continuum	24
2.2	Ecological services for human and surrounding environment benefits	27
2.3	Geometric figures indicate the shape of ecological patches with with monotonously increasing LSI (Landscape Shape Index) values	29
2.4	The concern of the ecological structure and function in landscape ecological study	34
2.5	Fragmentation of Natural Forest Cover in Peninsular Malaysia (1954-2000)	45
2.6	An illustration of how edge effect can reduce core ecological area in various size of natural landscape patch (adopted from Franklin, 1992 and Joan Voller, 1998)	49
2.7	An illustration of how natural landscape structure configuration influences the stability of core ecological area due to edge effect. (A) A circular natural patch tends to have a greater protection on core ecological area. (B) An irregularly shaped tends to have more edge and less core ecological area (adopted from Joan Voller, 1998)	49
2.8	The edge effects at regional scale and their influence distances (kilometres)	51
2.9	The edge effects at towards natural landscape patch and their influence distances (meters)	51

2.10	Edge effects on literature review divided into: (A) forest structure, (B) tree mortality, (C) forest microclimate and (D) biodiversity disturbance categories	53
3.1	The trend of focus of ecological impact on biotic and abiotic consequences	59
3.2	The concentration of edge effects studies in relation to land use and land cover change	59
3.3	Spatiotemporal landscape change studies in relation to ecological landscape structure and function assessment from 2000 to 2016	61
3.4	The use of landscape metrics in international peer-reviewed scientific papers: (a) evaluation of LULC pattern or changes; (b) habitat functions; (c) regulation functions; (d) information functions	81
4.1	The framework of the study	87
4.2	Temporal data analysis for landscape changes in Iskandar Malaysia	91
4.3	Relationship of MCA and GIS in decision making process	99
4.4	The contiguity filter used in this research	102
4.5	The process of analysis and components involved in spatiotemporal land use and land cover change analysis	102
4.6	Landscape ecological assessment in spatiotemporal landscape change in this study	105
4.7	Iskandar Malaysia region and context areas	112
4.8	Boundary of management areas for different local authorities in Iskandar Malaysia region	113
4.9	The population projection of Iskandar Malaysia by local authority	114
4.10	Population growth in Iskandar Malaysia region compared to Johor state	115

4.11	Composition of ethnics in 2006 for Iskandar Malaysia region	115
4.12	The policy dimension for the development of Johor Bahru from 1996 to 2011	117
4.13	The land use of Iskandar Malaysia in 2008	119
4.14	Urbanization promotion area in Iskandar Malaysia region	120
4.15	Land use of Iskandar Malaysia in 2025.	121
4.16	The current massive development projects that directly replace the ecological landscape areas in Iskandar Malaysia region.	122
4.17	Environment Sensitive Area in Iskandar Malaysia region	124
4.18	Blue and Green Corridor in Iskandar Malaysia region	125
4.19	Edge effects towards natural areas due to urban development and landscape mosaic change in Iskandar Malaysia	126
4.20	Natural and agriculture areas change to build up areas in Iskandar Malaysia	127
4.21	The components of spatial data involved in this research	129
4.22	Series of SPOT-5 satellite images of Iskandar Malaysia	135
4.23	Coordinate registration process for the satellite images	135
4.24	Mosaic process and extraction of images according to the boundary of study area	136
4.25	The classified images show the land cover of the study area	137
4.26	Geophysical spatial data involved in this study	139
4.27	The urban growth areas and natural landscape area loss in the past periods between 2007 and 2013	140

4.28	Distance of independent variables calculated in logistic regression analysis	141
4.29	Land use policy for development and protection zone	142
5.1	Comparison of classified images with land use data for accuracy assessment	146
5.2	Land cover map of Iskandar Malaysia region in 1994	148
5.3	Land cover map of Iskandar Malaysia region in 2000	150
5.4	Land cover map of Iskandar Malaysia region in 2007	152
5.5	Land cover map of Iskandar Malaysia region in 2013	155
5.6	Transitional of LULC change in different period (Hectares)	157
5.7	Transition map shows the conversion of the land cover between 1994 and 2000	159
5.8	Transition of land cover change from 2000 to 2007	160
5.9	Transition of land cover change from 2007 to 2013	162
5.10	Predicted probability of urban change from 1997 to 2013	166
5.11	Weightage of variables in urban land suitability	175
5.12	Decision Wizard Tool calibrates the overlay process by considering the AHP weightage for the selected variables	176
5.13	Suitability map for future landscape change in Iskandar Malaysia region	177
5.14	Projected land cover change in 2020	179
5.15	Transition of land cover change from 2013 to 2020	180
5.16	Projection of land cover change in 2025	183
5.17	Transition of land cover change from 2020 to 2025	184
5.18	Projected land cover change in 2030	186

5.19	Transition of land cover change from 2025 to 2030	187
5.20	Overall LULC maps and models involved in the study.	189
6.1	Percentage spatiotemporal LULC change for the whole study period	194
6.2	Spatiotemporal urban growth map between 1994 and 2030	196
6.3	Total class area and percentage of the urban landscape in Iskandar Malaysia region from 1994 to 2030	198
6.4	Graph showing number of patches (NP), largest patch index (LPI) and patch density (PD)	198
6.5	Graph showing interspersion and juxtaposition index (IJI), and Shannon's Diversity and Shannon's Evenness Indexes for the study area from 1994 to 2030.	199
6.6	Landscape matrices for spatiotemporal of urban area changes in the study area	201
6.7	Spatiotemporal of natural landscape area degradation (hectares) for the whole study period (1994 to 2030)	207
6.8	Configuration and composition of spatiotemporal natural landscape change in Iskandar Malaysia region between 1994 and 2013	208
6.9	The area (CA) change and percentage of landscape (PLAND) for natural forest landscape in past and future periods	207
6.10	The index shows fragmentation trend (NP), and largest patch index (LPI) of forest landscape in past and future periods	212
6.11	The graph shows the shape index (SHAPE_MN) forest landscape in past and future periods	214
6.12	Spatiotemporal landscape configuration change at mangrove areas in the study region between 1994 and 2030	216

6.13	The changes of the natural mangrove landscape areas in Sungai Pulai (WF3) between 1994 and 2030.	218
6.14	The changes of the natural mangrove landscape areas along the coastal at mukim Pulai (WF4) between 1994 and 2030.	219
6.15	The changes of the natural mangrove landscape areas in Sungai Skudai (WF5) between 1994 and 2030.	220
6.16	The changes of the natural mangrove landscape areas in Plentong (WF6) between 1994 and 2030.	221
6.17	The changes of the natural mangrove landscape areas in Pasir Gudang (WF6 and WF7) between 1994 and 2030.	222
6.18	Landscape structure characteristic changes for the inland forest areas in the landscape in the study area	225
6.19	The changes of the natural mangrove landscape areas in Gunung Pulai (IF1) between 1994 and 2030.	226
6.20	The distance of influence for edge effect in all ecological landscape zones in the study area.	229
6.21	The degradation of the natural ecological landscape areas due to edge effect from surrounding LULC change.	231
6.22	Erosion and soil contamination that degrading the natural mangrove areas in Tanjung Piai and Sungai Pulai.	232
6.23	Overall core area zones (TCA) compared to the total area of natural landscape (CA) at different types of natural landscape areas in the study area.	234
6.24	Number of patches change for core ecological area.	235
6.25	The consequences from edge effect towards some of the natural landscape areas in the study area.	238
6.26	Comparison of spatiotemporal intertidal forest intertidal forest landscape and core area zones in the study region between 1994 and 2030.	239

6.27	Comparison number of patches (NP) between intertidal forest landscape patches (CA) and core area patches (TCA) in the study region between 1994 and 2030.	241
6.28	Comparison of spatiotemporal intertidal forest intertidal forest landscape and core area zones in the study region between 1994 and 2030.	242
6.29	Comparison number of patches (NP) between inland forest landscape patches (CA) and core area patches (TCA) in the study region between 1994 and 2030.	243
7.1	Landscape indices pattern for spatiotemporal ecological landscape degradation in the study region between 1994 and 2030.	252
7.2	Strengthening multidisciplinary research with enhancement of landscape ecology assessment in land use development and adaptive management process	267

LIST OF APPENDICES

APPENDIX	TITLE	PAGE
A	Overall Landscape Index Result	295
B1	Pairwise Comparison Method (PCM) In Expert Choice Software	306
B2	The Result of Pairwise Comparison Method (PCM)	306
C	List of respondents for data analysis	307
D	GPS Data Collection	308
E	Survey Form – Expert Knowledge Criteria Comparison For LULC Change	309
F	Survey Form – Expert Knowledge Criteria For Ecological Landscape Response (Edge Effect)	323

LIST OF ACRONYMS

AHP	-	Analytical Hierarchical Process
ARSM	-	Agensi Remote Sensing Malaysia
CA	-	Cellular Automata
CAI	-	Core Area Index
CBD	-	Central Business District
CDP	-	Comprehensive Development Plan
CPLAND	-	Core Area Percentage of Landscape
DEI	-	Depth of Edge Influence
DCAD	-	Disjunct Core Area Density (DCAD)
ED	-	Edge Density
ESA	-	Environmental Sensitive Area
GIS	-	Geographic Information System
IF	-	Inland Forest
IJI	-	Interspersion and Juxtaposition Index
IRDA	-	Iskandar Region Development Authority
LPI	-	Largest Patch Index
MBJB	-	Majlis Bandaraya Johor Bahru
MC	-	Markov Chain
MCA	-	Multi-criteria Analysis
MCE	-	Multi-criteria Evaluation
MPPG	-	Malis Perbandaran Pasir Gudang
MPJBT	-	Majlis Perbandaran Johor Bahru Tengah
MPS	-	Mean Patch Size
NCA	-	Number of Core Area
NP	-	Number of Patches
LULC	-	Land Use and Land Cover
PCM	-	Pairwise Comparison Method
PD	-	Patch Density

PLAND	-	Percentage of Landscape
PRD	-	Patch Richness Density
PSS	-	Planning Support System
RSO	-	Rectified Skewed Orthomorphic
SHDI	-	Shannon's Diversity Index
SHEI	-	Shannon's Evenness Index
SHI	-	Shape Index
TCA	-	Total Core Area
LEA	-	Landscape Ecological Assessment
MAUT	-	MultiAttribute Theory
WF	-	Wetland Forest

CHAPTER 1

INTRODUCTION

1.0 Introduction

Landscape change is a spatial consequence that is the result of the complex dynamic land use change process and it is related to human activities, urban expansion and natural area loss. The dynamic land use and land cover (LULC) change for development is influenced by the rapid growth of the population as well as economic activities. The concentration of the population and economic activities in our urban areas creates demands for more land to be made available for development for new housing areas, commercial and industrial land, and public infrastructure. As a consequence, unexploited areas in and around urban areas are often likely to be converted into urban landscapes. The trade-off is a significant issue where many urban areas are expanding their boundaries and removing spaces that were previously open and natural landscape areas. In relation to the issue, the global urban population will increase in future with almost fifty percent of the world's population living in urban areas by 2025 (United Nation, 2009). Thus, urbanization will continue to modify the structure and function of natural landscape areas.

Our past experience shows that rapid landscape change has a significant negative impact on the environment. The environmental problems associated with the dynamic change of urbanization are biodiversity loss (Zhao *et al.*, 2005;

Matsushita *et al.*, 2006; Li *et al.*, 2010 and Nguyen, 2014), natural forest fragmentation (Abdullah and Nakagoshi, 2007; Li *et al.*, 2009), agriculture land loss (Pattanavibool and Derden, 2002; Gasparri *et al.*, 2009 and Su *et al.*, 2011), pollution (Nedeau *et al.*, 2003), microclimate degradation and ecosystem disturbance (Cumming *et al.*, 2012). From the perspective of ecological importance, land use and land cover change are deteriorating the ecological service and value at regional and local levels (Walters *et al.*, 2008). It is becoming increasingly difficult to ignore the issues of natural landscape degradation while urban development is in progress and degrading the serenity of our natural landscape.

The character of our natural landscape structure is related to the performance of ecological services (human's benefit) and values (ecosystem's benefit). The presence of the planet's natural areas provide ecosystem services and values, such as, carbon dioxide storage, stabilize microclimate, erosion protection, water catchment and, food resources and shelter for natural habitat (Tuan Vo *et al.*, 2012). However, natural area loss and fragmentation affect the structure of natural areas which consequently disrupt those ecological functions especially within the human dominated landscape area (Glennon and Kretser, 2013). Moreover, with rapid urbanization in the recent decades, a complex landscape mosaic between urban and natural land creates a more critical situation for the ecological landscape areas. The interaction between land use activities and natural landscape areas do not only represent a certain restructuring of the physical element of our natural landscape areas, it also affects the quality of core ecological areas in the remaining natural landscape patches. Natural areas adjacent to urban land uses are exposed to the edge effect and the depth of influence from complex external activities which subsequently reduce their core area through the deposition of species and the mortality of natural elements (Pattanavibool *et al.*, 2002 and Baker *et al.*, 2008).

Moreover, the fragmentation of natural areas reduces the size and connectivity of ecological zones and disturbs the meta-population process of species, prohibits energy transfer, instability of habitat, and increases competition in ecosystem (Hess and Fischer, 2001; Hersperger, 2006 and DeClerck, 2010). It is more critical because it impedes by the built up areas and could seriously deteriorate various ecological elements. As a result a certain depth of negative influence

restructures the core ecological zone (Cancino, 2005 and Lee *et al.*, 2008) which will harm the sustainability of the ecosystem. It can be considered as having a hidden impact towards a complex landscape mosaic that is not easy to interpret with the limited knowledge about the ecological response resulting from human land use activities. The quality of ecological function as it relates to the natural landscape has been seriously highlighted in the past and some theoretical frameworks were developed to resolve this challenge. Nevertheless the integration of these two components is not a simple matter without the combination of good knowledge and tools in terms of the decision-making process (Jogman, 2002; Corry *et al.*, 2005; Mortberg *et al.*, 2007; Reino *et al.*, 2009 and Llausas *et al.*, 2012). While in the process of considering and dealing with the issue, the different levels of knowledge on spatial land use change behaviours and specific ecological responses tend to increase the knowledge gap.

Many studies have been conducted around the world that highlight the issues of fragmentation of natural landscape and highlight the awareness of the changing landscape as it relates to the urbanization issue and the resulting ecological effect it has in terms of the spatiotemporal changes. However, a limited number of past studies integrate the spatiotemporal landscape change with the ecological function analysis. They revealed the spatiotemporal aspect of the landscape structure change but not been completely adequate to integrate the ecological response in regard to the LULC activities. This is an important information to justify the serenity of the patches of natural landscape and the capability of ecological areas to sustain their service and value. The different levels of knowledge on spatial land use and land cover change behaviours with specific ecological response increase the gap in addressing and dealing with the issue. Thus, a further understanding of the ecological consequences of the land use and land cover change area remains an important issue in term of the landscape change.

The issues highlighted above and past research experienced shows an apparent gap on the subject of land use planning with ecological consideration and its effect. Issues on the natural landscape structure change with the quality of the natural landscape patches should be carried out holistically in terms of landscape planning. Structure, function and change are three of the important components in landscape

ecology that must be integrated for any consideration in the urbanization process. Thus, this study is conducted to identify the approach and method to enhance the landscape ecological consideration by integrating spatiotemporal landscape change with an assessment of the ecological structure and function. While the ecological structure consequence refers to the natural landscape composition and configuration. Meanwhile the ecological function considers the natural landscape structure quality due to the interaction with its surrounding land use activities. The framework applied in this research is in response to the current environmental issues.

1.1 Research Background

Landscape can be defined as important resource components that consist of tangible (spatial elements) and intangible elements (human observer response) related to the characteristics of its spatial features (Sung *et al.*, 2001). Naveh (1995) and Jongman (2002) defined landscape resources as natural and cultural landscape components with regard to their interaction through economic condition, technical and social aspects, planning and policy. Based on this, it shows the changes of the global spatial dimension are highly related to the interaction and response between the natural landscape resource and spatial human development. The exploitation and degradation of ecological areas are part of the development process that restructures the physical form of the natural landscape areas. However, the ecological elements have been neglected as part of the ecosystem in development where most of the urbanization studies only consider the natural landscape structure dimension without having an understanding of their deteriorated functions (Rafiee *et al.*, 2009 and Stoate *et al.*, 2009).

Rapid urban development is growing and there is no sign of slowing down especially in the developing countries. It is becoming one of the most significant current discussions globally due to its impact towards landscape change and the resulting environmental degradation. On a global basis, almost 1.2 million kilometres square of forest and woodland, and 5.6 million kilometres square of grassland and pasture have been converted to other uses (Prato, 2005). This is followed by the

expansion of agricultural land that removed one part of the natural forest area in the world (Jongman, 2002) and reduced the diversity of natural landscapes; biotopes have either disappeared or have been isolated (Vuilleumire and Droux, 2002) with loss of heterogeneity (Gurrutxaga *et al.*, 2010). Wilderness areas have been encroached and river channels have been altered, and represent significant changes in terms of the important agriculture land and loss of forest areas that are rich with ecological services and values. The uncontrolled development and lack of consideration of the ecological importance of the development in the past have significantly caused a series of environmental problems, such as, the appearance of urban heat island, the altering of the hydrological characteristic, limited on carbon dioxide absorption, and the reduction of biodiversity which impacted on the ecological structure, function and dynamics (Han, 2009 and Yang *et al.*, 2011). In addition, it is expected that the urban population will be about 3.8 billion and predicted to reach 5.0 billion in 2030 (United Nation, 2009). The massive expansion of urban areas in the future will result in a tremendous change of the landscape structure and the function of the ecological areas (Jongman, 2007; Solon, 2009; Gurrutxaga *et al.*, 2010 and Vimal *et al.*, 2012).

The issues have been brought to the forefront in terms of the status of the current approaches with respect to understanding and awareness the ecological response in landscape change. The current decision- making process seems difficult to consider holistically in regard to the ecological response that result from the spatiotemporal landscape change interaction. The development of such model is one of the most important challenges as the urban areas are growing and restructuring the natural ecological areas. The limitation of the authority and platform for an ecologist to contribute in the decision planning process also creates a significant gap of understanding the ecological consequences related to landscape change. The stability and quality of ecological patches are neglected as it is challenging and difficult to measure in the development process and eventually the output plan negatively affects the natural landscape areas (Gattie *et al.*, 2007). The complex relationships between the dynamic land use change and ecological stability pose another question regarding how a different understanding of the various disciplines can be integrated along with the decision making process (Hazell *et al.*, 2000 and Musacchio *et al.*, 2005).

The interaction between land use planning and the ecological response becomes an obvious conflict since both disciplines present different levels of study. In land use planning exercise, while it is normally conducted at the macro-scale, an ecological study focuses on the specific habitat and site. Different perceptions on ecological areas and difficulty in translating the different techniques to a standard approach cause a conflict to achieve the integration (de Koning *et al.*, 2007 and Bishop, 2011). The landscape ecology however attempts to close the gap between ecological sustainability and urban landscape change, and serves to link the detailed characteristics of both components with the landscape changes. Nevertheless, since most of the past studies focus only on the effect of the structure of the natural landscape, it is difficult to describe the quality of the ecological function in response to change in the urban landscape. Specifically, the edge effect from the surrounding land use activities and the stability of the core ecological area in the remaining natural landscape areas have not been thoroughly considered in previous landscape change studies.

The shortcomings of the existing approach in terms of sustainable landscape planning attract the attention of and require decision makers to identify an achievable method. In order to maintain the ecological importance of the natural landscape, the modeling and enhancement of the approach ought to be considered in the landscape planning. This study is conducted to identify the approach and model to integrate the change of the landscape more in alignment to respond to the ecological aspect. It considers several spatial dimensions of the natural landscape structure change, such as, patch quality, patch context, boundaries, connectivity and landscape mosaic (land use activities in the surrounding area). Furthermore, the study evaluates the serenity and stability of the fragmented natural landscape areas by considering the composition and configuration of the landscape pattern. Landscape mosaic it also considers the relationship between the composition and configuration of the landscape pattern. This involves integrating the knowledge from the perspective of different disciplines and associating the knowledge into a standard platform. This is for the purpose of extending a more in-depth understanding of the landscape change scenario.

1.2 Problem Statement

In regard to the degradation of the natural landscape as temporal landscape change there is increasing concern that it has transformed the spatial patterns which has had an influence on the ecology and biological structure. Fragmentation, homogenization and the shrinking size of the natural areas are consequences from spatiotemporal LULC change particularly for urban development and agriculture expansion. Although many landscape change research studies have been conducted in the past it is difficult to understand of the ecological response due to the limited knowledge and approach (Sun *et al.*, 2012). In fact, past studies of landscape ecology have mostly focused on the impact of urban development and evaluate the static pattern of the consequences related to the natural landscape (landscape structure; composition and configuration). However, the ecological function degradation due the natural landscape structure change is been difficult to translate. The impact is apparent to the ecologist however it is difficult to explain to professionals of other disciplines, for example, designers or land use planners. Thus, in the process of urban development, attention to ecological sensitivity is not always given serious consideration.

The existing applications have limitations to incorporate ecological stability assessment in land use plan development such as the existing landscape planning and ecological assessment applications. The spatial cohesion of dynamic urban landscape change is difficult to translate the ecological responses (Ferreira *et al.*, 2013). Although many studies have successfully simulated the potential of future urban development and landscape changes, for example spatiotemporal urban landscape studies in Rome (Frondoni *et al.*, 2011), Wisconsin (Weng, 2007), North Carolina (Kirk *et al.*, 2012) and Shenzhen (Li and Yang, 2015), in the spatial changes of structure and function of the natural landscape present another topic for research. The interaction between the ecological responses of the LULC cover change is challenging to integrate in a similar platform. This is because the areas that remain fragmented natural landscape areas face immense pressure to sustain their quality whilst urban areas continue to expand their boundaries.

Then, poor consideration regarding the sensitivity of the ecological area in the landscape change leads to the degradation of the ecological value. Most of the current spatial planning applications only consider the natural landscape areas as a descriptive data without taking into account the biodiversity assets (Huang *et al.*, 2009; Li, *et al.*, 2010 and Miras *et al.*, 2014). The consideration of the urban expansion along with ecological sensitivity must be on a similar platform and requires the development of a better model. This main issue defines a requirement for the identification of the platform particularly engaging with the multidisciplinary expertise in the landscape change study. Any limitation in analyzing the ecological function in relation to its structure characteristics impairs the stability of the ecological area and consequently disturbs the quality thereof (Paudel and Yuan, 2012). Thus, consideration of the ecological function with structure characteristics is necessary to improve on the understanding of the ecological sensitivity and maintain their quality.

Ecological sensitivity and quality assessment in landscape change study requires an integrated model as a tool for holistic landscape change. Yet there is a gap between the methods and techniques in land use change analysis and ecological assessment where the parameter and measurement techniques are at different levels and it is difficult to support the holistic assessment of spatiotemporal landscape change and ecological response (Yue *et al.*, 2011 and He *et al.*, 2011). Moreover, the existing methods in ecological modeling are too specific and difficult to combine with other models especially at the landscape level (Table 3.2, Section 3.2.1). The probability of parameter and technique needs to be identified and should focus on how the ecological function assessment could fit in with a landscape change interaction study. This issue ought to be considered taking into account several technical aspects, such as, the spatial data integration, the accumulation of the input of experts, and the appropriate medium of analysis for output presentation.

Furthermore, landscape change is a dynamic process of human-driven land use change that requires continuity assessment in regard to the way it works. Temporal interpretation is essential for a better understanding of the

planning and management of future landscape change (Giam *et al.*, 2010). It is a powerful method that visualizes the characteristic of landscape change where historical experiences and future landscape scenarios are important to explain the quality status of natural landscape areas. Nevertheless, there is a dearth of knowledge to explain the rules and variables involved in landscape change and ecological response (Vreese *et al.*, 2016). Thus, the identification of the rules and variables is investigated in spatiotemporal landscape change and is outlined to predict the landscape change scenario.

The problem statements above form the direction for the research that attempts to improve spatiotemporal landscape change assessment. For this reason, the motivation for the study is to develop a method that incorporates urban development study with ecological structure and function assessment in order to extend an understanding of ecological stability and quality in dynamic landscape change. The effort of this research coupled with individual based analysis in a new platform comprises a different input knowledge. It is believed that the study will bridge the gap of LULC change with ecological interaction for the enhancement of the tool in consideration of the landscape change.

1.3 Research Questions

The problem statements highlighted above raise several research questions which reflect the approach and method involved in this research. The research questions are listed as follows:

- i. Which are the best measurements and indicators to evaluate the natural ecological landscape interaction and stability besides the natural landscape pattern analysis in spatiotemporal landscape change?
- ii. How to develop the approach and method in visualizing the temporal changes of ecological landscape to achieve a better interpretation and understanding?

- iii. What data are required and how might the database design be established to support the spatial dynamic modeling of land use and land cover change analysis and ecological sensitivity assessment?
- iv. What is the best platform for the landscape planning study that integrates the various multidisciplinary experts in regard to land use change interaction with ecological consequence?

1.4 Research Aim and Objectives

The aim of this study is to develop spatiotemporal landscape change model that integrates the assessment of the interaction between the spatiotemporal landscape change characteristics with ecological function.

Objectives:

- i. To design and establish a spatial dynamic model that incorporates both dynamic landscape change and ecological interaction
- ii. To identify the approach, technique and parameters that can be engaged for the integration of the landscape change model
- iii. To define the rules and variables involved in the spatial dynamic modelling of the landscape change and ecological function assessment
- iv. To formulate the database design and structure to support the parameters and techniques employed in the analysis of the spatiotemporal landscape change with deliberation of the ecological elements

1.5 Significance of Research

This research improves the understanding of spatiotemporal LULC landscape change with ecological area interaction. The visualization of the spatiotemporal landscape pattern and further assessment on the quality of ecological patches are important for adaptive management in landscape planning. The size, shape, connectivity, composition and configuration of the natural landscape scenario within the development area reveal the quality of the ecological aspect in the environment. The information then provides a better understanding of the ability of the fragmented natural areas as it relates to sustaining their services and values.

In addition, this research attempts to connect one of the important theoretical parts in landscape ecology. It instills ecological area function stability assessment through statistical analysis and expert knowledge in both ecology and land use planning. The method and technique used in this research are complementary to each other in order to strengthen the urban change analysis and ecological evaluation. This spatiotemporal landscape change with ecological function analysis is a time series assessment that improves the landscape planning method in adaptive management.

On the technical aspect, this research enriches the ability of spatial modeling by deploying the integration of GIS and Remote Sensing with related techniques that contribute to more intelligent application in landscape change analysis. Information technology used in this research provides a good basis for environmental modeling that can be implemented and upgraded for other environmental applications. Overall the study demonstrates and shows how important and significant of model development, information technology integration and ecological approach enhancement.

1.6 Scope of Research

Several aspects are considered in this research, such as, the level of the attributed detail in urban and ecological landscape change, the involvement of

expertise, and the scale of spatiotemporal dimension. This study focuses on the spatiotemporal landscape change as it highlights the LULC change, such as, built up areas, bare soil, agriculture land, water body, and natural forest areas that are divided into forest land and wetland area. Those spatial elements temporally represent the characteristics of landscape pattern in the study area.

The second aspect considered in this research is input from experts on the elements of urban landscape change and ecological sensitivity. The input contributes to the analysis for projecting future landscape change and ecological response. The input in the ecological aspect is transformed at the same level of the urban landscape change environment. With respect to the ecological aspect, this research collects ideas from the experts about the interaction of natural landscape with other land use activities. In this study, edge effect evaluation and core ecological zone determination are utilised and are based on the input from the experts.

On a spatial and time scale, this spatiotemporal landscape change study includes historical experience and future LULC. The selected durations for landscape change assessment are related to policy evolution in the study area. It refers to significant milestones of development policy in the past and in the projection of future landscape change that is based on the prospect of future development. The result of simulation indicates the spatial character of urban development behaviour and the ecological consequences of the landscape as they relate to the existing development policy.

The scope of this study fosters the ability of this research to be conducted in order to increase the understanding of ecological impact in landscape changes. The method used in this research could be a platform in order to extend knowledge for better landscape planning.

1.7 Research Outline

This study comprises several stages which involve various sub-activities in each stage. It includes the identification of the research framework, the model selection and development, the database design and development, the data analysis and recommendations. The outline of the overall process in this study is identified and illustrated in Figure 1.1.




Figure 1.1: The study process

Stage 1 - Preliminary Stage, Finalizing Framework

The first stage of the study focuses on the current situation of landscape change underpinning the theory of sustainable development and landscape ecological assessment. The trend of the urbanization process with ecological impact consideration is analysed to identify the limitation of current applications in landscape change studies. The existing approaches and methods used in ecological modelling are reviewed in relation to the landscape ecological structure and function change due to the interaction with the urban development. The identification of approaches to evaluate the relationship between ecological elements and other land use interaction is conducted to define the possibility of future integration in landscape change study. A comprehensive review based on referred journal, articles, theses, development plan documentation, government publication and books forms the idea of the research and supports the development of the goal and the objectives of the study.

Stage 2 - Model identification and integration

An analysis of the approaches and methods that are currently available, that have been applied in landscape change and landscape ecology studies in the past serves to highlight an approach that could be possibly be used in the integrated landscape change study. Specifically, the methodologies used in past studies are evaluated by taking into account their capabilities in managing multidisciplinary input and spatiotemporal landscape change. The advantages and disadvantages of previous methods and techniques are analysed that refer to the landscape ecological consideration in landscape change study. Although various methods have been used in land use planning and ecology assessment, it is a challenge to identify the possibility to integrate them in the dynamic process of urbanization and ecological quality assessment.

Nevertheless, several models are selected based on the requirement of the research, such as, Cellular Automata (CA), Logistic Regression, Markov Chain and Analytical Hierarchical Process (AHP) for the landscape change study. Meanwhile

landscape ecological assessment employs the landscape index as a tool to analyse the ecological structure and function in landscape change. There are several sub-models involved to analyse the ecological landscape structure change in terms of composition and configuration as well as ecological patches response due to the interaction with the surrounding land use. The quality of the ecological patches is evaluated utilising the Core Area Model that considers the edge effect of the surrounding land use activity. This model is useful to define the quality of remaining fragmented natural landscape areas within other land uses and land covers.

Those models are considered based on their capabilities in the integration with the spatial appearance of the landscape change analysis. Further, the data format is determined and designed to fit the models used in this study. GIS and Remote Sensing are the main tools utilized in this study and all of the data is transformed into their environments. The types and levels of the spatial and non-spatial data are identified in regard to the requirement of the models and research objectives.

Stage 3: Database design and collection

The next step of the study is designing the database for the spatial and non-spatial data. The design of the database is referred to as the requirement of the selected models used and the level of ecological consideration in this study. LULC maps are the main data where the input is derived from a series of satellite images to show the extent of landscape changes. Spatial topographic data is another important component that is used in the analysis and it would be in a standard spatial geo-reference and time period. The collection and development of the spatial data mostly from secondary sources, such as, the Department of Town and Country Planning, the local authorities and the Agencies of Remote Sensing Malaysia (ARSM). Some of the data should be updated and validated in order to increase the accuracy of the data and ensure it suits the requirement of the analysis models. While the Erdas Imagine software is used for processing the satellite images, ArcGIS 10.2 is utilized to analyse other geophysical data.

Since expert input is another component of the data that is important to support the models, therefore a questionnaire is prepared to acquire the input (Chapter 5, Appendix E and Appendix F). That input emphasizes the suitability and probability analysis of future urban growth areas where the projection of landscape changes is guided by their opinion. At the same time, the experts deliberate on the ecological impact of natural landscape areas in relation to the implication of the interaction between the urban landscape change and ecological sensitivity. This input is an important input because experts describe the depth of the influence of the ecological patches within the complex landscape mosaic. This input is normalized to ascertain a standard value of edge effect which is finally used to assess the quality of natural areas through the core ecological area. This is the loose-coupling method that is used within the GIS environment in this research.

Stage 4: Model Implementation and Data Analysis

At this stage, there are three major analytical processes involved in this study where the established model is implemented in regard to the case study area. The preliminary stage explores the trend of historical landscape change in the study area. The urban expansion pattern and natural landscape area degradation are analysed from 1994 to 2013. Furthermore, the degree of influence of spatial variables involved in the past landscape change is measured through the logistic regression model. The analysis highlights and ranks the criteria that have had a strong influence on the past changes. The historical experiences will serve as a basis for the next stage of the analysis to project the probability of future landscape change in the study area.

The following analysis extends the previous results to predict the future landscape change. Based on the logistic regression results, input from experts then consider strengthening the model of suitability analysis to define the suitability land for future urban land development. The Analytical Hierarchical Process (AHP) identifies the weightage for all significant variables in preparing the suitability map. The Pairwise Comparison Method is used in AHP and the Expert Choice software is used to run the model. At the similar stage, the probability of future changes is identified through the Markov Chain model. The historical experience in each period

is set up as a basis for the future projection of the landscape change. The simulation projects the probability of landscape changes in 2020, 2025 and 2030 by using the appropriate historical path. The interpolation of future spatial characteristics is translated using the Cellular Automata model.

In the final part of the analysis, the composition and configuration of the natural landscape area changes are measured using the selected landscape index. Several landscape matrices are utilised to evaluate the condition of natural landscape areas at class and landscape levels. The analysis focuses on the landscape ecological structure changes along the study period. The result shows the loss and fragmentation pattern of the natural landscape due to the spatiotemporal landscape changes in the study area. Moreover, the ecological stability of the remaining natural areas is evaluated based on the landscape structure, mosaic and edge effect from the adjacent land use activities. The edge effect from the surrounding land use activities towards natural areas is considered to delineate the ecological core zone in the natural areas. Input from experts on the depth of influence determines the quality function of ecological elements in the natural landscape areas. The FRAGSTATS software is used to run the analysis and the indexes indicate the condition of natural landscape structure and function within the dynamic landscape changes. The outcome of the analysis is discussed further in relation to LULC change pattern and ecological area consequences. The interpretation of the results emphasizes several aspects of land use planning, policy evolution and the most important aspects in landscape ecology.

Stage 5: Recommendation and Conclusion

Finally, as part of the contribution of the study, the recommendation for the application of this model is stated to enhance future landscape change studies with concern on the ecological elements. The recommendation also includes suggestions for improvement of the model for better application in future research. It could enhance better knowledge integration particularly in the process of analyzing a complex dynamic landscape change.

1.8 Thesis Organization

This thesis consists of seven chapters. The first chapter addresses the background of the study and highlights the issue of landscape change and ecological interaction. It describes the direction and focus of the research through the development of the goal and objectives of the study. The significance of the study is emphasized in this chapter and then the strategy to conduct the study follows.

The next chapter, Chapter 2 discusses the related literature about LULC change in urbanization studies as well as the landscape ecology approach and assessment in landscape change applications. Issues on past landscape change studies are highlighted and critiqued via the underpinning theories of landscape planning and landscape ecology. Then, the existing approaches, methods and techniques are discussed in Chapter 3 with regard to the spatiotemporal landscape change behaviors. Likewise, the discussion on the role of information technology in particular the geographic information system in spatial modeling is highlighted. It is the approach that is used in simulating landscape change in the study area. The established models and techniques with their spatial capabilities are deliberated upon towards the possibility of integration in the landscape change and ecological assessment. All the process involved is determined by formulating the framework of the study. At the end, this chapter formulates the concept and approach used in this study.

The Chapter 4 then discusses the methodology of the research. The development of the framework of the study is explained in the early part of the chapter. This is followed by an explanation of the detailed process involved in the study. Every single process is discussed starting from the data preparation to the analysis stage. The integration of the spatial and non-spatial data is displayed in the data preparation process and in the analysis stage. It is followed by the sequence of analysis processes in landscape change simulation and landscape ecology consideration.

Chapter 5 discusses the first part of the analysis in this study where it explains the implementation of the spatial dynamic model in landscape change at the study area. Every output within this stage is discussed in order to clarify the

spatiotemporal landscape change behaviors. The relationship of variables with the historical landscape change is highlighted as a basis to support input from experts in the preparation of suitability areas for future landscape change. Apart from that, further discussion on the result of the probability of future landscape changes shows the future character landscape change. It relates to the selected rules from the historical experience. The patterns of future landscape changes are visualized in this chapter particularly the landscape ecological area changes.

Meanwhile, the discussion in Chapter 6 focuses on the landscape ecological consequences due to the landscape changes in the past periods and future projection. The discussion emphasizes the landscape ecology structure and function changes. The composition and configuration of the natural landscape changes are justified and then related with the ecological function situation. It reveals the scenarios of the landscape ecological situation for the entire study period through spatial visualization. Graphs and statistical tables support the information of landscape ecological change behavior in the study.

A comprehensive discussion in Chapter 7 deliberates the output from Chapter 5 and Chapter 6 towards improving the landscape change analysis. It relates to achieving the goal of the research and the objectives wherein the landscape ecological function is now highly considered in dynamic landscape change. Furthermore, the results of the study are linked to the implication of policy changes in the study area. It reveals detailed aspects of the ecological consequences from rapid urban development not considered before. This final chapter then synthesizes the findings of the entire study by discussing the achievements of the study approach, the methodology applied and the techniques used in this study. The advantages and disadvantages of the spatial modeling in this research are deliberated upon excluding the limitations of the study. At the end, the chapter suggests possible future research that could be conducted by applying the spatial dynamic modeling method from this study along with improvements.

Decision makers are confronted with many issues in landscape change as demand for spatial development are increasing and reducing significant ecological areas. Even though there are theoretical frameworks that have been

established, it seems that due to the limitation of knowledge and technical support it hinders a better comprehensive decision that could be made. It is vitally important that information on the phenomena in regard to the urbanization and ecological impacts be understood at all levels of the decision-making process. And the prediction in regard to the probability concerning the way it happens must be considered by decision makers in landscape planning. It requires advance technology from information system and the appropriate spatial modelling to manage, organise and address the issues. Thus, this research explores the use of spatial modelling in the LULC change with better landscape ecological assessment. It is hoped that the significance of this study will be able to improve the decision-making process in landscape planning and consequently enable it to sustain ecological values and services in the environment.

REFERENCES

- Aburas, M. M., Ho, Y. M., Ramli, M. F. and Ash'aari, Z. H. (2017). Improving the capability of an integrated CA-Markov model to simulate spatio-temporal urban growth trends using an Analytical Hierarchy Process and Frequency Ratio. *International Journal of Applied Earth Observation and Geoinformation*, 59, 65–78.
- Albert, C., Galler, C., Hermes, J., Neuendorf, F., Haaren, C. V. and Lovett, A. (2016). Applying ecosystem services indicators in landscape planning and management: The ES-in-Planning framework. *Ecological Indicators*, 61, 100-113.
- Alexandrino, E. R., Buechley, E. R., Karr, J. R., de Barros Ferraz, K. M. P. M. de Barros Ferraz, S. F., do Couto, H. T. Z. and Sekercioglu, C. H. (2017). Bird based Index of Biotic Integrity: Assessing the ecological condition of Atlantic Forest patches in human-modified landscape. *Ecological Indicators*, 73, 662-675.
- Alig, R. J., Kline, J. D. and Lichtenstein, M. (2004). Urbanization on the US landscape : looking ahead in the 21st century. *Landscape and Urban Planning*, 69, 219–234.
- Antrop, M. (2001). The language of landscape ecologists and planners: A comparative content analysis of concepts used in landscape ecology. *Landscape and Urban Planning*, 55, 163–173.
- Abdullah, S. A. and Nakagoshi, N. (2006). Changes in landscape spatial pattern in the highly developing state of Selangor, Peninsular Malaysia. *Landscape and Urban Planning*, 77, 263–275.
- Abdullah, S.A. and Nakagoshi, N. (2007). Forest fragmentation and its correlations to human land use change in the state of Selangor, peninsular Malaysia. *Forest Ecology and Management*, 241, 39-48.

- Ahris, Y., Mansor, I., Susilawati, S. and Zulherman, M. S. (2000). Assessment of urban development and land use changes' impaction on the environment: A case study of urban development in Klang Valley region, Malaysia. *Proceeding SENVAR 2000*, Surabaya, Indonesia.
- An, L. (2012). Modeling human decisions in coupled human and natural systems: Review of agent-based models. *Ecological Modelling*, 229, 25– 36.
- Avon, C., Berges, L, Dumas, Y and Dupouey, J. L. (2010). Does the effect of forest roads extend a few meters or more into the adjacent forest? A study understory plant diversity in managed oak stands. *Forest Ecology and Management*, 259, 1546-1555.
- Al-Badaii, F., Othman, M. S. and Gasim, M. B. (2013). Water quality assessment of the Semenyih River, Selangor, Malaysia. *Journal of Chemistry*, 2013.
- Arsanjani, J. J., Helbichb, M., Kainz, W., Bolorani, A. D. (2013). Integration of logistic regression, Markov chain and cellular automata models to simulate urban expansion. *International Journal of Applied Earth Observation and Geoinformation*, 21, 265–275.
- Ayad, Y. M. (2005). Remote sensing and GIS in modeling visual landscape change: a case study of the northwestern arid coast of Egypt. *Landscape and Urban Planning*, 73, 307–325.
- Backhaus, N. (2011). Regional Environmental Governance: Interdisciplinary Perspectives, Theoretical Issues. Landscape, spatial totalities or special regions? *Social and Behavioral Sciences*, 14, 193–202.
- Baker, S. C., Barmuta, L. A., McQuillan, P. B. and Richardson, A. M.M. (2008). Estimating edge effects on ground-dwelling beetles at clearfeled non-riparian stand edges in Tasmanian wet eucalypt forest. *Forest Ecology and Management*, 239, 92-101.
- Barau, A. S. (2017). Tension in the periphery: An analysis of spatial, public and corporate views on landscape change in Iskandar Malaysia. *Landscape and Urban Planning*, 165, 256–266.
- Basnou, C., Iguzquiza, J. and Pino, J. (2015). Examining the role of landscape structure and dynamic in alien plant invansion from urban Mediterranean coastal habitats. *Landscape and Urban Planning*, 136, 156-164.
- Beardsley, K., Thorne, J. H., Roth, N. E., Gao, S. and McCoy, M. C. (2009). Assessing the influence of rapid urban growth and regional policies on biological resources. *Landscape and Urban Planning*, 93, 172–183.
- Bishop. I. D. (2011). Landscape planning is not a game: Should it be? *Landscape Urban and Planning*, 100, 390-392.

- Bjo, C. and Lo, K. (2002). Biotope patterns in urban areas: a conceptual model integrating biodiversity issues in spatial planning. *Landscape and Urban Planning*, 58, 223–240.
- Bolte, J. P., Hulse, D. W., Gregory, S. V. and Smith, C. (2007). Modeling biocomplexity – actors, landscapes and alternative futures. *Environmental Modelling & Software*, 22(5), 570–579.
- Bone, C. and Dragicevic, S. (2010). Incorporating spatio-temporal knowledge in an Intelligent Agent Model for natural resource management. *Landscape and Urban Planning*, 96, 123–133.
- Broadbent, E. N., Asner, G. P., Keller, M., Knapp, D. E., Oliveira, P. J. C. and Silva, J. N. (2009). Forest fragmentation and edge effects from deforestation and selective logging in the Brazilian Amazon. *Biological Conservation*, 141, 1745–1757.
- Bruun, M. and Smith, H. G. (2003). Landscape composition affects habitat use and foraging flight distances in breeding European starlings. *Biological Conservation*, 114, 179–187.
- Burger, J. and Gochfeld, M. (2016). Initiating events, functional remediation, and assessment of risk to ecological resources. *Ecological Indicators*, 71, 32–40.
- Cancino, J. (2005). Modelling the edge effect in even-aged Monterey pine (*Pinus radiata* D. Don) stands. *Forest Ecology and Management*, 210, 159–172.
- Caniani, D., Labella, A., Lioi, D. S., Mancini, I. M. and Masi, S. (2016). Habitat ecological integrity and environmental impact assessment of anthropic activities: A GIS-based fuzzy logic model for sites of high biodiversity conservation interest. *Ecological Indicators*, 67, 238–249.
- Castillo, E. M., Martin, A. G., Aladren, L. A. L., Luis, M. M. (2015). Evaluation of forest cover change using remote sensing techniques and landscape metrics in Moncayo Natural Park (Spain). *Applied Geography* 62, 247–255.
- Chaffin, B. C., Garmestani, A. S., Angeler, D. G., Herrmann, D. L., Stow, A. C., Nystron, M., Sendzimir, J., Hopton, M. E., Kolasa, J. and Allen, C. R. (2016). *Journal of Environmental Management*, 183, 399–407.
- Chazal, J. D. and Rounsevell, M. D. A. (2009). Land-use and climate change within assessment of biodiversity change: A review. *Global Environmental Change*, 19, 306–315.
- Chan, Clifton, Barrow, Chang, Wong and Ting (2003). Development issues and prospects in Cameron Highlands. Bahagian Geografi, Pusat Pengajian Ilmu Kemanusiaan, Universiti Sains Malaysia, Pulau Pinang.

- Chan, K. M. And Vu, T. T. (2017). A landscape ecological perspective of the impacts of urbanization on urban green spaces in the Klang Valley. *Applied Geography*, 85, 89-100.
- Chang, H., Li, F., Li, Z., Wang, R., Wang, R. and Wang, Y. (2011). Urban landscape pattern design from the viewpoint of networks: A case study of Changzhou city in Southeast China. *Ecological Complexity*, 8, 51-59.
- Chen, G., Powers, R. P., Carvalho, L. M. T. and Mora. (2015). Spatiotemporal patterns of tropical deforestation and forest degradation in response to the operation of the Tucuruí hydroelectric dam in the Amazon basin. *Applied Geography*, 63, 1-8.
- Chen, L. and Nuo, W. (2013). Dynamic simulation of land use changes in port city: a case study of Dalian, china Liang. *Procedia - Social and Behavioral Sciences* 96, 981 – 992.
- Chiao, L. Y., Hsieh, C. and Chiu, T. S. (2012). Exploring spatiotemporal ecological variations by the multiscale interpolation. *Ecological Modelling*, 246, 26-33.
- Clec'h, S. L., Oszwald, J., Decaens, T., Desjardins, T., Dufour, S., Grimaldi, M., Jegou, N. and Lavelle, P. (2016). Mapping multiple ecosystem services indicators: Toward an objective-oriented approach. *Ecological Indicators*, 69, 508-521.
- Clement, F., Ruiz, J., Rodriguez, M. A., Blais, D. and Campeau, S. (2017). Landscape diversity and forest edge density regulate stream water quality in agricultural catchments. *Ecological Indicators*, 72, 627-639.
- Collinge, S. K. (1996). Ecological consequences of habitat fragmentation: implications for landscape architecture and planning. *Landscape and Urban Planning*, 36(1), 59–77.
- Conway, T. M. and Lathrop, R. G. (2005). Alternative land use regulations and environmental impacts: assessing future land use in an urbanizing watershed. *Landscape and Urban Planning*, 71, 1–15.
- Corry, R. C. and Nassauer, J. I. (2005). Limitations of using landscape pattern indices to evaluate the ecological consequences of alternative plans and designs. *Landscape and Urban Planning*, 72, 265-280.
- Cotter, M., Hauser, I., Harich, F. K., He, P., Sauerborn, J., Treydte, A. C., Martin, K. and Cadisch, G. (2017). Biodiversity and ecosystem services—A case study for the assessment of multiple species and functional diversity levels in a cultural landscape. *Ecological Indicators*, 75, 111–117.
- Cumming, G. S., Southworth, J., Rondon, X. J. and Marsik, M. (2012). Spatial complexity in fragmenting Amazonian rainforests towards an ecological threshold? *Ecological Complexity*, 11, 67-74.

- Costanza, R. and Ruth, M. (1998). Using Dynamic Modeling to Scope Environmental Problems and Build Consensus. *Environmental Management*, 22, 183-195.
- Cumming, G. S., Southworth, J., Rondon, X. J. and Marsik, M (2012). Spatial complexity in fragmenting Amazonian rainforest: Do feedbacks from edge effects push forests towards an ecological threshold? *Ecological Complexity*, 11, 67-74.
- Dajoz, R. (1977). Introduction to Ecology. Great Britain: Fletcher & Son Ltd.
- Dalloz, M. F., Crouzeilles, R., Gomes, M. A., Papi, B. and Prevedello, J. A. (2017). Incorporating landscape ecology metrics into environmental impact assessment in the Brazilian Atlantic Forest. *Perspectives in Ecology and Conservation*, 15, 216–220.
- DeClerk, F. A. J., Chadzon, R., Holl, K. D., Milder, J. C., Finegan, B., Martinez-Salinas, A., Imbach, P., Canet, L. and Ramos, Z. (2010). Biodiversity conservation in human-modified landscape of Mesoamerica: Past, present and future. *Biological Conservation*, 143, 2301-2313.
- De, H. (2000). Demanding more of landscape research. *Landscape and Urban Planning*, 47, 105–109.
- De Hierro, L. G. L., Moleon, M., Lupianaz, D. G., Virgos, E. and Jimenez, R. (2013). Positive and negative unintended human-induced effects on Iberian mole abundance at the edge of its distribution area. *Mammalian Biology*, 78, 276–282.
- De Groot, R. S., Alkemade, R., Braat, L., Hein, L. and Willemsen, L. (2010). Challenges in integrating the concept of ecosystem services and values in landscape planning, management and decision making. *Ecological Complexity*, 7, 260-272.
- de Koning, G.H.J, Benitez, P.C., Munoz, F. and Olschewski, R. (2007). Modelling the impacts of payments for biodiversity conservation on regional land-use patterns. *Landscape and Urban Planning*, 83, 255-267.
- de Paula, M. D., Groeneveld, J. and Huth, A. (2016). The extent of edge effects in fragmented landscapes: Insights from satellite measurements of tree cover. *Ecological Indicators*, 69, 196-204.
- Department of Town and Country Planning (JPBD) (2010). National Physical Plan-2. Ministry of Housing and Local Government, Malaysia.
- Department of Town and Country Planning (JPBD – Jabatan Perancangan Bandar dan Desa). (1998). Laporan Pemeriksaan: Rancangan Struktur (Pengubahan) Daerah Johor Bahru (1998-2020).

- DiBari, J. N. (2007). Evaluation of five landscape-level metrics for measuring the effects of urbanization on landscape structure: the case of Tucson, Arizona, USA. *Landscape and Urban Planning*, 79, 308–313.
- Didham, R. K. and Ewers, R. M. (2012). Predicting the impacts of edge effects in fragmented habitats: Laurance and Yensen's core area model revisited. *Biological Conservation*, 155, 104-110.
- Domon, G. (2011). Landscape as resource: Consequences, challenges and opportunities for rural development. *Landscape and Urban Planning*, 100, 338–340.
- Dupuch, A. and Fortin, D. (2013). The extent of edge effects increases during post-having forest succession. *Biological Conservation*, 162, 9-16.
- Du, X. and Huang, Z. (2017). Ecological and environmental effects of land use change in rapid urbanization: The case of Hangzhou, China. *Ecological Indicators*, 8, 243-251.
- Fadzil Rashid (2004). Pengintegrasian Model Simulasi/Kualiti Air dan Sistem Maklumat Geografi Menggunakan Pendekatan Analisis Keputusan Multikriteria Dalam Kajian Kesan Gunatanah Setinggal Terhadap Kualiti Air. Universiti Teknologi Malaysia: *Master Thesis (Geoinformatic)*.
- Fang, S., Gertner, G. Z., Sun, Z. and Anderson, A. A. (2005). The impact of interactions in spatial simulation of the dynamics of urban sprawl. *Landscape and Urban Planning*, 73, 294-306.
- Feng, Li., Wang, R., Paulussen, J. and Liu, X. (2005). Comprehensive concept planning of urban greening based on ecological principles: a case study in Beijing, China. *Landscape and Urban Planning*, 72, 325-336.
- Feng, Y., Liu, Y., Tong, X., Liu, M. and Deng, S. (2011). Modeling dynamic urban growth using cellular automata and particle swarm optimization rules. *Landscape and Urban Planning*, 102(3), 188–196.
- Ferreira, P. A., Boscolo, D., Viana, B. F. (2013). What do we know about the effects of landscape changes on plant-pollinator interaction networks? *Ecological Indicators*, 31,35-40.
- Frank, S., Furst, C., Koschke, L. and Makeschin, F. (2012). A contribution towards a transfer of the ecosystem service concept to landscape planning using landscape metrics. *Ecological Indicators*, 21, 30-38.
- Frondoni, R., Mollo, B. and Capotorti, G. (2011). A landscape analysis of land cover change in the Municipality of Rome (Italy): Spatiotemporal characteristics and ecological implications of land cover transitions from 1954 to 2001. *Landscape and Urban Planning*, 100(1-2), 117–128.

- Fry, G. L. A. (2001). Multifunctional landscapes-towards transdisciplinary research. *Landscape and Urban Planning*, 57, 159–168.
- Fu, P. and Weng, Q. (2016). A time series analysis of urbanization induced land use and land cover change and its impact on land surface temperature with Landsat imagery. *Remote Sensing of Environment*, 175, 205-214.
- Gasparri, N. G. and Grau, H. R. (2009). Deforestation and fragmentation of Chaco dry in NW Argentina (1972-2007). *Forest Ecology and Management*, 258, 913-921.
- Gattie, D. K., Kellam, N. N. and Turk, H. J. (2007). Informing ecological engineering through ecological network analysis, ecological modeling, and concepts of systems and engineering ecology. *Ecological Modelling*, 208, 25-40.
- Geneletti, D. (2008). Incorporating biodiversity assets in spatial planning: Methodological proposal and development of a planning support system. *Landscape and Urban Planning*, 84, 252–265.
- Geri, F., Amici, V., Rocchini, D., 2010. Human activity impact on the heterogeneity of a Mediterranean landscape. *Applied Geography*, 30, 370–379.
- Gilbert, A. J. and Jansen, R. (1998). Use of environmental functions to communicate the values of mangrove ecosystem under different management regimes. *Ecological Economics*, 25, 323-346.
- Giam, X., Bradshaw, C. J. A., Tan, H. W. and Sodhi, N. S. (2010). Future habitat loss and the conservation of plant biodiversity. *Biological Conservation*, 143, 1594-1602.
- Grabaum, R. and Meyer, B. C. (1998). Multicriteria optimization of landscapes using GIS-based functional assessments. *Landscape and Urban Planning*, 43, 21–34.
- Glennon, M. J. and Kretser, H. E. (2013). Size of the ecological effect zone associated with exurban development in the Adirondack Park, NY. *Landscape and Urban Planning*, 112, 10-17.
- Gong, C., Yu, S., Joesting, H. and Chen, J. (2013). Determining socioeconomic drivers of urban forest fragmentation with historical remote sensing images. *Landscape and Urban Planning*, 117, 57-65.
- Green, D. G. and Sadedin, S. (2005). Interactions matter—complexity in landscapes and ecosystems. *Ecological Complexity*, 2(2), 117–130.
- Grimm, V., Wyszomirski, T., Aikman, D. and Uchman, J. (1999). Individual-based modelling and ecological theory: synthesis of a workshop. *Ecological Modelling*, 115, 275–282.

- Gurrutxaga, M., Lozano, P. J. and Barrio, G. D. (2010). GIS-based approach for incorporating the connectivity of ecological networks into regional planning. *Journal for Nature Conservation*, 18, 318-326.
- Hadac, E. (1977). Complex interdisciplinary investigation of landscape. *Landscape and Urban Planning*, 4, 333-348.
- Hardt, E., Pereira-Silva, E., Dos Santos, R. F., Tamashiro, J. Y., Ragazzi, S. and Lins, D. B. (2013). The influence of natural and anthropogenic landscape on edge effects. *Landscape and Urban Planning*, 120, 59-69.
- Han, J., Hayashi, Y., Cao, X. and Imura, H. (2009). Application of an integrated system dynamics and cellular automata model for urban growth assessment: A case study of Shanghai, China. *Landscape and Urban Planning*, 91, 133-141.
- Haaren, C. v., Kretzschmar, B. W., Milos, C. and Werthmann, C. (2014). Opportunities for design approaches in landscape planning. *Landscape and Urban Planning*, 130, 159-170.
- Haas, J. and Ban, Y. (2015). Urban growth and environmental impacts in Jing-Jin-Ji, the Yangtze, River Delta and the Pearl River Delta. *International Journal of Applied Earth Observation and Geoinformation*, 30, 42-55.
- Haas, J., Furberg, D. and Ban, Y. (2015). Satellite monitoring of urbanization and environmental impacts-A comparison of Stockholm and Shanghai. *International Journal of Applied Earth Observation and Geoinformation*, 38, 138-149.
- Hazell, P. and Wood, S. (2000). From science to technology adoption: the role of policy research in improving natural resource management. *Agriculture, Ecosystems and Environment*, 82, 385-393.
- He, C., Tian, J., Shi. and Hu, D. (2011). Simulation of the spatial stress due to urban expansion on the wetlands in Beijing, China using a GIS-based assessment model. *Landscape and Urban Planning*, 101, 269-277.
- Hendrix, W. G., Fabos, J. G., and Price, J. E (1988). An ecological approach to landscape planning using Geographic Information System Technology. *Landscape and Urban Planning*, 15, 211-225.
- Herold, M., Scepan, J., Clarke, K.C., 2002. The use of remote sensing and landscape metrics to describe structures and changes in urban land uses. *Environmental Planning*, 34, 1443-1458.
- Hersperger, A. M. (2006). Spatial adjacencies and interaction: Neighbourhood mosaics for landscape ecological planning. *Landscape and Urban Planning*, 77, 227-239.
- Hess, G. R. and Fischer, R. A. (2001). Communicating clearly about conservation corridors. *Landscape and Urban Planning*, 55, 195-208.

- Hierro, L. G., Moleon, M., Lupianez, D. G., Virgos, E. and Jimenez, R. (2013). Positive and negative unintended human-induced effects on Iberian mole abundance at the edge of its distribution area. *Mammalian Biology*, 78, 276-282.
- Hobbs, R. (1997). Future landscapes and the future of landscape ecology. *Landscape and Urban Planning*, 37, 1–9.
- Hong, S. K., Nakagoshi, H. and Morimoto, Y. (2007). Landscape Ecological Applications In Man-Influenced Areas – Linking Man And Nature Systems. In: Hong, S. K., Nakagoshi, H., Morimoto, Y. Landscape Ecological Applications in Man-Influenced Areas. The Netherlands. 1-5; 2007.
- Hong, W., Jiang, R., Yang, C., Zhang, F., Su, M. and Liao, Q. (2016). Establishing an ecological vulnerability assessment indicator system for spatial recognition and management of ecologically vulnerable areas in highly urbanized regions: A case study of Shenzhen, China. *Ecological Indicators* 69, 540–547.
- Hossein, S. M. and Helbich, M. (2013). Spatiotemporal urbanization processes in the megacity of Mumbai, India: A Markov Chains-cellular automata urban growth model. *Applied Geography*, 40, 140-149.
- Howit, R. *Rethinking Resource Management: Justice, Sustainability and Indigenous Peoples*. London: Routledge. 2001
- Huang, S. L., Yeh, C. T. and Chang, L. F. (2010). The transition to an urbanizing world and the demand for natural resources. *Current Opinion in Environmental Sustainability*, 2, 136-143.
- Huber, P. R., Shilling, R. Thorne, J. H. and Greco, S. E. (2012). Municipal and regional habitat connectivity planning. *Landscape and Urban Planning*, 105, 15–26.
- Iskandar Region Development Authority (IRDA). (2008). Comprehensive Development Plan (CDP) for Iskandar Malaysia Region 2006-2025.
- Iskandar Region Development Authority (IRDA). (2011). Integrated Land Use Blueprint 2025.
- Jiang, P., Cheng, L., Li, M., Zhao, R. and Huang, Q. (2014). Analysis of landscape fragmentation processes and driving forces in wetlands in arid areas: A case study of the middle reaches of the Heihe River, China. *Ecological Indicators*, 46, 240-252.
- Jorgensen, S. E. (2009). Structurally dynamic models. In : Jorgensen, S. E., Chon, T. S., Recknagel, F. Handbook of Ecological Modelling And Informatics. Southampton, Boston: WIT Press. 221-239; 2009.
- Jorgensen, S. E. and Chon, T. S. (2009). Overview of the model types available for ecological modelling. In : Jorgensen, S. E., Chon, T. S., Recknagel, F. Handbook

- of Ecological Modelling And Informatics. Southampton, Boston: WIT Press; 9-40; 2009.
- Jongman, R. H. G. (2002). Homogenisation and fragmentation of the European landscape: ecological consequences and solutions. *Landscape and Urban Planning*, 58, 211–221.
- Jongman, R. H. G. (2007). Ecological Networks, From Concept To Implementation. In: Hong, S. K., Nakagoshi, H., Morimoto, Y. ed. *Landscape Ecological Applications in Man-Influenced Areas*. The Netherlands: Springer. 57-69; 2007.
- Knowlton, J. L. and Graham, C. H. (2010). Using behavioral landscape ecology to predict species' responses to land use and climate change. *Biological Conservation*, 143(6), 1342–1354.
- Karlson, M. and Mortberg, U. (2015). A spatial ecological assessment of fragmentation and disturbance effects of the Swedish road network. *Landscape and Urban Planning*, 134, 53–65.
- Kirk, R. W., Bolstad, P. V. and Manson, S. M. (2012). Spatio-temporal trend analysis of long-term development patterns (1900-2030) in a Southern County. *Landscape and Urban Planning*, 104, 47–58.
- Koffi, K. J., Deblauwe, V., Sibomana, S., Neuba, D. F. R., Champluver, D., De Canniere, C., Barbier, H., Traore, D., Habonimana, B., Robbrecht, E., Lejoly, J. and Bogaert, J. (2007). Spatial Pattern Analysis As A Focus Of Landscape Ecology To Support Evaluation Of Human Impact On Landscapes And Diversity. In : Hong, S. K., Nakagoshi, H., Morimoto, Y.ed. *Landscape Ecological Applications in Man-Influenced Areas*. The Netherlands: Springer. 1-5; 2007.
- Koschke, L., Furst, C., Frank, S. and Mekeschin, F. (2012). A multi-criteria approach for an integrated land-cover-based assessment of ecosystem services provision to support landscape planning. *Ecological Indicators*, 21, 54-66.
- Kuttner, M., Herman, A. and Wrбка, T. (2013). Do landscape structural patterns reflect Ecosystem Service provision? – A comparison between protected and unprotected areas throughout the Neusiedler See region. *5th Symposium for Research in Protected Areas*, 437-442.
- Lasanta, T. and Gonz, C. (2006). Using landscape ecology to evaluate an alternative management scenario in abandoned Mediterranean mountain areas. *Landscape and Urban Planning*, 78, 101–114.
- Lathrop, R. G., Tulloch, D. L., and Hatfield, C. (2007). Consequences of land use change in the New York – New Jersey Highlands , USA: Landscape indicators of forest and watershed integrity. *Landscape and Urban Planning*, 79, 150–159.
- Laurance, W. F. (1991). Predicting the Impacts of Edge Effects in Fragmented Habitats. *Biological Conservation*, 55, 77-92.

- Le, Q. B., Park, S. J. and Vlek, P. L. G. (2010). Land Use Dynamic Simulator (LUDAS): A multi-agent system model for simulating spatio-temporal dynamics of coupled human–landscape system 2. Scenario-based application for impact assessment of land-use policies. *Ecological Informatics*, 5, 203–221.
- Lee, S. W., Ellis, C. D., Kweon, B. S. and Hong, S. K. (2008). Relationship between landscape structure and neighborhood satisfaction in urbanized areas. *Landscape and Urban Planning*, 85, 60–70.
- Leitao, A. B. and Ahern, J. (2002). Applying landscape ecological concepts and metrics in sustainable landscape planning. *Landscape and Urban Planning*, 59, 65–93.
- Levin, N., Lahav, H., Ramon, U., Heller, A., Nizry, G., Tsoar, A. and Sagi, Y. (2007). Landscape continuity analysis: A new approach to conservation planning in Israel. *Systematics*, 79, 53–64.
- Li, F., Wang, R., Hu, D., Ye, Y., Yang, W. and Liu, H. (2014). Measurement methods and applications for beneficial and detrimental effects of ecological services. *Ecological Indicators*, 47, 102–111.
- Li, L., Sato, Y. and Zhu, H. (2003). Simulating spatial urban expansion based on a physical process. *Landscape and Urban Planning*, 64, 67–76.
- Li, J., Jiang, H., Bai, Y., Alatalo, J. M., Li, X., Jiang, H., Liu, G. and Xu, J. (2016). Indicators for spatial-temporal comparisons of ecosystem service status between regions: A case study of the Taihu River Basin, China. *Ecological Indicators*, 60, 1008–1016.
- Li, M., Huang, C., Zu, Z., Shi, H., Lu, H. and Peng, S. (2009). Assessing rates of forests change and fragmentation in Alabama, USA, using vegetation change tracker model. *Forest Ecology and Management*, 257, 1480–1488.
- Li, Y., Zhu, X., Sun, X. and Wang, F. (2010). Landscape effects of environmental impact on bay-area wetlands under rapid urban expansion and development policy: A case study of Lianyungang, China. *Landscape and Urban Planning*, 94, 218–227.
- Li, S. and Yang, B. (2015). Introducing a new method for assessing spatially explicit processes of landscape fragmentation. *Ecological Indicators*, 56, 116–124.
- Lien, J. K. (2003). *Integrated Environmental Planning*. United Kingdom: Blackwell Science Ltd.
- Lin, Y., Hong, N., Wu, P., Wu, C. and Verburg, P. H. (2007). Impacts of land use change scenarios on hydrology and land use patterns in the Wu-Tu watershed in Northern Taiwan. *Landscape and Urban Planning*, 80, 111–126.

- Linehan, J. R. and Gross, M. (1998). Back to the future, back to basics: the social ecology of landscapes and the future of landscape planning. *Landscape and Urban Planning*, 42, 207-233.
- Ling, O. H. L., Ting, K. H., Shaharudin, A., Kadarudin A. and Yaakob, M. J. (2010). Urban growth and air water quality in Kuala Lumpur City, Malaysia. *Environment Asia* 3(2), 123-128.
- Liu, D., Cao, C., Dubovyk, O., Tian, R., Chen, W., Zhuang, Q., Zhao, Y. and Menz, G. (2017). Using fuzzy analytic hierarchy process for spatio-temporal analysis of eco-environmental vulnerability change during 1990–2010 in Sanjiangyuan region, China. *Ecological Indicators*, 73, 612–625.
- Liu, H., Zhi, W., Li, X., Jie, L., and Bin, Z. (2009). Characterizing landscape dynamics of a small catchment under ecological rehabilitation interventions in Northwestern China. *Landscape and Urban Planning*, 93, 201–209.
- Llausàs, A. and Nogué, J. (2012). Indicators of landscape fragmentation: The case for combining ecological indices and the perceptive approach. *Ecological Indicators*, 15, 85–91.
- Locke, C. M. and Rissman, A. R. (2012). Unexpected co-benefits: Forest connectivity and property tax incentives. *Landscape and Urban Planning*, 104, 418-425.
- Malanson, G. P. (2003). Dispersal across continuous and binary representations of landscapes. *Ecological Modelling*, 169(1), 17–24.
- MacLean, M. G. (2017). Edge influence detection using aerial LiDAR in Northeastern US deciduous forests. *Ecological Indicators*, 72, 310-314.
- Masum, K. M., Mansor, A., Mohd Sah, S. A. and Lim, H. S. (2017). Effect of differential forest management on land-use change (LUC) in a tropical hill forest of Malaysia. *Journal of Environmental Management*, 200, 468-474.
- Marker, M., Angeli, L., Bottai, L., Costantini, R., Ferrari, R., Innocenti, L. and Siciliano, G. (2008). Assessment of land degradation susceptibility by scenario analysis: A case study in Southern Tuscany, Italy. *Geomorphology*, 93, 120–129.
- Mather, A. S. and Chapman, K. (1995). *Environmental Resources*. Logman Sciences and Technical, Essex, England.
- Matsushita, B., Xu, M. and Fukushima, T. (2006). Characterizing the changes in landscape structure in Lake Kasumigaura Basin, Japan using a high quality GIS dataset. *Landscape and Urban Planning*, 78, 241-250.
- McDonal, R. I., Forman, T. T., Kareiva, P., Busing, R. T. and Neugarten, R., Salzer, D. and Fisher, J. (2009). Urban effects, distance and protected areas in an urbanizing world. *Landscape and Urban Planning* 93, 63–75.

- McGarigal, K., Cushman, S.A., Neel, M.C., Ene, E. (2002). FRAGSTATS: spatial pattern analysis program for categorical maps. www.umass.edu/landeco/research/fragstats/fragstats.html. Access date: 21st August 2012.
- Malczewski, J. (1999). *GIS And Multicriteria Decision Analysis*. Department of Geography, University of Western Ontario, Canada.
- Mcrae, B. H., Schumaker, N. H., Mckane, R. B., Busing, R. T., Solomon, A. M., Burdick, C. A. and Service, F., et al. (2008). A multi-model framework for simulating wildlife population response to land-use and climate change, *Landscape Ecology*, 9, 77–91.
- Meyer, B. C., Wolf, T. and Grabaum, R. (2012). A multifunctional assessment method for compromise optimization of linear landscape elements. *Ecological Indicators*, 22, 53-63.
- Mernerd, A. and Marceau, D. J. (2006). Simulating the impact of forest management scenarios in an agriculture and landscape of southern Quebec, Canada, using a geographic cellular automata. *Landscape and Urban Planning*, 79, 253-265.
- Millward, A. A. and Mersey, J. E. (2006). Conservation strategies for effective land management of protected areas using an erosion prediction information systems (EPIS). *Environmental Management*, 61, 329-343.
- Montis, A. D., Caschili, S., Mulas, M., Ganciu, A., Bardi, A., Ledda, A., Dessena, L., Dessena, L., Laudari, L. and Fichera, C. R. (2016). Urban-rural ecological networks for landscape planning. *Land Use Policy*, 50, 312-327.
- Mortberg, U. M., Balfors, B. and Knol, W. C. (2007). Landscape ecological assessment: A tool for integrating biodiversity issues in strategic environmental assessment and planning. *Journal of Environmental Management*, 82, 457–470.
- Mirasa, P. Z., Palomo, I., Baggethun, E. G., López, B. M., Lomas, P. L. and Montes, C. Effects of land-use change on wetland ecosystem services: A case study in the Donana marshes (SW Spain). *Landscape and Urban Planning*, 122, 160-174.
- Murcia, C. (1995). Edge effects in fragmented forests—implications for conservation. *Trends Ecological. Evolution*, 10, 58–62.
- Musacchio, L., Ozdenerol, E., Bryant, M., and Evans, T. (2005). Changing landscapes, changing disciplines: seeking to understand interdisciplinarity in landscape ecological change research. *Landscape and Urban Planning*, 73, 326–338.
- Mörtberg, U. M., Balfors, B., and Knol, W. C. (2007). Landscape ecological assessment: a tool for integrating biodiversity issues in strategic environmental assessment and planning. *Journal of environmental management*, 82(4), 457–70.

- Moreno, M. L., Bernaschini, M. L., Harguindeguy, N. P. and Valladers, G. (2014). Area and edge effects on leaf-litter decomposition in a fragmented subtropical dry forest. *Acta Oecologica*, 60, 26-29.
- Moser, D., Zechmeister, H.G., Plutzer, C., Sauberer, N., Wrбка, T., Grabherr, G., 2002. Landscape patch shape complexity as an effective measure for plant species richness in rural landscapes. *Landscape Ecology*, 17, 657–669.
- Morisette, J. T., Richardson, A. D., Knapp, A. K., Fischer, J., Graham, A. G., Abatzoglou, J., et al. (2009). Tracking the rhythm of the seasons in the face of global change: Phenological research in the 21st century. *Frontiers in Ecology and the Environment*, 7(5), 253–260.
- Naveh, Z. (1995). Interactions of landscapes and cultures. *Landscape and Urban Planning*, 32(94), 43–54.
- Ndibisi, F., DeMeo, T. and Ditto, N. D. (1995). Environmentally sensitive areas: a template for developing greenway corridors. *Landscape and Urban Planning*, 33, 159-177.
- Nedeau, E. J., Merritt, R. W. and Kaufman, M. G. (2003). The effect of an industrial effluent on an urban stream benthic community: water quality vs habitat quality. *Environmental Pollution*, 123, 1–13.
- Nguyen, H. (2014). The relation of coastal mangrove changes and adjacent land use: A review in Southeast Asia and Kien Giang, Vietnam. *Ocean and Coastal Management*, 90, 1-10.
- Normann, C., Tschardtke, T. and Scherber, C. (2016). Interacting effects of forest stratum, edge and tree diversity on beetles. *Forest Ecology and Management*, 361, 421–431.
- Palacio, C. V., Berrouet, L., Lopez, C., Ruiz, A. and Upegui, A. (2016). *Ecosystem Services*, 22, 297-308.
- Pattanavibool, A. and Derden, P. (2002). Fragmentation and wildlife in montane evergreen forests, northern Thailand. *Biological Conservation*, 107, 155-164.
- Paudel, S and Yuan, F. (2012). Assessing landscape changes and dynamics using patch analysis and GIS modeling. *International Journal of Applied Earth Observation and Geoinformation*, 16, 66-75.
- Pavlacky Jr, D. C., Possingham, H. P. and Goldizen, A. W. (2015). Integrating life history traits and forest structure to evaluate the vulnerability of rainforest birds along gradients of deforestation and fragmentation in eastern Australia. *Biological Conservation*, 188, 89–99.

- Peng, J., Tian, L., Liu, Y., Zhao, M., Hu, Y. and Wu, J. (2017). Ecosystem services reponse to urbanization in metropolitan areas: *Thresholds identification*. *Science of the Total Environment*, 607-608, 706-714.
- Pimentel, D. and Pimentel, M. (2006). Global environmental resources versus world population growth. *Ecological Economics*, 59, 195–198.
- Piessens, K., Honnay, O., Devlaeminck, R. and Hermy, M. (2006). Biotic and abiotic edge effects in highly fragmented heathlands adjacent to cropland and forest. *Agriculture, Ecosystems and Environment*, 114, 335-342.
- Poelmans, L. and Rompaey, A. V. (2009). Detecting and modelling spatial patterns of urban sprawl in highly fragmented areas: A case study in the Flanders – Brussels region. *Landscape and Urban Planning*, 93, 10–19.
- Prato, T. (2005). Modeling ecological impacts of landscape change. *Environmental Modelling & Software*, 20(10), 1359–1363.
- Primdahl, J. (1990). Heterogeneity in agriculture and landscape: From segregation to integration. *Landscape and Urban Planning*, 18(3-4), 221–228.
- Rabe, N. S., Osman, M. M. and Bachok, S. (2014). Economics of Local People: Iskandar, Malaysia. *Procedia - Social and Behavioral Sciences*, 153, 463 – 478.
- Rafiee, R., Mahini, A. S. and Khorasani, N. (2009). Assessment of changes in urban green spaces of Mashad city using satellite data. *International Journal of Applied Earth Observation and Geoinformation*, 11, 431-438.
- Rafikul Islam and Saifulizan Tahir (2002). Geographic Information Systems and Decision Support System. Module 1: Fundamental of Multi-Criteria Decision Making (MCDM) Problems. *Short Course Paperwork*, UIAM.
- Reddy, C. S., Sreelekshmi, Jha, C. S. and Dadhwal, V. K. (2013). National assessment of forest fragmentation in India: Landscape indices as measures of the effects of fragmentation and forest cover change. *Ecological Engineering*, 60, 453-464.
- Rees (1989). Types of environmental resources. Mather, A. S. and Chapman, K. (1995). *Environmental Resources*. Logman Sciences & Technical, Essex, England. 14–16.
- Reino, L., Beja, P., Osborne, P. E., Morgado, R., Fabiao, A. and Rotenberry, J. T. (2009). Distance to edges, edge contrast and landscape fragmentation: Interactions affecting farmland birds around forest plantations. *Biological Conservation*, 142, 824-838.
- Rempel, R. S., Naylor, B. J., Elkie, P. C., Baker, J., Churcher, J. and Gluck, M. J. (2016). An indicator system to assess ecological integrity of managed forests. *Ecological Indicators*, 60, 860-869.

- Rescia, A.J., Willaarts, B.A., Schmitz, M.F., Aguilera, P.A., 2010. Changes in land uses and management in two Nature Reserves in Spain: evaluating the social–ecological resilience of cultural landscapes. *Landscape & Urban Planning*, 98, 26–35.
- Riutta, T., Slade, E. M., Bebbler, D. P., Taylor, M. E., Malhi, Y., Riordan, P., Macdonald, D. W. and Morecroft, M. D. (2012). Experimental evidence for the interacting effects of forest edge, moisture and soil macrofauna on leaf litter decomposition. *Soil Biology and Biochemistry*, 49, 124-131.
- Rocchini, D., Perry, G. L. W., Salerno, M., Maccherini, S., & Chiarucci, A. (2006). Landscape change and the dynamics of open formations in a natural reserve. *Landscape and Urban Planning*, 77, 167–177.
- Rodríguez, M. P., Kuemmerle, T., Segura, D. A., Milla, R. Z. and Cabello, J. (2012). Future land use effects on the connectivity of protected area networks in southeastern Spain. *Journal for Nature Conservation*, 20, 326– 336.
- Saaty, T.L. (1980). *The Analytic Heirarchy Process*. New York: McGraw-Hill.
- Saaty, T.L. dan Kerns, K.P. (1985). *Analytical Planning: The Organisations of Systems*. Oxford. New York. Toronto. Sydney. Frankfurt, Pergamon Press.
- Schlaefter, R., Iorgulescu, I. and Glenz, C. (2001). Management of forested landscape in mountain areas: an ecosystem-based approach. *Forest Policy and Economics*, 4, 89-99.
- Shaharudin, I. and Abdul Samad, H. (2004). Urban land use change and the Langat Basin ecosystem health. *Journal of the Malaysian Institute of Planners*, II, 51-68.
- Shi, Y., Sun, X., Zhu, X., Li, Y. and Mei, L. (2008). Characterizing growth types and analyzing growth density distribution in response to urban growth patterns in peri-urban areas of Lianyungang City. *Landscape and Urban Planning*, 105, 425-433.
- Sleeter, B. M., Sohl, T. L., Bouchard, M. A., Reker, R. R., Soulard, C. E., Acevedo, W. and Griffith, G. E., et al. (2012). Scenarios of land use and land cover change in the conterminous United States: Utilizing the special report on emission scenarios at ecoregional scales. *Global Environmental Change*.
- Silva, E. A., Ahern, J. and Wileden, J. (2008). Strategies for landscape ecology: An application using cellular automata models. *Progress in Planning*, 70, 133-177.
- Silva, A. M., Huang, C. H., Francesconi, W., Saintil, T. and Vilegas, J. (2015). Using landscape metrics to analyze micro-scale soil erosion processes. *Ecological Indicators* 56, 184–193.
- Sisk, T. D., Margules, C. R. (1993). Habitat edges and restoration: methods for quantifying edge effects and predicting the results of restoration efforts. In

- Nature Conservation 3: Reconstruction of Fragmented Ecosystems, ed. DA Saunders, RJ Hobbs, PR Ehrlich, pp. 57–69. Sydney, Aust.: Surrey, Beatty & Sons.
- Solon, J. (2009). Spatial context or urbanization: Landscape pattern and changes between 1950 and 1990 in the Warsaw metropolitan area, Poland. *Landscape and Urban Planning*, 93, 250-261.
- Soliveres, S., DeSoto, L., Maestre, F.T. and Olano, J.M. (2010). Spatio-temporal heterogeneity in abiotic factors modulate multiple ontogenetic shifts between competition and facilitation. *Perspectives in Plant Ecology, Evolution and Systematics*, 12, 227–234.
- Stratford, J. A. and Stouffer, P. C. (2015). Forest fragmentation alters microhabitat availability for Neotropical terrestrial insectivorous birds. *Biological Conservation*, 188, 109–115.
- Stoate, C., Baldi, A., Beja, P., Boatman, N. D., Herzon, I., van Doorn, A., de Snoo, G. R., Rakosy, L. and Ramwell, C. (2009). Ecological impacts of early 21st century agricultural change in Europe - A review. *Journal of Environmental Management*, 91, 22-46.
- Sun, X., He, J., Shi, Y., Zhu, X. and Li, Y. (2012). Spatiotemporal change in land use patterns of coupled human-environment system with an integrated monitoring approach: A case study of Lianyungang, China. *Ecological Complexity*, 12, 23-33.
- Su, M., Hui, C., Zhang, Y. and Li, Z. (2009). How does the spatial structure of habitat loss affect the eco-epidemic dynamics? *Ecological Modelling*, 12, 51-59.
- Su, S., Jiang, Z., Zhang, Q. and Zhang, Y. (2011). Transformation of agricultural landscapes under rapid urbanization: A threat to sustainability in Hang-Jia-Hu region, China. *Applied Geography*, 31, 439-449.
- Su, W., Gu, C., Yang, G., Chen, S. and Zhen, F. (2010). Measuring the impact of urban sprawl on natural landscape pattern of the Western Taihu Lake watershed, China. *Landscape and Urban Planning*, 95, 61–67.
- Sung, D., Lim, S., Ko, J., and Cho, G. (2001). Scenic evaluation of landscape for urban design purposes using GIS and ANN. *Landscape and Urban Planning*, 56, 75-85.
- Sui, D. Z. and Zeng, H. (2002). Modelling the dynamics of landscape structure in Asia's emerging desakota: a case study in Shenzhen. *Landscape and Urban Planning*, 53, 37-52.
- Termorshuizen, J. W., Opdam, P., and Brink, A. V. D. (2007). Incorporating ecological sustainability into landscape planning. *Landscape and Urban Planning*, 79, 374–384.

- Teixeira, A. M., Soares-Filho, B. S., Freitas, S. R. and Metzger, J. P. (2009). Modeling landscape dynamics in an Atlantic Rainforest region: Implications for conservation. *Forest Ecology and Management*, 257, 1219-1230.
- Tengku Adeline Adura Tengku Hamzah (2004). Landslides hazard and its management in Cameron Highlands, Malaysia: Playing the same old tune? Facing Changing Conditions. Proceedings 2nd Bangi World Conference on Environmental Management. Universiti Kebangsaan Malaysia, Malaysia.
- Terra, T. N., dos Santos, R. F. and Costa, D. C. (2014). Land use changes in protected areas and their future: The legal effectiveness of landscape protection. *Land Use Policy*, 38, 378-387.
- Tigas, L. A., Van Vuren, D. H. and Sauvojat, R. M. (2002). Behavioral responses of bobcats and coyotes to habitat fragmentation and corridors in an urban environment. *Biological Conservation*, 108, 299-306.
- Tjallingii, S. P. (2000). Ecology on the edge: Landscape and ecology between town and country. *Landscape and Urban Planning*, 48, 103-119.
- Tong, S. T. Y., Sun, Y., Ranatunga, T., He, J. and Yang, Y. J. (2012). Predicting plausible impacts of sets of climate and land use change scenarios on water resources. *Applied Geography*, 32, 477-489.
- Tuan Vo, Q., Kuenzer, C., Minh Vo., Moder, F. and Oppelt, N. (2012). Review of valuation methods for mangrove ecosystem services. *Ecological Indicators*, 23, 431-446.
- Turner, M.G., 1990. Spatial and temporal analysis of landscape patterns. *Landscape Ecology*, 4, 21-30.
- Urban, D.L., (1993). Landscape ecology and ecosystem management. In Proc. Of Sustainable Ecological Systems: Implementing an Ecological Approach to Land Management. USDA Forest Service.
- United Nation (2009). Population Division of the Department of Economic and Social Affairs of the United Nations Secretariat. World Population Prospects: The 2008 Revision and World Urbanization Prospects: The 2009 Revision. Available at http://esa.un.org/unpd/wup/Maps/maps_1_2025.htm. Access Date: 15th July 2012.
- Uuemaa, E., Mander, U. and Marja, R. (2013). Trends in the use of landscape spatial matrices as landscape indicators: A review. *Ecological Indicators*, 28, 100-106.
- Verburg, P. H., Sopeboer, W., Veldkamp, A., Limpiada, R., Espaldon, V. and Mastura, S. S. S. (2002). Modelling the spatial dynamic of regional land use. *Environmental Management*, 30, 391-405.

- Vreese, R. D., Leys, M., Fontaine, C. M. and Dendocker, N. (2016). Social mapping of perceived ecosystem service supply – The role of social landscape metrics and social hotspots for integrated ecosystem services assessment, landscape planning and management. *Ecological Indicators*, 66, 517-533.
- Vimal, R., Geniaux, G., Pluvinet, P., Napoleone, C. and Lepar, J. (2012a). Detecting threatened biodiversity by urbanization at regional and local scales using an urban sprawl simulation approach: Application on the French Mediterranean region. *Landscape and Urban Planning*, 104(3-4), 343–355.
- Vimal, R., Mathevet, R. and Thompson, J. D. (2012b). The changing landscape of ecological networks. *Journal for Nature Conservation*, 20(1), 49–55.
- Vuilleumire, S. and Droux, P. R. (2002). Map of ecological networks for landscape planning. *Landscape and Urban Planning*, 58, 157-170.
- Vuilleumire, S. and Fontanillas, P. (2002). Landscape structure affects dispersal in greater white-toothed shrew: Inference between genetic and ecological distance. *Ecological Modelling*, 201, 369-376.
- Voller, J. (1998). Biodiversity and Interior Habitat: The Need to Minimize Edge Effects – Extension Note. Biodiversity Management Concepts in Landscape Ecology. Ministry of Forest Research Program. British Columbia.
- Walters, B. B., Ronnback, P., Kovacs, J. M., Crona, B., Hussain, S. A., Badola, R., Primareva, J. H., Barbier, E. and Dahdouh-Guebes, F. (2008). Ethnobiology, socio-economics and management of mangrove forests: A review. *Aquatic Botany* 89, 220-236.
- Wang, H., Li, X., Long, H., Qiao, Y., and Li, Y. (2011). Development and application of a simulation model for changes in land-use patterns under drought scenarios. *Computers and Geosciences*, 37(7), 831–843.
- Weber, A., Fohrer, N. and Moller, Y. (2001). Long-term land uses changes in a mesoscale watershed due to socio-economic factors-effects on landscape structures and functions. *Ecological Modelling*, 140, 125–140.
- Weng, Y. C. (2007). Spatio-temporal changes of landscape pattern in response to urbanization. *Landscape and Urban Planning*, 81, 341–353.
- White, M. A., Brunsell, N., & Schwartz, M. D. (2003). Vegetation phenology in global change studies. In M. D. Schwartz (Ed.), *Phenology: An integrative environmental science*. New York, NY: Kluwer Academic Publishers, pp. 453–466.
- Wiens, J. A., Horne, B. V and Noon, B. R. (2002). Integrating landscape structure and scale into natural resource management. In :Liu, J. and William, W. T. *Integrating Landscape Ecology into natural Resource Management*. Cambridge University Press. 23-67; 2002.

- Willis, K. J., Jeffers, E. S., Tovar, C., Long, P. R., Caithness, N., Smit, M. G. D. and Hagemann, R. (2012). Determining the ecological value of landscapes beyond protected areas. *Biological Conservation*, 147(1), 3–12.
- With, K.A., 1997. The application of neutral landscape models in conservation biology. *Conservation Biology*, 11, 1069–1080.
- Wollenberg, E., Edmunds, D. and Buck, L. (2000). Using scenarios to make decisions about the future: anticipatory learning for the adaptive co-management of community forest. *Landscape and Urban Planning*, 47, 65-77.
- Wu, K., Ye, X., Qi, Z. and Zhang, H. (2013). Impacts of land use/land cover change and socioeconomic development on regional ecosystem services: The case of fast-growing Hangzhou metropolitan area, China. *Cities*, 31, 276-284.
- Wu, M. Y., Xue, L., Jin, W. B., Xiong, Q. X., Ai, T. C. and Li, B. L. (2012). Modelling the Linkage Between Landscape Metrics and Water Quality Indices of Hydrological Units in Sihui Basin, Hubei Province, China: An Allometric Model. *Procedia Environmental Sciences*, 13, 2131-2145.
- Wu, Q., Li, H., Wang, R. S., Paulussen, J., He, Y., Wang, M., Wang, B. and Wang, Z. (2006). Monitoring and predicting land use change in Beijing using remote sensing and GIS. *Landscape and Urban Planning*, 78, 322–333.
- Xi, F., He, H. S., Clarke, K. C., Hu, Y., Wu, X. and Liu, M. (2012). The potential impacts of sprawl on farmland in Northeast China — Evaluating a new strategy for rural development. *Landscape and Urban Planning*, 104(1), 34–46.
- Xu, L., You, H., Li, D. and Yu, K. (2016). Urban green spaces, their spatial pattern, and ecosystem service value: The case of Beijing. *Habitat International*, 56, 84-95.
- Xun, B., Yu, D., Liu, Y., Hao, R. and Sun, Y. (2014). Quantifying isolation of urban growth on key ecological areas. *Ecological Engineering*, 69, 46-54.
- Yang, W., Li, F., Wang, R. and Hu, D. (2011). Ecological benefits assessment and spatial modeling of urban ecosystem for controlling urban sprawl in Eastern Beijing, China. *Ecological Complexity*, 8(2), 153–160.
- Yeh, C. and Huang, S. (2009). Investigating spatiotemporal patterns of landscape diversity in response to urbanization. *Landscape and Urban Planning*, 93, 151–162.
- Yeh, A. and Li, X. (2003). Simulation of development alternatives using Neural Networks, Cellular Automata, and GIS for Urban Planning. *Photogrammetric Engineering & Remote Sensing*, 69(9), 1043-1052.
- Yue, T. X., Jorgensen, S. E. and Larocque, G. R. (2011). Progress in global ecological modeling. *Ecological Modelling*, 222, 2172-2177.

- Zang, Z., Zou, X., Zuo, P., Song, Q., Wang, C. and Wang, J. (2017). Impact of landscape patterns on ecological vulnerability and ecosystem values. An empirical analysis of Yancheng Nature Reserve in China. *Ecological Indicators*, 72, 142-152.
- Zhao, C., Zhongren, N. and Guodong, C. (2005). Methods for modeling of temporal and spatial distribution of air temperature at landscape scale in the southern Qilian mountains, China. *Ecological Modelling*, 189, 209-220.
- Zheng, D. and Chen, J. (2000). Edge effects in fragmented landscape: a generic model for delineating area of edge influence (D-AEI). *Ecological Modelling*, 132, 175-190.