

MUSHROOM CULTIVATION BY USING AGRICULTURAL WASTES

FATIMAH HAFIFAH BINTI MOHD HANAFI

A thesis submitted in fulfilment of the
requirements for the award of the degree of
Master of Philosophy

School of Civil Engineering
Faculty of Engineering
Universiti Teknologi Malaysia

JULY 2018

*“To my beloved mother, Salmiah binti Abdullah,
father, Mohd Hanafi bin Ab. Rahman,
and my siblings,
Mohd Shakir bin Mohd Hanafi and Shazlin binti Mohd Hanafi
for their everlasting love, support, pray and concern.”*

ACKNOWLEDGEMENT

In the name of Allah, the Most Beneficent, the Most Merciful

I am profoundly grateful to my supervisors, Assoc. Prof. Ir. Dr. Mohd Fadhil Md Din (main), Dr. Shazwin Mat Taib, Prof. Dr. Masahito Yamauchi, Emeritus Prof. Dr. Fumio Yagi, Assoc. Prof. Dr. Sakamoto Mariko, and Assoc. Prof. Dr. Hirofumi Hara for their thoughtful guidance, both intentional and unintentional, on all matters of science and life. I am also grateful to receive financial support from Ministry of Higher Education (MyBrain 15), and Research University Grants (GUP) Tier 2 Vote, Ministry of Higher Education (MOHE) Fundamental Research Grant Scheme Vote No. 10H25.

A million thanks to my fiancé, Mohamad Faizal Bin Mahat, friends and fellow lab-mates especially Ms. Nur Azmira Zainuddin, Mr. Ooi Theam Yiew, Ms. Dianah Mazlan, Mrs. Fatin Asyikin Alias and Mrs. Nur Nadia Farhana Azman for their great support, guidance, encouragement and sincere enthusiasm which helped me to stay on-track.

Finally, I wish to extend my sincere gratitude towards the students in National Institute of Technology Kagoshima College (NITKC), Kagoshima University and Mr. Saiful Amri Suloh from mushroom commercial industry, Saifulam Agrofarm Sdn. Bhd of their thoughtful guidance for the mushroom cultivation techniques.

ABSTRACT

Agricultural waste also known as lignocellulosic residue is a renewable resource that can be used for mushroom cultivation. Empty fruit bunch (EFB) and paddy straw (PS) are categorized as agricultural waste and abundant in every Malaysian palm oil plantation and paddy field. Mushroom cultivation using agricultural waste promises nutritious mushroom and spent mushroom substrate (SMS) that can be used for producing beneficial products such as ruminant feedstock and fertilisers in post-harvest cultivation. SMS is defined as residual biomass generated by commercial mushroom industry after harvesting period. This study aims to reuse agricultural waste generated from the two largest agricultural sectors i.e. palm oil plantation and paddy field, in mushroom cultivation and application of SMS in the post-harvest cultivation which have potential to be used as supplement for ruminant feedstock. The techniques used in this study were named as Process A, B, C and D which consist of drying, grinding, substrates mixing, sterilization, spawning, incubation and harvesting process. Then, the optimum process was used in subsequent cultivation to determine the optimum ratio between ratio A, B, C and D. At the end of harvesting period, SMS was tested for nutrient composition, feeding analysis, hazardous metal composition and bacteriological properties. The results showed novelty in Process D, which obtained optimum yield for three substrates i.e. sawdust, EFB and PS substrates as 232.5 ± 50.3 g, 134.8 ± 82.4 g, and 127.7 ± 25.6 g, respectively. In addition, ratio B comprises 76.0% EFB, 20.0% rice bran and 4.0% agriculture hydrated lime obtained the highest percentage of recovering EFB at the end of mushroom cultivation of 63.2%. Another finding indicated that EFB based-SMS contains adequate nutrients to be applied as supplement for ruminant feedstock when compared to animal feed pellets. Furthermore, EFB-based SMS contains desirable feeding analysis and acceptable amount of *Escherichia coli* which is suitable to be used as ruminant feeding. In summary, this study shows that both, EFB and PS are suitable agricultural waste to be reused in mushroom cultivation, henceforth, reduced the generation of waste in oil palm plantations and paddy fields. EFB was the most applicable substrate to be used for commercialization purposes compared to PS due to high production of mushroom. Furthermore, EFB substrate can be applied further as supplement to ruminant feedstock in the post-harvest cultivation of SMS. Hence, this cycle promotes zero waste discharge. This study can be extended using other abundant agricultural waste in Malaysia such as cocoa or pineapple waste in order to minimise agricultural waste generation.

ABSTRAK

Sisa pertanian juga dikenali sebagai sisa lignoselulosa adalah sumber yang boleh diperbaharui yang mana boleh digunakan untuk penanaman cendawan. Tandan buah kosong (EFB) dan jerami padi (PS) dikategorikan sebagai sisa pertanian dan dihasilkan secara meluas di setiap ladang kelapa sawit dan sawah padi Malaysia. Penanaman cendawan menggunakan sisa pertanian menjanjikan cendawan yang berkhasiat dan sisa substrat cendawan (SMS) yang boleh digunakan untuk menghasilkan produk yang bermanfaat seperti bahan makanan haiwan dan baja dalam pasca penanaman. SMS ditakrifkan sebagai sisa biomas yang dihasilkan oleh industri komersil cendawan selepas tempoh penuaian cendawan. Kajian ini bertujuan untuk menggunakan semula sisa pertanian yang dihasilkan dari dua sektor pertanian terbesar iaitu ladang kelapa sawit dan sawah padi, dalam penanaman cendawan dan aplikasi SMS pada pasca penanaman yang berpotensi digunakan sebagai makanan tambahan untuk bahan makanan haiwan. Teknik yang digunakan dalam kajian ini dinamakan sebagai Proses A, B, C dan D yang terdiri daripada proses pengeringan, pengisaran, pencampuran substrat, pensterilan, pemijahan, pengeraman dan penuaian. Kemudian, proses yang optimum digunakan dalam penanaman berikutnya untuk menentukan nisbah optimum di antara nisbah A, B, C dan D. Pada akhir tempoh penuaian, SMS telah diuji untuk komposisi nutrien, analisis makanan, komposisi logam berbahaya dan sifat bakteriologi. Keputusan mendapati hasil pembaharuan dalam Proses D, dimana hasil optimum dari ketiga-tiga substrat; habuk kayu, EFB dan PS, direkodkan sebagai 232.5 ± 50.3 g, 134.8 ± 82.4 g, dan 127.7 ± 25.6 g. Di samping itu, nisbah B yang terdiri daripada 76.0% EFB, 20.0% dedak beras dan 4.0% kapur pertanian, memperolehi peratusan tertinggi pemulihan EFB pada penghujung penanaman cendawan iaitu 63.2%. Keputusan lain menunjukkan bahawa SMS berasaskan EFB mengandungi nutrien yang mencukupi untuk digunakan sebagai bahan tambah makanan haiwan apabila dibandingkan dengan pelet makanan haiwan. Tambahan pula, SMS berasaskan EFB mengandungi analisis makanan yang dikehendaki dan jumlah *Escherichia coli* yang boleh diterima dalam makanan haiwan. Secara ringkasnya, kajian ini menunjukkan bahawa kedua-dua sisa, EFB dan PS adalah sisa pertanian yang sesuai untuk digunakan semula dalam penanaman cendawan, dan seterusnya, mengurangkan penjanaaan sisa di ladang kelapa sawit dan sawah padi. EFB merupakan substrat yang paling sesuai digunakan untuk tujuan pengkomersilan berbanding PS disebabkan penghasilan cendawan yang tinggi. Tambahan pula, substrat EFB boleh digunakan semula sebagai bahan tambah makanan haiwan dalam pasca penanaman SMS. Oleh itu, kitaran ini menggalakkan pelepasan buangan sifar. Kajian ini boleh diperkembangkan dengan menggunakan sisa pertanian lain yang banyak di Malaysia seperti sisa koko atau nenas untuk meminimumkan penjanaaan sisa pertanian.

TABLE OF CONTENT

CHAPTER	TITLE	PAGE
	DECLARATION	ii
	DEDICATION	iii
	ACKNOWLEDGEMENTS	iv
	ABSTRACT	v
	ABSTRAK	vi
	TABLE OF CONTENTS	vii
	LIST OF TABLES	xiii
	LIST OF FIGURES	xv
	LIST OF ABBREVIATIONS	xvii
	LIST OF APPENDICES	xviii
1	INTRODUCTION	1
	1.1 Background of study	1
	1.2 Problem Statement	4
	1.3 Objectives of the study	5
	1.4 Scope of the study	6
	1.5 Significance of the study	7
2	LITERATURE REVIEW	9
	2.1 Introduction	9
	2.2 Agricultural wastes	10
	2.2.1 Expansion of oil palm in Malaysia	12

2.2.2	Empty fruit bunch (EFB)	13
2.2.3	Paddy straw	15
2.3	Environmental Status in Malaysia	16
2.3.1	Environmental issues due to production of agricultural wastes	16
2.3.2	Disposal of EFB and PS	18
2.3.3	Agricultural waste management	20
2.3.4	Recovery of agricultural wastes	22
2.4	Mushroom	24
2.4.1	Types of mushroom	24
2.4.2	Composition of mushroom fruiting bodies	25
2.4.3	Global mushroom production trends	28
2.5	<i>Pleurotus sajor-caju</i> cultivation techniques	29
2.5.1	Types of mushroom substrates	31
2.6	Spent mushroom substrates (SMS)	33
2.6.1	Composition of spent mushroom substrates (SMS)	34
2.6.2	Application of spent mushroom substrates (SMS)	36
2.6.3	Agro-based spent mushroom substrates (SMS) as animal feedstock	38
2.7	Concluding remarks	41
3	RESEARCH METHODOLOGY	42
3.1	Introduction	42
3.2	Materials and chemical	42
3.3	Apparatus and equipment	44
3.4	Research framework	43
3.5	Preparation of materials for mushroom cultivation	46
3.6	Mushroom cultivation process	47
3.6.1	Moisture content analysis	48
3.6.2	Mixing and packaging	48
3.6.3	Sterilization	50
3.6.4	Inoculation	51
3.6.5	Incubation	52

3.6.6	Mycelia colonization observation	53
3.6.7	Scraping and watering	53
3.6.8	Harvesting process	54
3.6.9	Spent mushroom substrates (SMS) classification	55
3.6.10	Mushroom media composition ratio	56
3.7	Proximate analysis of mushroom fruiting bodies and spent mushroom substrates (SMS)	58
3.7.1	Energy	58
3.7.2	Crude protein	59
3.7.3	Crude fat	59
3.7.4	Ash	60
3.7.5	Crude fiber	60
3.7.6	Nitrogen	61
3.7.7	Potassium (K), phosphorus (P), calcium (Ca), sodium (Na), magnesium (Mg)	61
3.7.8	Calculation of mushroom fruiting bodies	62
3.8	Feeding analysis of spent mushroom substrate (SMS)	62
3.8.1	Neutral-detergent fiber (NDF)	62
3.8.2	Acid-detergent fiber (ADF)	63
3.8.3	Cellulose and lignin	63
3.8.4	Hemicellulose	63
3.9	Hazardous composition and bacteriological analyses in spent mushroom substrate (SMS)	64
4	RESULT AND DISCUSSION	66
4.1	Introduction	66
4.2	Mycelial colonization	67
4.3	Determination of mushroom cultivation process	69
4.3.1	Sawdust (SD) substrate	70
4.3.1.1	Mushroom harvesting interval and flushes	70
4.3.1.2	Mushroom fruiting bodies	71
4.3.1.3	Fresh mushroom yield	72
4.3.2	Empty fruit bunch (EFB) substrate	74
4.3.2.1	Mushroom harvesting interval and flushes	75

4.3.2.2	Mushroom fruiting bodies	76
4.3.2.3	Fresh mushroom yield	77
4.3.3	Paddy straw (PS) substrate	79
4.3.3.1	Mushroom harvesting interval and flushes	80
4.3.3.2	Mushroom fruiting bodies	81
4.3.3.3	Fresh mushroom yield	82
4.3.4	Mushroom performances in Process A, B, C and D of SD, EFB and PS substrates	84
4.3.4.1	Mushroom harvesting period and number of flushes	84
4.3.4.2	Average of mushroom fruiting bodies	86
4.3.4.3	Average of fresh mushroom yield	89
4.3.4.4	Summary on agro-wastes performance as mushroom substrate	91
4.4	Determination of optimum ratio for EFB and PS substrates	94
4.4.1	Mixture of SD and EFB in local ratio	94
4.4.2	Mixture of SD and EFB in modified ratio	97
4.4.3	Mixture of EFB and PS cultivation	99
4.4.4	<i>Pleurotus sajor-caju</i> cultivation by using EFB substrate with variation of rice bran	101
4.5	EFB waste recovery rate in mushroom cultivation	103
4.6	Nutrients composition of mushroom fruiting bodies	106
4.7	Potential application of EFB-based spent mushroom substrate (SMS) as a supplement for ruminant feedstock	111
4.7.1	Animal feed nutrition of EFB-based SMS	111
4.7.2	Feeding analysis of EFB-based SMS	113
4.7.3	Hazardous metal composition of SMS	115
4.7.4	Bacteriological properties of SMS	116
4.7.5	Quantitative calculations for zero waste discharge	117
4.7.5.1	Empty fruit bunch	117
4.7.5.2	Paddy straw	118

5	CONCLUSION AND RECOMMENDATION	120
5.1	Conclusion	120
5.2	Recommendation	122
	REFERENCES	124
	Appendices A-B	144

LIST OF TABLES

TABLE NO.	TITLE	PAGE
2.1	The production rate and waste management of agricultural wastes	20
2.2	Mushroom nutrient composition for various types of species using different substrates	27
2.3	Techniques and ratios involved in mushroom cultivation	30
2.4	Types of substrate and organisms applied in mushroom cultivation	32
2.5	SMS composition and application in various types of mushroom	35
2.6	Potential usage of SMS generation discovered in the previous study	37
2.7	Application of agro-based SMS in feeding trial to ruminants	39
3.1	List of materials and chemical	43
3.2	List of apparatus and equipment	44
3.3	The ratios for comparison of suitable process	57
3.4	Media composition for PS and EFB mixtures	57
3.5	Media composition with different percentage of rice bran	58
3.6	Procedure to determine hazardous chemical composition and bacteriological properties	64
4.1	Mycelial colonization in SD, EFB and PS substrates on 15 th , 30 th and 46 th days during incubation period	68
4.2	SD, EFB and PS-based substrates after end of flushes	68

4.3	Number of fruiting bodies in SD substrate	72
4.4	Amounts of fresh mushroom yield in SD substrate	74
4.5	Number of fruiting bodies for EFB substrate	77
4.6	The amount of fresh mushroom yield of EFB substrate	79
4.7	Number of mushroom fruiting bodies for PS substrate	82
4.8	The amount of fresh mushroom yield by using PS substrates	84
4.9	Number of flushes of the three experimented substrates; SD, EFB and PS	86
4.10	Total mushroom fruiting bodies of the three experimented substrates; SD, EFB and PS	88
4.11	The observation of the mushroom fruiting bodies in SD, EFB and PS substrates	93
4.12	Local media composition for mixture of SD and EFB	95
4.13	Mushroom performances of the local ratio of SD and EFB mixtures	96
4.14	Modified media composition for mixture of SD and EFB	97
4.15	Mushroom performances of the modified ratio of SD and EFB mixtures	99
4.16	Media composition of PS and EFB mixtures	99
4.17	Mushroom performances of PS and EFB mixtures	101
4.18	Media composition with different percentage of rice bran	102
4.19	Mushroom performance of EFB substrate with varies of rice bran	103
4.20	The EFB waste recovered rate	105
4.21	Proximate analysis of animal pellet and EFB substrate	112
4.22	Feeding analysis of EFB-based SMS	114
4.23	Hazardous metal composition of EFB-based SMS	115
4.24	Bacteriological properties of EFB-based SMS	116

LIST OF FIGURES

FIGURE NO.	TITLE	PAGE
2.1	Skeleton of literature review	9
2.2	Palm oil plantations in Malaysia	14
2.3	Empty fruit bunch (EFB)	14
2.4	Empty fruit bunch as mulching in palm oil plantations	15
2.5	Disposal of spent mushroom substrates (SMS) per day	19
3.1	Experimental design layout	45
3.2	The collection of EFB and paddy straw	46
3.3	Preparation of paddy straw substrate	46
3.4	Preparation of EFB substrate	47
3.5	Mixing of mushroom substrates in an exposed area	49
3.6	Mixing of substrates in polyethylene bag	49
3.7	The hole for inoculation purposes	50
3.8	The inoculation in local process	51
3.9	The inoculation process in modified process	52
3.10	The mushroom beds during incubation process	53
3.11	The scraping and watering process	54
3.12	The harvesting process of fruiting bodies	55
3.13	The contaminated mushroom bed	56
3.14	The expired mushroom beds	56
4.1	Mushroom harvesting intervals by using SD substrate	71
4.2	Number of fruiting bodies by using SD substrate	72

4.3	The production of fresh mushroom by using SD substrate	74
4.4	Mushroom harvesting intervals by using EFB substrate	76
4.5	Mushroom fruiting bodies by using EFB substrate	77
4.6	The production of fresh mushroom by using EFB substrate	79
4.7	Mushroom harvesting intervals by using PS as mushroom substrate	81
4.8	Mushroom fruiting bodies by using PS substrate	82
4.9	The production of fresh mushroom by using PS substrate	83
4.10	Mushroom harvesting intervals of the three experimented substrates; SD, EFB and PS	86
4.11	Mushroom fruiting bodies of the three experimented substrates; SD, EFB and PS	88
4.12	Fresh mushroom yield of the three experimented substrates; SD, EFB and PS	90
4.13	EFB waste recoveries in mushroom life-cycle process	105
4.14	The mushroom substrate after lignin degradation by fungi	106
4.15	Energy level	109
4.16	Protein content	109
4.17	Fiber content	109
4.18	Carbohydrates content	109
4.19	Moisture content	109
4.20	Ash content	109
4.21	Magnesium content	110
4.22	Nitrogen content	110
4.23	Potassium content	110
4.24	Phosphorus content	110
4.25	Sodium content	110
4.26	Calcium content	110
4.27	EFB quantitative calculations of zero waste discharge	119
4.28	PS quantitative calculations of zero waste discharge	119

LIST OF ABBREVIATIONS

EFB	Empty fruit bunch
PS	Paddy straw
SMS	Spent mushroom substrate
SD	Sawdust
N	Nitrogen
K	Potassium
P	Phosphorus
CO ₂	Carbon dioxide
Ca	Calcium
Na	Sodium
NDF	Neutral detergent fiber
ADF	Acidic detergent fiber
C	Carbon
Mg	Magnesium
Zn	Zinc
MC	Moisture content
CP	Crude protein
CFT	Crude fat
CF	Crude fiber
CaCO ₂	Lime hydrated agriculture
EE	Ether extract
WRF	White-rot fungi
N/A	Not available

LIST OF APPENDICES

APPENDIX	TITLE	PAGE
A	Moisture content	145
B	Calculation of mushroom fruiting bodies	146
C	List of publications	147
D	Results	148
E	Result of EFB-spent mushroom substrate (EFB-SMS)	155

CHAPTER 1

INTRODUCTION

1.1 Background of Study

Agricultural waste is an abundant raw material of crops dry matter that is easy to obtain. They are also called as lignocellulosic biomass which brings definition of wastes generated in agricultural activities prior process and vice versa. The term “lignocellulosic” is due to the existence of cellulose and hemicellulose in their structure of plant (Mahesh and Mohini, 2013). Interestingly, the lignocellulosic materials not just known as renewable resource, but at the meantime, they are low cost and abundantly available (Sarkar et al, 2012). Typically, agricultural wastes that often generated in the fields were obtained from straws, wheat, paddy, oats, corncob, meanwhile, remnants of plantation activities were most of the time come from oil palm wastes; empty fruit bunch (EFB), trunks, palm kernel shells, and pineapple (Adebayo, 2015; Nicolcioiu et al. 2016). These wastes are known possess high nutrient levels of nitrogen (N), potassium (K), and phosphorus (P) which able to increase crop yields (Elly, 2011).

Embrandiri (2013) in his review stated that, palm oil and paddy plantation are among of important agricultural products which spurred the growth of economic condition in most ASEAN countries including Malaysia, Thailand, and Indonesia. The oil palm industry has become a significant icon to Malaysia and Indonesia as both countries become the major contributors in world's palm oil production (Sudirman et al., 2011). Furthermore, Malaysia as a tropical climate country experiencing mild humid weather throughout the year, has encouraged the growth

and development of the oil palm plantation (Pei et al., 2012). These agricultural activities can play important role in stabilizing the climate change and act as food security for the countries (Schaffnit-chatterjee et al., 2011). Every mill of oil palm generated solid waste such as EFB, palm oil trunks, palm fiber and palm kernel shell. Oil palm plantations produced 1.27 tons of EFB per hectare per year of the surplus mill (Elbersen et al., 2013).

Other sectors that also become the larger contributor for agricultural waste in Malaysia is rice paddy plantation-Malaysia recorded 730.0 thousand hectares of rice crop by 2015 (DOSM, 2016) and this expected to keep expanding along with other contributors in the agricultural sector throughout the year. Typically, in Malaysia, paddy field are double-cropping throughout the year, during off season and main season. From this cultivation process, about 80% of rice straw is produced by improper disposal management, which results in a series of pollution (Shafie et al., 2014a). The paddy straw disposal is usually done through open burning known as the cheapest and easiest way to prepare the field for the next cultivation season (Rosmiza et al., 2014; Shafie et al., 2014b).

Waste disposal was introduced into the plantation and enforced under the Environmental Quality (Amendment) Act 2012 to control the open burning and open waste dumping that contributed to the release of carbon dioxide (CO₂) and methane (CH₄) into the atmosphere (MPOB, 2015). Conventionally, from previous practice, EFB will either incinerated or applied on top of soil as mulch, a direct composting system for plantations (Ali et al., 2013). Wan and Surya (2012) in their study stated, the mulching was able to improve soil conditions by maintaining soil moisture and has also been applied to cover crops for oil palm plantation.

An alternative method and application to recycle these wastes is seen as so important in order to recover this waste resource through its value proposition toward zero waste discharge. Reuse of agricultural wastes is more worthwhile as this not only provides another new application but also minimizes the waste generated in the field. Thus, this step can protect the environment from pollution. This waste recovery also contributes significantly other industries as it introduces low-cost primary

source and potentially producing profitable products in future. Therefore, a number of researches on reuse agricultural wastes have been successfully carried out such as fertilisers, animal feedstock, and coal tailing (Elly, 2011; Mahesh and Mohini, 2013; Pattanayak, 2015).

Agricultural wastes can also be converted into valuable food through mushroom cultivation (Lalithadevy and Many, 2014). Since two decades ago, paddy straw was introduced and widely used as substrate media in mushroom cultivation of *P. sajor-caju*, *P. platypus* and *P. citrinopileatus* species (Ragunathan et al., 1996; Gurudevan, 2012). EFB, paddy straw, wheat straw, sugarcane bagasse, and maize cob were reported as great mushroom substrates media which produced higher yield compared to present practices substrates, sawdust (Chukwurah, 2012; Gurudevan, 2012; Kavitha et al., 2013). Out of listed media, paddy straw is the growth media that provided the shortest duration to complete mycelium colonization compared to other substrates (Lalithadevy and Many, 2014).

Additionally, the medium-scale mushroom industry is capable of producing approximately 13.6 million tonnes per year of spent mushroom substrate (SMS) after the harvest cycle (Phan and Sabaratnam, 2012). In Korea, SMS production is estimated about 2 million tons yearly, with almost 1.2 million tons generated from *Pleurotus sajor-caju* (Lim et al., 2013). SMS defined as biomass generation from mushroom industry remaining after a crop of mushroom is harvested (Moon et al., 2012).

In fact, SMS contains enough digestible nutrition for ruminant feeding and reuse of SMS can protect the environment as well as increase growers' income, (Danny et al., 2004). As reported by Khattab et al., (2013), paddy straw-SMS can be forages for ruminants and possess the high possibility of replacing practices ruminant feedstock at high levels of up to 50% or 90% from diets. Besides, lignocellulosic characteristics in mushroom substrate suitable to be used as animal feed due to its upgraded properties and digestible carbohydrates (Georgios et. al., 2014; Kuijk et al., 2015). Both targeted waste residues; EFB and paddy straw, contain high protein and carbohydrates which is very suitable to be used as mushroom media for *Pleurotus*

spp. production and supplement for ruminant feedstock (Silvana et al., 2006; Kowalski et al., 2014; Wyngaard et al., 2015).

Therefore, this study is purposely to reuse and recover agricultural wastes; EFB and paddy straw through mushroom media as to produce edible *Pleurotus* spp. and recover again the residues produced from cultivation of mushrooms called as SMS as supplement for ruminant animals.

1.2 Problem Statement

Oil palm and paddy plantation are the main important commodity product that become as transformation agent to the scenario of agricultural sector and economy in Malaysia (DOSM, 2016). The presence of wastes from oil palm plantation has created a major disposal problem such as open burning and *in situ* dumping. Currently, most of EFB are used as soil mulching as organic fertiliser to the plantation, otherwise, dumped in the same manner as palm oil mill effluent (POME) (Pei et al., 2012).

According to available literature, production of paddy straw is over 13 tons per hectares and reported mostly in North of Malaysia (Perlis, Kedah, Penang, and Perlis) and Central of Malaysia (Selangor, Negeri Sembilan, and Malacca) (Shafi et al., 2013c). Presently, paddy straw is disposed of by open burning which sparked lots of environmental issues, hence affecting weather and local communities (Rosmiza et al., 2014). An alternative application of agricultural wastes is needed in order to minimize the waste generation and protect the environment from being polluted.

Moreover, series of problems also arises in mushroom industries whereby generation and management of spent mushroom substrate (SMS) become a big challenge to the farmers. The application of SMS is still lacking as the current practices substrate which is sawdust that containing low nutrient composition is still become an option, hence limiting the potential its usage.

Various efforts have been spent to increase the potential of SMS application in the agricultural industry, given the use of present substrate; sawdust substrate is not suitable to be used as animal feedstock and fertilisers, and it ends up dumped and burned *in situ* (Park et al., 2012). Hence, to minimize the problem, sawdust substrate should be substituted with lignocellulosic substrates such as paddy straw, palm wastes, or crop wastes, as to minimize the wastes from mushroom industry.

Another problem arises in the mushroom industries is the cultivation techniques of *Pleurotus* spp. which caused high contamination rate and put the farmers in the worrisome state. Mushroom commercial industries in Ulu Tiram, and Pontian, Johor, experienced approximately 20% and 14%, respectively, contamination rate in every 1000 beds per production. Contamination occurs probably because of the techniques and handling methods used are unhygienic. Besides, *Pleurotus* spp. is categorized as fungi and very prone to contamination. Contamination rate must be reduced by improving the process in the mushroom cultivation.

Most of the studies conducted in Malaysia only emphasized on a single output, for instance, mushroom cultivation by using agricultural wastes (Ali, et. al., 2013). There are limitations in research especially on the use of agricultural wastes as mushroom substrate, the percentage of waste recovered from the process and utilization of agro wastes-SMS as ruminant feedstock. Besides, the study of minimizing contamination rate in mushroom cultivation process also still scarce.

1.3 Objectives of The Study

The main purpose of this research is to recycle and recover agricultural wastes generated from two (2) largest agricultural sectors; palm oil plantation and paddy field, through mushroom life-cycle. The following are the objectives to achieve the research aim:

- i. To study the performance of reuse agricultural wastes; empty fruit bunch and paddy straw in *Pleurotus* spp. cultivation by comparing mushroom cultivation process,
- ii. To determine the percentage of waste recovery by using optimum ratio of recycled agro-wastes substrates; empty fruit bunch and paddy straw,
- iii. To compare the nutrient composition of mushroom fruiting bodies to the three substrates; sawdust, empty fruit bunch and paddy straw,
- iv. To investigate the composition and feeding analysis of spent mushroom substrates (SMS) as a supplement for ruminant feedstock in order to promote zero waste discharge.

1.4 Scope of The Study

This study focused on reuse and recovery agricultural wastes mainly from two (2) largest agricultural sectors; palm oil plantation and paddy field through mushroom cultivation of well-known species; *Pleurotus* spp.,. The study was started from the observation of problem arises in plantation and mushroom industries. Then, continued to the laboratory set up to study the performance of *Pleurotus* spp. on the recycled agro-wastes substrates and its optimum ratio of agro-wastes. The parameter of mushroom harvesting interval, mushroom fruiting bodies and mushroom yield of three substrates; sawdust (SD), empty fruit bunch (EFB) and paddy straw (PS) were investigated and percentage of waste recovered was calculated.

Furthermore, this study compared the nutrient composition of *Pleurotus* spp. fruiting bodies of these three experimented substrates; SD, EFB and PS in order to determine the most nutritious mushroom. In the post-harvest cultivation, composition and feeding analysis of chosen substrate (EFB-SMS) were examined to analyze the compatibility as a supplement for ruminant feedstock.

Controllable parameters in the study were moisture content, substrates (SD, EFB, and PS) and additional nutrients used in cultivation (rice bran and lime) and *Pleurotus sajor-caju* spawn.

1.5 Significance of The Study

The significance of this study is to utilize abandoned waste from palm oil plantations and paddy field as mushroom substrate in order to reduce the generation of agricultural wastes from the prohibited activities; burning and open dumping. Besides, these wastes can be used as a bio-remediation solution in SMS production for ruminant feedstock (Foluke et al., 2014; Phan and Sabaratnam, 2012).

Recently, under the Economic Transformation Program (ETP), both the oil palm and biomass industries have been highlighted as the nation's premier niche National Key Economic Areas (NKEAs) (JPM, 2013). The utilization of palm biomass is increasing significantly over time, which creates a symbiotic situation where the "previous waste" serves as the input for other industries, leading the palm oil industry to a zero waste path (Pei et al., 2012). Hence, Malaysian Economic Transformation Programme (ETP) encourages the utilization of oil palm and biomass for another application in order to promote zero waste paths.

Mushroom cultivation is well-known as the easiest way to reduce generation of agricultural wastes, moreover, the result obtained in previous study indicated that high yield of mushroom can be produced by using agricultural wastes as mushroom substrate (Ali et al., 2013; Kavitha et al., 2013; Marlina et al., 2015). Furthermore, some techniques to improve local mushroom cultivation process are provided in this study in order to minimize contamination rate due to unhygienic handling and techniques. The result obtained in this study showed significant differences in terms of harvesting interval, flushes, production of fruiting bodies, fresh mushroom yield, and nutrients of mushroom compared to local practices cultivation. An optimum ratio and nutrient composition of SD, EFB and PS substrates is provided.

Moreover, this study provides an alternative to recycle and recover abundant of agricultural waste to another beneficial products. Reuse and recovering agricultural wastes in mushroom life-cycle can be concluded as promoting a zero waste discharge; hence, this is important to the farmers to practice in agriculture sector. In addition, this study hopes to help the country in controlling environmental pollution through the zero waste initiatives, thereby enabling Malaysia to take a step further towards sustaining sustainable growth.

REFERENCES

- Abdel-Aziz, N. A., Salem, A. Z., El-Adawy, M. M., Camacho, L. M., Kholif, A. E., Elghandour, M. M., & Borhami, B. E. (2015). *Biological treatments as a mean to improve feed utilization in agriculture animals—An overview*. *Journal of Integrative Agriculture*, 14(3), 534-543.
- Abdullah, N., and Sulaiman, F. (2013). The oil palm wastes in Malaysia. *Biomass Now – Sustainable Growth and Use*. 75–100.
- Adebayo, E. A. and Martinez-Carrera, D. (2015). Oyster mushrooms (*Pleurotus*) are useful for utilizing lignocellulosic biomass. *African Journal of Biotechnology*. 14(1), 52–67.
- Adedokun, O. M. (2014b). *Oyster mushroom: Exploration of additional agro-waste Substrates in Nigeria*. *International Journal of Agricultural Research*, 9(1), 55-59.
- Adedokun, O. M., and Orluchukwu, J. A. (2013a). *Pineapple: organic production on soil amended with spent mushroom substrate*. *Agr Biol JN Am*, 4, 590-593.
- Adegoke, A. E., Kola, O. J., Yadav, A., and Chandral, B. T. (2012). *Improvement of laccase production in *Pleurotus pulmonarius*-LAU 09 by mutation*. *Journal of Microbiology Research*, 2(1), 11-17.
- Ahlawat, O. P., and Tewari, R. P. (2007). *Cultivation technology of paddy straw mushroom (*Volvariella volvacea*)*, 213, 36.
- Ahlawat, O. P., and Sagar, M. P. (2007). *Management of spent mushroom substrate*. National Research Centre for Mushroom, Indian Council of Agricultural Research.
- Aida, F. M. N. A., Shuhaimi, M., Yazid, M., and Maaruf, A. G. (2009). *Mushroom as a potential source of prebiotics: a review*. *Trends in food science & technology*, 20(11), 567-575.

- Alam, N., Amin, R., Khan, A., Ara, I., Shim, M. J., Lee, M. W., and Lee, T. S. (2008). *Nutritional analysis of cultivated mushrooms in Bangladesh- Pleurotus ostreatus, Pleurotus sajor-caju, Pleurotus florida and Calocybe indica*. Mycobiology, 36(4), 228-232.
- Alananbeh, K. M., Bouqellah, N. A., and Kaff, N. A. (2014). *Cultivation of oyster mushroom Pleurotus ostreatus on date-palm leaves mixed with other agro-wastes in Saudi Arabia*. Saudi Journal of Biological Sciences. 21(6), 616-625.
- Alborés, S., Pianzola, M. J., Soubes, M., and Cerdeiras, M. P. (2006). *Biodegradation of agroindustrial wastes by Pleurotus spp for its use as ruminant feed*. Electronic Journal of Biotechnology, 9(3), 0-0.
- Alder, V. G., Brown, A. M., and Gillespie, W. A. (1966). *Disinfection of heat-sensitive material by low-temperature steam and formaldehyde*, 19(1), 83–89.
- Ali, N., Tabi, A. N. M., Zakil, F. A., Mohd Fauzai, W. N. F., and Hassan, O. (2013). *Yield performance and biological efficiency of empty fruit bunch (EFB) and palm pressed fibre (PPF) as substrates for the cultivation of pleurotus ostreatus*. Jurnal Teknologi (Sciences and Engineering). 64(1), 93–99.
- Amerah, A. M. (2015). *Interactions between wheat characteristics and feed enzyme supplementation in broiler diets*. Animal Feed Science and Technology, 199, 1-9.
- Amin, M. Z. M., Harun, A., and Wahab, M. A. M. A. (2014). *Status and potential of mushroom industry in Malaysia*. Economic and Technology Management Review, 9b, 103-111.
- Amiruddin, M. N., Faizah, S., Ayat, K. A. R., and Basiron, Y. (2005). *Impact of commercialization of biofuel on prices of related commodities*. Kuala Lumpur: MPOB.
- AOAC. (1995). *Official methods of analysis* (16th Ed.). Arlington VA, USA; Association of Official Analytical Chemists.
- Awalludin, M., Sulaiman, O., and Hashim, R. (2015). *An overview of the oil palm industry in Malaysia and its waste utilization through thermochemical conversion, specifically via liquefaction*. Renewable and Sustainable Energy Review. 50, 1469-1484.
- Arvanitoyannis, I. and Tserkezou, P. (2008). *Corn and rice waste: A comparative and critical presentation of methods and current and potential uses of treated waste*. International Journal of Food Science and Technology. 43(6), 958-988.

- Ayala, M. et al., (2011). *Fibrolitic potential of spent compost of the mushroom *Agaricus bisporus* to degrade forages for ruminants*. African Journal of Microbiology Research, 5(3), 241–249.
- Ayhan Demirbas. (2018). Waste management, waste resource facilities and waste conversion process. Energy Conversion and Management. 1280-1287.
- Babasaki, K., Masuno, K., and Murata, H. (2003). *Interactions of heterologous mycelia colonized in the substrate govern fruit body production in the cultivated homobasidiomycete *Pholiota nameko**. Bioscience, biotechnology, and biochemistry, 67(1), 100-106.
- Bakker, R. R., Jenkins, B. M., and Williams, R. B. (2002). *Fluidized bed combustion of leached rice straw*. Energy & fuels, 16(2), 356-365.
- BAM, F., Hunt, J., Abeyta, C., and Tran, T. (1998). *FDA Bacteriological Analytical Manual*. Chapter 4
- BAM, F., Hunt, J., Abeyta, C., and Tran, T. (1998). *FDA Bacteriological Analytical Manual*.
- Bandyopadhyay, S. (2006). *Source composite curve for waste reduction*. Chemical Engineering Journal, 125(2), 99-110.
- Barros, L., Cruz, T., Baptista, P., Leticia, M. E., and Isabel, C. F. R. F. (2008). *Wild and comemrcial mushrooms as source of nutrients and nutraceuticals*. Food and Chemical Toxicology. 46(8), 2742-2747.
- Baysal, E., Peker, H., Yalinkiliç, M. K., and Temiz, A. (2003). *Cultivation of oyster mushroom on waste paper with some added supplementary materials*. Bioresource Technology. 89(1), 95-97.
- Berne, S., Pohleven, J., Vidic, I., Rebolj, K., Pohleven, F., Turk, T., and Sepčić, K. (2008). *Ostreolysin enhances fruiting initiation in the oyster mushroom (*Pleurotus ostreatus*)*. Mycological Research, 111(12), 1431–1436.
- Binod, P., Sindhu, R., Singhanian, R. R., Vikram, S., Devi, L., Nagalakshmi, S., Kurien, N., Sukumaran, R. K., and Pandey, A. (2010). *Bioethanol production from rice straw: An overview*. Bioresource Technology. 101(13), 4767-4774.
- Bisaria, R., Vasudevan, P., and Bisaria, V. S. (1990). *Utilization of spent agro-residues from mushroom cultivation for biogas production*. Applied Microbiology and Biotechnology. 33, 607-609.
- Çağlarirmak, N. (2007). *The nutrients of exotic mushrooms (*Lentinula edodes* and*

- Pleurotus species*) and an estimated approach to the volatile compounds. *Food Chemistry*. 105(3), 1188-1194.
- Carabajal, M., Laura, L., Edgardo, A., and Bernardo, L. (2012). *Effect of co-cultivation of two Pleurotus species on lignocellulolytic enzyme production and mushroom fructification*. *International Biodeterioration & Biodegradation*. 66(1), 71-76.
- Chae, H.J. and Ahn, J.H. (2013). *Optimization of rice bran and food waste compost contents in mushroom culture medium to maximize mycelial growth rate and fruit body yield of Pleurotus ostreatus*. *International Biodeterioration and Biodegradation*, 80, 66–70.
- Chang, K. L., Chen, X. M., Sun, J., Liu, J. Y., Sun, S. Y., Yang, Z. Y., and Wang, Y. (2017). *Spent mushroom substrate biochar as a potential amendment in pig manure and rice straw composting processes*. *Environmental technology*, 38(13-14), 1765-1769.
- Chaturvedi, V. and Verma, P., (2013). *An overview of key pretreatment processes employed for bioconversion of lignocellulosic biomass into biofuels and value added products*. *3 Biotech*, 3(5), 415–431.
- Childers, D. L., Corman, J., Edwards, M., and Elser, J. J. (2011). *Sustainability challenges of phosphorus and food: solutions from closing the human phosphorus cycle*. *Bioscience*, 61(2), 117-124.
- Chukwurah, N. F. (2012). *Performance of oyster mushroom (Pleurotus ostreatus) in different local agricultural waste materials*. *African Journal of Biotechnology*. 11(37), 8979–8985.
- Cunha, T. J., and McDowell, L. R. (2012). *Nutrition of grazing ruminants in warm climates*. Academic Press. 22-25
- Danny L. Rinker, and ZERI, S. W. K. (2004). *Recycling of spent oyster mushroom substrate*. *Mushroom Grower's Handbook* 1.
- Das, N. and Mukherjee, M. (2007). *“Cultivation of Pleurotus Ostreatus on Weed Plants.”* *Bioresource Technology* 98(14): 2723–26.
- De Carvalho, C. S. M., Sales-Campos, C., De Andrade, M. C. N. (2010). *Mushrooms of the Pleurotus genus: A review of cultivation techniques*. *nterciencia*. 35(3), 177-182.
- Dobermann. A. and Fairhurst, T. H. (2002). *Rice straw management*. *Better Crops*

- International. 16(5), 7-11.
- DOSM. (2016). Department of Statistics Malaysia Press Release, (December). Retrieved from <https://www.dosm.gov.my>
- Dundar, A., Acay, H., & Yildiz, A. (2009a). *Effect of using different lignocellulosic wastes for cultivation of Pleurotus ostreatus (Jacq.) P. Kumm. on mushroom yield, chemical composition and nutritional value*. African Journal of Biotechnology, 8(4).662-666.
- Dundar, A., Acay, H., and Yildiz, A. (2008b). Yield performances and nutritional contents of three oyster mushroom species cultivated on wheat stalk. *Journal of Biotechnology*, 7(19), 3497–3501.
- Elbersen, W., Meesters, K., and Bakker, R. (2013). *Valorization of palm oil (mill) residues*. Netherlands.
- Elly Sabiiti. (2011). *Utilising Agricultural Waste to Enhance Food Security and Conserve the Environment*. African Journal of Food, Agriculture, Nutrition and Development, 11(6), 1–9.
- Embrandiri, A., Ibrahim, M. H., and Singh, R. P. (2013). *Palm Oil Mill Wastes Utilization; Sustainability in the Malaysian Context*. International Journal of Scientific and Research Publications, 3(1), 2250–3153.
- Ezeibekwe, I.O, Ogonnaya, C.I, Unamba, C.I.N and Osuala, O. . (2009). *Proximate Analysis and Mineral Composition of Edible Mushrooms in Parts of South Eastern Nigeria*. Report and Opinion, 1(4), 32–36.
- Falandysz, J., 2013. Review: *On published data and methods for selenium in mushrooms*. Food Chemistry, 138(1), 242–250.
- Farhat A. Avin. (2017). Growing a mushroom business. The Star Online. Available online on 8th November 2017.
- FAO. (2014). Food and agriculture organization of the United Nations. www.fao.org/.
- Fazaeli, H., Shafyee-Varzeneh, H., Farahpoor, A., and Moayyer, A. (2014). *Recycling of mushroom compost wheat straw in the diet of feedlot calves with two physical forms*. International Journal of Recycling of Organic Waste in Agriculture, 3(3),3.
- Finney, K. N., Ryu, C., Sharifi, V. N., and Swithenbank, J. (2009). *The reuse of spent mushroom compost and coal tailings for energy recovery: comparison of thermal treatment technologies*. Bioresource technology, 100(1), 310-315.

- Foluke, A., Olutayo, A., and Olufemi, A. (2014). *Assessing spent mushroom substrate as a replacement to wheat bran in the diet of broilers*. American International Journal of Contemporary Research, 4(4), 178–183.
- Fraga, C. G. (2005). *Relevance, essentiality and toxicity of trace elements in human health*. Molecular aspects of medicine, 26(4), 235-244.
- Gao, W., Liang, J., Pizzul, L., Feng, X. M., Zhang, K., and del Pilar Castillo, M. (2015a). *Evaluation of spent mushroom substrate as substitute of peat in Chinese biobeds*. International Biodeterioration & Biodegradation, 98, 107-112.
- Gao, L., Yang, H., Wang, X., Huang, Z., Ishii, M., Igarashi, Y., and Cui, Z. (2008b). *Rice straw fermentation using lactic acid bacteria*. Bioresource Technology, 99(8), 2742–2748. <http://doi.org/10.1016/j.biortech.2007.07.001>
- García-Delgado, C., D’Annibale, A., Pesciaroli, L., Yunta, F., Crognale, S., Petruccioli, M., and Eymar, E. (2015b). *Implications of polluted soil biostimulation and bioaugmentation with spent mushroom substrate (Agaricus bisporus) on the microbial community and polycyclic aromatic hydrocarbons biodegradation*. Science of the Total Environment, 508, 20-28.
- Garg, V.K., Gupta, R., 2009. *Vermicomposting of Agro-Industrial Processing Waste*. In: *Biotechnology for Agro-Industrial Residues Utilization*. Springer, Netherlands, pp. 431–456.
- Gbolagade, J., Ajayi, A., Oku, I., and Wankasi, D., (2006). *Nutritive value of common wild edible mushrooms from Southern Nigeria*. Global Journal of Biotechnology & Biochemistry. 1(1), 16-21.
- Georgios, K., Konstantinos, C. M., Iordanis, C., and Georgios, I. Z. (2014). *Bioconversion of lignocellulosic residues by Agrocybe cylindracea and Pleurotus ostreatus mushroom fungi - Assessment of their effect on the final product and spent substrate properties*. Food Chemistry. 161, 127-135.
- Gimeno, A., Al Alami, A., Toral, P. G., Frutos, P., Abecia, L., Fondevila, M., and Castrillo, C. (2015). *Effect of grinding or pelleting high grain maize-or barley-based concentrates on rumen environment and microbiota of beef cattle*. Animal Feed Science and Technology, 203, 67-78.
- Glass, N. L., Carolyn, R., Roca, M. G., and Read, N. D. *Hyphal homing, fusion and mycelial interconnectedness*. Trends in Microbiology, Elsevier. 12(3), 135-141.
- Graminha, E., Gonçalves, A., Pirota, R. D. P. B., Balsalobre, M. A. A., Silva, R. D.

- and Gomes, E. (2008). *Enzyme production by solid-state fermentation: Application to animal nutrition*. *Animal Feed Science and Technology*. 144(1-2), 1-22.
- Gregori, A., Švagelj, M., Pahor, B., Berovič, M., and Pohleven, F. (2008). *The use of spent brewery grains for Pleurotus ostreatus cultivation and enzyme production*. *New Biotechnology*, 25(2-3), 157–161.
- Gurudevan, T. (2012). *Improved techniques to enhance the yield of paddy straw mushroom (Volvariella volvacea) for commercial cultivation*. *African Journal of Biotechnology*. <http://doi.org/>
- Harith, N., Abdullah, N., and Sabaratnam, V. (2014). *Cultivation of Flammulina velutipes mushroom using various agro-residues as a fruiting substrate*. *Pesquisa Agropecuária Brasileira*, 49(3), 181-188.
- Hartley, C. W. S. (1967). *The oil palm*.
- He, Y., Pang, Y., Liu, Y., Li, X., and Wang, K. (2008). *Physicochemical characterization of rice straw pretreated with sodium hydroxide in the solid state for enhancing biogas production*. *Energy & Fuels*. 22(4), 2775-2781.
- Heriansyah. (1991). *Optimizing the use of oil palm by-product (EFB) as fertilizer supplement for oil palm*. 1-11.
- Herman, T., Murchie, E. H., and Warsi, A. A. (2015). *Rice production and climate change: A case study of Malaysian rice*. *Pertanika Journal of Tropical Agricultural Science*. 38(3), 321-328.
- Hernández, D., Sánchez, J.E. and Yamasaki, K., 2003. *A simple procedure for preparing substrate for Pleurotus ostreatus cultivation*. *Bioresource Technology*, 90(2), pp.145–150.
- Herrero-Hernández, E., Andrades, M. S., Rodríguez-Cruz, M. S., and Sánchez-Martín, M. J. (2011). *Effect of spent mushroom substrate applied to vineyard soil on the behaviour of copper-based fungicide residues*. *Journal of environmental management*, 92(7), 1849-1857.
- Ho, H. T., Wang, C. L., and Wang, C. H. (2015). *The effects of different substrates on the growth, yield, and nutritional composition of two oyster mushrooms (Pleurotus ostreatus and Pleurotus cystidiosus)*. *Mycobiology*, 43(4), 423–434.
- Hu, B. B., and Zhu, M. J. (2017). *Enhanced hydrogen production and biological saccharification from spent mushroom compost by Clostridium thermocellum*

- 27405 supplemented with recombinant β -glucosidases. *International Journal of Hydrogen Energy*, 42(12), 7866-7874.
- Imran, M., Anwar, Z., Irshad, M., Asad, M. J., and Ashfaq, H. (2016). *Cellulase Production from Species of Fungi and Bacteria from Agricultural Wastes and Its Utilization in Industry : A Review*. *Advances in enzyme Research*. 4, 44-55.
- Islam, M. Z., Rahman, M. H., and Hafiz, F. (2009). *Cultivation of oyster mushroom (Pleurotus flabellatus) on different substrates*. *Int. J. Sustain. Crop Prod*, 4(1), 45-48.
- Jabatan, P. M. (2013). *Program Transformasi Ekonomi "Laporan Tahunan 2012."* Jabatan Perdana Menteri.
- Jafari, M. A., Nikkhah, A., Sadeghi, A. A., and Chamani, M. (2007). *The effect of Pleurotus spp. fungi on chemical composition and in vitro digestibility of rice straw*. *Pakistan Journal of Biological Sciences*.
- Jami, E. and Mizrahi, I., (2012). *Composition and similarity of bovine rumen microbiota across individual animals*. *PLoS ONE*, 7(3), 1–8.
- Jayakumar, T. et al., (2011). *In-vitro and in-vivo antioxidant effects of the oyster mushroom Pleurotus ostreatus*. *Food Research International*, 44(4), 851–861.
- Jr, A.B.V., Kalaw, S. P., Nitural, P. S., Abella, E. A., and Reyes, R. G. (2006). *Cultivation of Thai and Japanese strains of Pleurotus sajor – caju on rice straw-based Volvariella volvacea mushroom spent and composted rice straw in Central Luzon Region , Philippines*. *Journal of Agricultural Technology*. 69–75.
- Juris, B., Yahya, J., Mait, Kriipsalu, Z., Vincevica-Gaile, Z., Fabio, K., Gunita, C., Ruta, O., Laine, R., Vita, R., Marika, H., Arturs, V., Kaur-Mikk, P., William, H., Maris, K. (2018). *On the way to 'zero waste' management: Recovery potential of elements, including rare earth elements, from fine fraction of waste*. *Journal of Cleaner Production*. 81-90.
- Kavitha, B., Rajannan, G., and Jothimani, P. (2013). *Utilization of Empty Fruit Bunch of Oil Palm as Alternate Substrate for The Cultivation of Mushroom*. *International Journal of Science*, 2(5), 839–846.
- Kerdsuwan, S., and Laohalidanond, K. (2010). *Renewable energy from palm oil empty fruit bunch*. 29.
- Khan, M.W., Ali, M. A., Khan, N. A., Khan, M. A., Rehamn, A., and Javed, N. (2013). *Effect of different levels of lime and pH on mycelial growth and*

- production efficiency of oyster mushroom (Pleurotus spp)*. Pak.J.Bot, 45(1), 297–302.
- Khattab, H.M. , Gado H. M., Salem A.Z.M., Camacho L.M., El-Sayed, M.M., Kholif, A.M., El-Shewy, A. A. and. K. A. E. (2013). *Chemical Composition and In Vitro Digestibility of Rice Straw Treated with Pleurotus ostreatus, Pleurotus pulmonarius and Pleurotus tuber-regium*. Slovak J. Anim. Sci, 45(1), 14–20.
- Kim, Y. I., Cho, W. M., Hong, S. K., Oh, Y. K., and Kwak, W. S. (2011a). *Yield, nutrient characteristics, ruminal solubility and degradability of spent mushroom (Agaricus bisporus) substrates for ruminants*. Asian-Australasian Journal of Animal Sciences. 24(11), 1560-1568.
- Kim, Y. I., Lee, Y. H., Kim, K. H., Oh, Y. K., Moon, Y. H., and Kwak, W. S. (2012b). *Effects of supplementing microbially-fermented spent mushroom substrates on growth performance and carcass characteristics of Hanwoo steers (a field study)*. Asian-Australasian journal of animal sciences, 25(11), 1575.
- Kim, J. S., Lee, Y. H., Kim, Y. I., Ahmadi, F., Oh, Y. K., Park, J. M., and Kwak, W. S. (2016c). *Effect of microbial inoculant or molasses on fermentative quality and aerobic stability of sawdust-based spent mushroom substrate*. *Bioresource technology*, 216, 188-195.
- Kim, Y. I., and Kwak, W. S. (2012d). *Isolation and identification of lactic acid bacteria from spent mushroom substrate for silage making and determination of optimal medium conditions for growth*. Journal of Animal Science and Technology, 54(6), 435-442.
- Kossmann, R. J., Peterson, D. C., and Andrews, H. L. (1965). *Studies in neuromuscular disease I. Total body potassium in muscular dystrophy and related diseases*. Neurology, 15(9), 855-855.
- Kowalski, Z., Ludwin, J., Gorka, P., Rinne, M., Weisbjerg, M. R., and Jagusiak, W. (2014). *The use of cellulase and filter abg technique to predict digestibility of forages*. Animal Feed Science and Technology. 198, 49-56.
- Kuijk, V. S. J. a., Sonnenberg, A. S. M., Baars, J. J. P., Hendriks, W. H., and Cone, J. W. (2015). *Fungal treated lignocellulosic biomass as ruminant feed ingredient: A review*. Biotechnology Advances. 33(1), 191-202.
- Koutrotsios, G., Mountzouris, K. C., Chatzipavlidis, I., and Zervakis, G. I. (2014). *Bioconversion of lignocellulosic residues by Agrocybe cylindracea and*

- Pleurotus ostreatus* mushroom fungi–Assessment of their effect on the final product and spent substrate properties. *Food chemistry*, 161, 127-135.
- Kwak, W. S., Jung, S. H., and Kim, Y. I. (2008a). *Broiler litter supplementation improves storage and feed-nutritional value of sawdust-based spent mushroom substrate*. *Bioresource technology*, 99(8), 2947-2955.
- Kwakman, P. H., te Velde, A. A., de Boer, L., Speijer, D., Vandenbroucke-Grauls, C. M., and Zaat, S. A. (2010). *How honey kills bacteria*. *The FASEB Journal*, 24(7), 2576-2582.
- Lalithadevy, V., and Many, J. N. (2014). *Yield performance of fruits and vegetables peels as substrates for the cultivation of oyster mushroom (Pleurotus florida)*, 1(1).
- Lau, H. L., Wong, S. K., Bong, C. F. J., and Rabu, A. (2014). *Suitability of oil palm empty fruit bunch and sago waste for Auricularia polytricha cultivation*. *Asian Journal of Plant Sciences*, 13(3), 111-119.
- Lee, C. Y., Park, J. E., Kim, B. B., Kim, S. M., and Ro, H. S. (2009). *Determination of mineral components in the cultivation substrates of edible mushrooms and their uptake into fruiting bodies*. *Mycobiology*, 37(2), 109-113.
- León, B. D., Gómez-Aldapa, C. A., Rangel-Vargas, E., Vázquez-Barrios, E., and Castro-Rosas, J. (2013). *Frequency of indicator bacteria, Salmonella and diarrhoeagenic Escherichia coli pathotypes on ready-to-eat cooked vegetable salads from Mexican restaurants*. *Letters in applied microbiology*, 56(6), 414-420.
- Li, X., Pang, Y., and Zhang, R. (2001). *Compositional changes of cottonseed hull substrate during P. ostreatus growth and the effects on the feeding value of the spent substrate*. *Bioresource Technology*. 80(2), 157-161.
- Liang, Z. C., Wu, C. Y., Shieh, Z. L., and Cheng, S. L. (2009). *Utilization of grass plants for cultivation of Pleurotus citrinopileatus*. *International Biodeterioration & Biodegradation*, 63(4), 509-514.
- Lim, S. H., Lee, Y. H., and Kang, H. W. (2013). *Efficient recovery of lignocellulolytic enzymes of spent mushroom compost from oyster mushrooms, Pleurotus spp., and potential use in dye decolorization*. *Mycobiology*, 41(4), 214–220.

- Lin, Y., Ge, X., and Li, Y. (2014a). *Solid-state anaerobic co-digestion of spent mushroom substrate with yard trimmings and wheat straw for biogas production*. *Bioresource technology*, 169, 468-474.
- Liu, Z., Xu, A., and Zhao, T. (2011a). *Energy from combustion of rice straw: Status and challenges to China*. *Energy and Power Engineering*. 3(3), 325-331.
- Liu, F., Ooi, V. E. C., and Chang, S. T. (1997b). *Free radical scavenging activities of mushroom polysaccharide extracts*. Elsevier. 60(10), 763-771.
- Loss, E., Royer, A. R., Barreto-Rodrigues, M., and Barana, A. C. (2009). *Use of maize wastewater for the cultivation of the Pleurotus spp. mushroom and optimization of its biological efficiency*. *Journal of Hazardous Materials*. 166(2-3), 1522-1525.
- Lou, Z., Sun, Y., Zhou, X., Baig, S. A., Hu, B., and Xu, X. (2017a). *Composition variability of spent mushroom substrates during continuous cultivation, composting process and their effects on mineral nitrogen transformation in soil*. *Geoderma*, 307, 30-37.
- Mahaffey, K. R., Capar, S. G., Gladen, B. C., and Fowler, B. A. (1981). *Concurrent exposure to lead, cadmium, and arsenic*. *The Journal of laboratory and clinical medicine*, 98(4), 463-481.
- Mahesh, M. S., and Mohini, M. (2013). *Biological treatment of crop residues for ruminant feeding : A review*. *African Journal of Biotechnology*, 12(27), 4221–4231.
- Macias-Corral, M., Zohrab, S., Adrian, H., Geoffrey, S., Paul, F., Hui, Y., and John, L., (2008). *Anaerobic digestion of municipal solid waste and agricultural waste and the effect of co-digestion with dairy cow manure*. *Bioresource Technology*. 99(17), 8288-8293.
- Mamman, M. and Folorunsho, J.O. 2016. “*Assessment of open burning of agricultural waste in Biu Local Government Area of Borno State , Nigeria.*” *Advances in Social Sciences Research Journal*. 3(6): 52–60.
- Marlina, L., Sukotjo, S., and Marsudi, S. (2015). *Potential of Oil Palm Empty Fruit Bunch (EFB) as Media for Oyster Mushroom, Pleurotus ostreatus Cultivation*. *Procedia Chemistry*, 16, 427–431.
- Mazaheri, H, Lee, K.T., Bhatia, S., and Mohamed, A. R. (2010). *Subcritical water liquefaction of oil palm fruit press fiber for the production of bio-oil: Effect of catalysis*. *Bioresource Tehcnology*. 101(23), 9335-9341.

- Medina, E. et al., (2009). *Spent mushroom substrates as component of growing media for germination and growth of horticultural plants*. Bioresource Technology, 100(18), pp.4227–4232.
- Mekhilef, S., Siga, S., and Saidur, R. (2011). *A review on palm oil biodiesel as a source of renewable fuel*. Renewable and Sustainable Energy Reviews. 15(4), 1937-1949.
- Miles, P. G., and Chang, S. T. (2004). *Mushrooms: cultivation, nutritional value, medicinal effect, and environmental impact*. CRC press.
- Montañez-Valdez, O. D., Avellaneda-Cevallos, J. H., Guerra-Medina, C. E., Reyes-Gutiérrez, J. A., Peña-Galeas, M. M., Casanova-Ferrín, L. M., and del Carmen, R. (2015). *Chemical Composition and Ruminal Disappearance of Maize Stover Treated with Pleurotus Djamor*. Life Science Journal, 12(2s).
- Moonmoon, M., Uddin, M. N., Ahmed, S., Shelly, N. J., and Khan, M. A. (2010a). *Cultivation of different strains of king oyster mushroom (Pleurotus eryngii) on saw dust and rice straw in Bangladesh*. Saudi Journal of Biological Sciences, 17(4), 341-345.
- Moonmoon, M., Shelly, N. J., Khan, M. A., Uddin, M. N., Hossain, K., Tania, M., and Ahmed, S. (2011b). *Effects of different levels of wheat bran, rice bran and maize powder supplementation with saw dust on the production of shiitake mushroom (Lentinus edodes (Berk.) Singer)*. Saudi Journal of Biological Sciences, 18(4), 323–328.
- Moon, Y. H., Shin, P. G., and Cho, S. J. (2012). *Feeding Value of Spent Mushroom (Pleurotus eryngii). Substrate*. Journal of Mushroom Science and Production, 10(4), 236–243.
- Nadhim, Mustafa, Idham Ali, Sajid Salahuddin, and Saleem Al-saeedi. (2017). *Applicable Properties of the Bio-Fertilizer Spent Mushroom Substrate in Organic Systems as a Byproduct from the Cultivation of Pleurotus Spp*. Information Processing in Agriculture 4(1): 78–82.
- Naraian, R. et al. (2009). *Influence of Different Nitrogen Rich Supplements during Cultivation of Pleurotus Florida on Corn Cob Substrate*. Environmentalist 29(1): 1–7.
- Nicolcioiu, M. B., Popa, G., and Matei, F. (2016). *Mushroom mycelia cultivation on different agricultural waste substrates*. Scientific Bulletin. Series F. Biotechnologies.

- Nur Aziera Abd Rasib, Zarina Zakaria, Mohammad Fahrurrazi Tompong, Ridzwan Abdul Rahman, and Hakimah Othman. (2015). *Characterization of Biochemical Composition for Different Types of Spent Mushroom Substrates in Malaysia*. Malaysian Journal of Analytical Sciences 19(1): 41–45.
- Nurhayati, A. and Fauziah, S. (2013). *The properties of the washed empty fruit bunches of oil palm*. Journal of Physical Science. 24(2), 117-137.
- Oei, P., Hui, Z., Jianhua, L., Jianqing, D., Meiyuan, C., and Yi, C. (2007). *The alternative uses of spent mushroom compost*. ECO Consult Foundation, Tiel, Netherlands.
- Olisa, Y. and Kotingo, K. (2014). *Utilization of palm empty fruit bunch (PEFB) as solid fuel for steam boiler*. European Journal of Engineering and Technology. 2(2), 1-7.
- Ouzouni, P. K., Petridis, D., Wolf-Dietrich, K., and Kyriakos, A. R. (2016). *Nutritional value and metal content of wild edible mushrooms collected from West Macedonia and Epirus, Greece*. Food Chemistry. 115 (4), 1575-1580.
- Pant, D., Reddy, U. G., and Adholeya, A. (2006). *Cultivation of oyster mushrooms on wheat straw and bagasse substrate amended with distillery effluent*. World Journal of Microbiology and Biotechnology, 22(3), 267-275.
- Paredes, C., Medina, E., Bustamante, M. A., and Moral, R. (2016). *Effects of spent mushroom substrates and inorganic fertilizer on the characteristics of a calcareous clayey-loam soil and lettuce production*. Soil Use and Management, 32(4), 487-494.
- Park, J. H., Kim, S. W., Do, Y. J., Kim, H., Ko, Y. G., Yang, B. S., and Cho, Y. M. (2012). *Spent mushroom substrate influences elk (Cervus Elaphus Canadensis) hematological and serum biochemical parameters*. Asian-Australasian Journal of Animal Sciences, 25(3), 320–324.
- Pathmashini, L., Arulnandhy, V. and Wijeratnam, R.S.W. (2008). *Cultivation of Oyster Mushroom (Pleurotus Ostreatus) on Sawdust*. Ceylon Journal of Science (Biological Science). 37(2), 177–182.
- Paula, F. S., Tatti, E., Abram, F., Wilson, J., and O'Flaherty, V. (2017). *Stabilisation of spent mushroom substrate for application as a plant growth-promoting organic amendment*. Journal of Environmental Management. 196, 476-486.
- Pei, W., Ng, Q., Loong, H., Yuen, F., Kamal, M., Heng, J., and Lim, E. (2012). *Waste-to-wealth : green potential from palm biomass in Malaysia*. Journal of

- Cleaner Production, 34(September 2011), 57–65.
- Peng, J.-T., Lee, C.-M., and Tsai, Y.-F. (2000). *Effect of Rice Bran on the Production of Different King Oyster Mushroom Strains During Bottle Cultivation*.
- Phan, C. W., & Sabaratnam, V. (2012). Potential uses of spent mushroom substrate and its associated lignocellulosic enzymes. *Applied Microbiology and Biotechnology*, 96(4), 863–873.
- Poulose, A. M., Elnour, A. Y., Anis, A., Shaikh, H., Al-Zahrani, S. M., George, J., ... & Sarmah, A. K. (2018). *Date palm biochar-polymer composites: An investigation of electrical, mechanical, thermal and rheological characteristics*. Science of The Total Environment, 619, 311-318.
- Purnomo, A. S., Mori, T., Kamei, I., Nishii, T., and Kondo, R. (2010). *Application of mushroom waste medium from Pleurotus ostreatus for bioremediation of DDT-contaminated soil*. International Biodeterioration & Biodegradation, 64(5), 397-402.
- Qiao, J. J., Zhang, Y. F., Sun, L. F., Liu, W. W., Zhu, H. J., and Zhang, Z. (2011). *Production of spent mushroom substrate hydrolysates useful for cultivation of Lactococcus lactis by dilute sulfuric acid, cellulase and xylanase treatment*. Bioresource technology, 102(17), 8046-8051.
- Quaicoe, E.H. Amoah, C. Oboda, M., and Odamntten, G. T. (2014). *Nutrient Requirements And Environmental Conditions For The Cultivation Of The Medicinal Mushroom (Lentinula Edodes) (Berk.) In Ghana*. International Journal of Scientific & Technology Research, 3(12), pp.45–50.
- Quartey, E. T. and Chýlková, J. (2012). *Challenges and opportunities in managing agricultural waste*. Advances in Environment Biotechnology and Biomedicine. 235-239.
- Radke, S. M., and Alocilja, E. C. (2005). *A high density microelectrode array biosensor for detection of E. coli O157: H7*. Biosensors and Bioelectronics, 20(8), 1662-1667.
- Ragunathan, R., Gurusamy, R., Palaniswamy, M., and Swaminathan, K. (1996). *Cultivation of Pleurotus spp. on various agro-residues*. Food Chemistry, 55(2), 139–144.
- Rahman, M. M., Alam, M. R., Amin, M. R., and Das, N. G. (2010). *Comparative study of the nutritive values of the different varieties of rice straw*. Bangladesh

- Journal of Animal Science. 39(1&2), 75-82.
- Raikwar, M. K., Kumar, P., Singh, M., and Singh, A. (2008). *Toxic effect of heavy metals in livestock health*. Veterinary world, 1(1), 28-30.
- Ramírez-Bribiesca, J.E. et al., 2010. *Influence of Pleurotus ostreatus spent corn straw on performance and carcass characteristics of feedlot Pelibuey lambs*. Indian Journal of Animal Sciences, 80(8), pp.754–757.
- Reis, F. S., Barros, L., Martins, A., and Ferreira, I. C. (2012). *Chemical composition and nutritional value of the most widely appreciated cultivated mushrooms: an inter-species comparative study*. Food and Chemical Toxicology, 50(2), 191-197.
- Reyes, R. (2000). *Indoor cultivation of paddy straw mushroom, Volvariella volvacea, in crates*. Mycologist. 14(4), 174-176.
- Rezaei, J., Rouzbehan, Y., Zahedifar, M., and Fazaeli, H. (2015). *Effects of dietary substitution of maize silage by amaranth silage on feed intake, digestibility, microbial nitrogen, blood parameters, milk production and nitrogen retention in lactating Holstein cows*. Animal Feed Science and Technology, 202, 32-41.
- Ribas, L.C.C. et al., 2009. *Use of spent mushroom substrates from Agaricus subrufescens (syn. A. blazei, A. brasiliensis) and Lentinula edodes productions in the enrichment of a soil-based potting media for lettuce (Lactuca sativa) cultivation: Growth promotion and soil bioremediati*. Bioresource Technology, 100(20), pp.4750–4757.
- Rizki, M., and Tamai, Y. (2011). *Effects of different nitrogen rich substrates and their combination to the yield performance of oyster mushroom (Pleurotus ostreatus)*, (68), 1695–1702.
- Rosmiza, M., Rosniza, A. C., Mazdi, M., and Jabil, M. (2014). *Farmers' knowledge on potential uses of rice straw: An assessment*. Malaysian Journal of Society and Space, 5(5), 30–43.
- Ryu, J. S., Kim, M. K., Im, C. H., and Shin, P. G. (2015). *Development of cultivation media for extending the shelf-life and improving yield of king oyster mushrooms (Pleurotus eryngii)*. Scientia Horticulturae, 193(2), 121–126.

- Saidu, M., Salim, M. R., and Yuzir, M. A. M. (2011). *Cultivation of oyster mushroom (Pleurotus spp.) on palm oil mesocarp fibre*. African Journal of Biotechnology, 10(71), 15973-15976.
- Sánchez, C. (2009). *Lignocellulosic residues: biodegradation and bioconversion by fungi*. Biotechnology advances, 27(2), 185-194.
- Sarkar, N., Ghosh, S. K., Bannerjee, S., and Aikat, K. (2012). Bioethanol production from agricultural wastes: An overview. *Renewable Energy*, 37(1), 19–27.
- Sarnklong, C., Cone, J. W., Pellikaan, W., and Hendriks, W. H. (2010). *Utilization of rice straw and different treatments to improve its feed value for ruminants: a review*. Asian-Australasian Journal of Animal Sciences, 23(5), 680-692.
- Schaffnit-chatterjee, A. C., Kahn, B., Editor, D., Schneider, S., Assistant, T., Peter, M., and Mayer, M. T. (2011). Mitigating climate change through agriculture. <http://doi.org/>
- Seaby, D. A. (1996). *Investigation of the epidemiology of green mould of mushroom (Agaricus bisporus) compost caused by Trichoderma harzianum*. Plant pathology, 45(5), 913-923.
- Sendi, H., Mohamed, M. T. M., Anwar, M. P., and Saud, H. M. (2013). *Spent mushroom waste as a media replacement for peat moss in Kai-Lan (Brassica oleracea var. Alboglabra) production*. The Scientific World Journal, 2013. <https://doi.org/10.>
- Shafie, S. M., Mahlia, T. M. I., and Masjuki, H. H. (2013c). *Life cycle assessment of rice straw co- fi ring with coal power generation in Malaysia*. Energy, 57, 284–294.
- Shafie, S. M., Masjuki, H. H., and Mahlia, T. M. I. (2014b). *Life cycle assessment of rice straw-based power generation in Malaysia*. Energy, 70, 401–410.
- Shafie, S. M., Masjuki, H. H., and Mahlia, T. M. I. (2014a). *Rice straw supply chain for electricity generation in Malaysia: Economical and environmental assessment*. Applied Energy, 135, 299–308.
- Shahzad, F., Abdullah, M., Chaudhry, A. S., Bhatti, J. A., Jabbar, M. A., Ahmed, F., ... and Irshad, I. (2016). *Effects of varying levels of fungal (Arachniotus sp.) treated wheat straw as an ingredient of total mixed ration on growth performance and nutrient digestibility in Nili Ravi Buffalo calves*. Asian-Australasian journal of animal sciences, 29(3), 359.

- Sher, H., Al-Yemeni, M., and Khan, K. (2011). *Cultivation of the oyster mushroom (Pleurotus ostreatus (Jacq.) P. Kumm.) in two different agroecological zones of Pakistan*. African Journal of Biotechnology, 10(2), 183-188.
- Shumaila, G. and Mahapara, S. (2009). *Proximate composition and mineral analysis of cinnamon*. Pakistan Journal of Nutrition. 8(9), 1456-1460.
- Sidana, A., and Farooq, U. (2014). *Sugarcane bagasse: a potential medium for fungal cultures*. Chinese Journal of Biology, 2014, 1-5.
- Singh, C.S., 2009. *Recycling of spent oyster mushroom substrate to recover additional value*. Journal of Science, Engineering and Technology.
- Singh, M. P., and V. K. Singh. 2012. "Biodegradation of Vegetable and Agrowastes by Pleurotus Sapidus: A Novel Strategy to Produce Mushroom with Enhanced Yield and Nutrition." Cellular and Molecular Biology, 58(1): 1-7.
- Silvana, A., Maria, J. P., Matilde, S., Maria, P. C. (2006). *Biodegradation of agroindustrial wastes by Pleurotus spp for its use as ruminant feed*. Electronic Journal of Biotechnology, 9(3), 215-220.
- Simard, R. R. (1993). *Ammonium acetate-extractable elements*. In: Carter, M.R. (Ed.), *Soil Sampling and Methods of Analysis*. Canadian Society of Soil Sciences, pp. 39-42.
- Sindhu, N., and Shehrawat, P. (2015). *Agricultural waste utilization for healthy environment and sustainable lifestyle*. Annals of Agri Bio Research. 20(1), 110-114.
- Singh, M. P., and Kumar, V. (2012). *Biodegradation of vegetable and agrowastes by Pleurotus sapidus: A noble strategy to produce mushroom with enhanced yield and nutrition*. Cell. Mol. Biol, 58(1), 1-7.
- Sonali, D. R. (2012). *Cultivation and study of growth of oyster mushroom on different agricultural waste substrate and its nutrient analysis*. Advances in Applied Science Research. 3(4), 1938-1949.
- Song, Q., Li, J., & Zeng, X. (2015). *Minimizing the increasing solid waste through zero waste strategy*. Journal of Cleaner Production, 104, 199-210.
- Stamets, Paul. (1993). *Growing gourmet and medicinal mushrooms*. Ten Speed Press. New York.
- Sudirman, L. I., Sutrisna, A., Listiyowati, S. R. I., Fadli, L., and Tarigan, B. (2011). *The potency of oil palm plantation wastes for mushroom production*, 378-384.
- Sugimoto, H. H., Barbarosa, A. Dekker, R., Castro-Gomez, R. J. (2001) *Veratryl*

- alcohol stimulates fruiting body formation in the oyster mushroom, Pleurotus ostreatus*. FEMS microbiology letters, 194(2), 235–238.
- Sumathi, S., Chai, S., and Mohamed, A. (2008). *Utilization of oil palm as a sources of renewable energy in Malaysia*. Renewable and Sustainable Energy Reviews. 12(9), 2404-2421.
- Sung, C. T. B., Goh, K. J., and Khairun, N. K. (2010). *Physical changes to oil palm empty fruit bunches (EFB) and EFB mat (Ecomat) during their decomposition in the field*. Pertanika Journal of Tropical Agricultural Science. 33(1), 39-44.
- Thy, P. Jenkins, B. M., Williams, R. B., Lesher, C. E.c, and Bakker, R. R. (2010). *Bed agglomeration in fluidized combustor fueled by wood and rice straw blends*. Fuel Processing Technology. 91(11), 1464-1485.
- Toptas, A., Demierege, S., Mavioglu Ayan, E., and Yanik, J. (2014). *Spent mushroom compost as biosorbent for dye biosorption*. CLEAN–Soil, Air, Water, 42(12), 1721-1728.
- Tuhy, L., Samoraj, M., Witkowska, Z., Wilk, R., and Chojnacka, K. (2015a). *Using spent mushroom substrateas the base for organic-mineral micronutrient fertilizer–field tests on maize*. BioResources, 10(3), 5709-5719.
- Upadhyay, R. C., Verma, R. ., Singh, S. ., and Yadav, M. C. (2002). *Effect of organic nitrogen supplementation in Pleurotus species*. The 4 Th ICMBMP, (February), 1–11.
- Van Soest PJ, Robertson JB, Lewis BA (1991). *Methods for dietary fibre, neutral detergent fibre, and non-starch carbohydrates in relation to animal nutrition*. J. Dairy Sci., 74: 3583-3597.
- Veena, S. S., and Pandey, M. (2011). *Paddy Straw as a Substrate for the Cultivation of Lingzhi or Reishi Medicinal Mushroom, Ganoderma lucidum (W.Curt. :Fr.) P. Karst. in India*. International Journal of Medicinal Mushrooms, 13(4), 397–400.
- Wales, A. D., Allen, V. M., and Davies, R. H. (2010). *Chemical treatment of animal feed and water for the control of Salmonella*. Foodborne pathogens and disease, 7(1), 3-15.
- Wan, A. I. and Surya, P. C. (2012). *Converting Waste Oil Palm Trees Into a Resource*. Osaka, Japan: United Nations Programme, International Environmental Technology Centre, Osaka, Japan. <http://doi.org/>

- Wang, S., Xu, F., Li, Z., Zhao, S., Song, S., Rong, C., and Liu, Y. (2015). *The spent mushroom substrates of Hypsizigus marmoreus can be an effective component for growing the oyster mushroom Pleurotus ostreatus*. *Scientia Horticulturae*, 186, 217-222.
- Watt, B. K., and Merrill, A. L. (1975). *Composição química de cogumelos comestíveis (Pleurotus spp.) originados de cultivos em diferentes substratos*. *Ciência e Tecnologia de Alimentos* 15(2): 189-193.
- Williams, B.C., McMullan, J.T. and McCahey, S. (2001). *An initial assessment of spent mushroom compost as a potential energy feedstock*. *Bioresource Technology*, 79(3), 227–230.
- Wisitrassameewong, K., Samantha, C. K., Naritsada, T., and Ruilin, Z. (2012). *Review: Agaricus subrufescens*. *Saudi Journal of Biological Sciences*. 19(2), 131-146.
- Witte, F., Hort, N., Vogt, C., Cohen, S., Kainer, K. U., Willumeit, R., and Feyerabend, F. (2008). *Degradable biomaterials based on magnesium corrosion*. *Current opinion in solid state and materials science*, 12(5), 63-72.
- Wyngaard, V., Meeske, J. D. V. R., and Erasmus, L. J. (2015). *Effect of palm kernel expeller as supplementation on production performance of Jersey cows grazing kikuyu-ryegrass pasture*. *Animal Feed Science and Technology*, 199, 29-40.
- Yang, D., Liang, J., Wang, Y., and Sun, F. (2016). *Tea waste: An effective and economic substrate for oyster mushroom cultivation*. *Journal of the Science of Food and Agriculture*, 96(2), 680–684.
- Yodkhum, S., Sampattagul, S., & Gheewala, S. H. (2018). *Energy and environmental impact analysis of rice cultivation and straw management in northern Thailand*. *Environmental Science and Pollution Research*, 1-11.
- Zadrazil, F., Kamra, D. N., Isikhuemhen, O. S., Schuchardt, F., and Flachowsky, G. (1996). *Bioconversion of Lignocellulose into Ruminant Feed with White Rot Fungi—Review of Work Done at the FAL, Braunschweig*. *Journal of Applied Animal Research*, 10(2), 105–124.
- Zaffrie, M., Amin, M., and Harun, A. (2014). “*Status and Potential of Mushroom Industry in Malaysia*.” 9: 103–11.
- Zhang, R., Li, X. and Fadel, J. (2002). *Oyster mushroom cultivation with rice and wheat straw*. *Bioresource Technology*, 82(3), 277–284.

- Zhu, H. J., Liu, J. H., Sun, L. F., Hu, Z. F., and Qiao, J. J. (2013c). Combined alkali and acid pretreatment of spent mushroom substrate for reducing sugar and biofertilizer production. *Bioresource technology*, *136*, 257-266.
- Zhu, H. J., Sun, L. F., Zhang, Y. F., Zhang, X. L., and Qiao, J. J. (2012a). Conversion of spent mushroom substrate to biofertilizer using a stress-tolerant phosphate-solubilizing *Pichia farinose* FL7. *Bioresource technology*, *111*, 410-416.
- Zhu, H., Sheng, K., Yan, E., Qiao, J., and Lv, F. (2012b). Extraction, purification and antibacterial activities of a polysaccharide from spent mushroom substrate. *International Journal of Biological Macromolecules*, *50*(3), 840-843.
- Zhu, Y., Chang, Y., Guan, J., Shangguan, G., and Xin, F. (2016d). Butanol production from organosolv treated spent mushroom substrate integrated with in situ biodiesel extraction. *Renewable Energy*, *96*, 656-661
- Ziyad Tareq Aldoori, Ahmed Sinan Ahmed Al-Obaidi, Abdullah Hasan Abdulkareem, M.K.A., (2015). *Effect of Dietary Replacement of Barley with Mushroom Cultivation on Carcass Characteristics of Awassi Lambs*. *Journal of Animal Health and Production*, *3*(4), pp.94–98.
- ZWIA, 2015.. Zero Waste International Alliance (ZWIA). Zero Waste Definition. <http://zwia.org/standards/zw-definition>. accessed 26.08.2015.