

MATHEMATICAL MODEL OF LEAD TRANSFER IN HUMAN BODY

MOHD SALLEHUDDIN BIN YAHYA

A dissertation submitted in partial fulfilment of the
requirements for the award of the degree of
Master of Science (Mathematics)

Faculty of Science
Universiti Teknologi Malaysia

AUGUST 2015

To my beloved mother and father

Acknowledgement

I would like to express my deep gratitude to Associate Professor Dr. Shamsuddin Ahmad, my research supervisor, for his patient guidance, enthusiastic encouragement and useful critiques of this research work. I would also like to thank to Dr. Yeak Su Hoe and Graduate School staffs, for their advice and assistance in keeping my progress on schedule. Finally, I wish to thank my parents and friends for their support and encouragement throughout my study.

Abstract

A mathematical model of lead transfer in human body is developed by using compartmental model based on previous studies and formulated in mathematical terms to calculate and predict the level of lead in human body. The Batschelet's three-compartmental model being used in this study is adopted from the Rabinowitz *et al.* (1976). The experimental data adopted in this study is drawn from the published researches to test the mathematical model for solving the real biological system illustrated by the Batschelet's three-compartmental model and to ascertain the mathematical model fit the experimental data. Some examples of a real biological system of lead transfer in human body are studied such as the amount of lead and the equilibrium state of lead level stored in several major parts of human body. A mathematical model derived from the compartmental model which involves first-order ordinary differential equations is solved analytically by using a mathematical technique which utilize some basic concepts from the algebra of matrices. The mathematical general solution obtained is employed by using Microsoft Excel program to produce tables and plot graphs.

Abstrak

Model bermatematik bagi peralihan jisim plumbum dalam tubuh manusia dibangunkan dengan menggunakan model berpetak berdasarkan kajian lepas dan dirumus dalam sebutan bermatematik untuk menghitung dan meramal aras jisim plumbum dalam tubuh manusia. Model berpetak-tiga Batschelet dalam kajian ini mengguna pakai model yang dipelopori oleh Rabinowitz *et al.* (1976). Data kajian yang digunakan pula diperoleh daripada kajian lepas untuk diuji pada model bermatematik tersebut bagi menyelesaikan sistem biologi sebenar yang diterjemahkan oleh model berpetak-tiga Batschele dan untuk menentusahkan sama ada model bermatematik tersebut bersesuaian dengan data kajian. Beberapa contoh sistem biologi sebenar bagi peralihan jisim plumbum dalam tubuh manusia yang boleh dikaji ialah amoun plumbum dan takat keseimbangan aras plumbum tersimpan dalam beberapa bahagian utama tubuh manusia. Model bermatematik yang diterbitkan daripada model berpetak tersebut mengandungi persamaan-persamaan terbitan biasa darjah-pertama diselesaikan secara analitik dengan teknik bermatematik yang melibatkan beberapa konsep asas matriks aljabar. Penyelesaian umum bermatematik yang diperoleh digembleng dengan mengguna pakai Microsoft Excel untuk menghasilkan jadual dan memplot graf.

TABLE OF CONTENTS

CHAPTER	TITLE	PAGE
	DECLARATION	ii
	DEDICATION	iii
	ACKNOWLEDGEMENTS	iv
	ABSTRACT	v
	ABSTRAK	vi
	TABLE OF CONTENTS	vii
	LIST OF TABLES	ix
	LIST OF FIGURES	x
	LIST OF SYMBOLS	xii
	LIST OF APPENDICES	xiii
1	INTRODUCTION	1
	1.1 Background of The Problem	1-2
	1.2 Statement of The Problem	2
	1.3 Objectives of The Study	3
	1.4 Scope of The Study	3
	1.5 Significance of The Study	3-4
2	LITERATURE REVIEWS	5
	2.1 Introduction	5
	2.2 Lead in Human Body	5-8
	2.3 Compartmental Model Theories	9-10
	2.4 Compartmental Model in Studies of Lead in Human Body	11-12
	2.5 Conclusion	12-13

TABLE OF CONTENTS

CHAPTER	TITLE	PAGE
3	RESEARCH METHODOLOGY	14
3.1	Introduction	14
3.2	Research Methodology Flow Chart	15
3.3	Batschelet's Three-Compartmental Model	16-17
3.4	Mathematical Model of Batschelet's Three-Compartmental Model	18-22
3.5	Identification of The Model	22-25
3.6	An ODE System (A Continuous Model)	26-42
3.7	Conclusion	42-43
4	RESULTS AND DISCUSSIONS	44
4.1	Introduction	44
4.2	Results	45-46
4.2.1	Lead Environment	46-58
4.2.2	Non-Lead Environment	58-72
4.3	Discussions	72-73
4.4	Conclusion	73
5	CONCLUSION AND RECOMMENDATIONS	74
5.1	Introduction	74
5.2	Conclusion of Study	75
5.3	Recommendations	76
	REFERENCES	77-79
	Appendices A-E	80-101

LIST OF TABLES

TABLE NO.	TITLE	PAGE
3.1	Amount of lead in compartment x_i daily exchange rate of lead y_{ij} , daily lead output rate, z_i , daily lead input rate,	24
3.2	Values of coefficients in (9), (12), (13), (14) and (15)	25
4.1	Distribution of lead at interval of 10 days in lead	48
4.2	Distribution comparison of lead levels in each compartment for $\alpha=49\mu\text{g/d}$ (represent by b , d and f) and $98\mu\text{g/d}$ (represent by a , c and e)	56
4.3	Distribution of lead at interval of 10 days in non-lead environment	62
4.4	Distribution of lead at interval of 10 days in non-atmospheric lead environment	70

LIST OF FIGURES

FIGURE NO.	TITLE	PAGE
3.1	The Batschelet's Three-Compartmental Model	15
1	Lead level during lead intake in blood compartment	49
2	Lead level during lead intake in soft tissues compartment	49
3	Lead level during lead intake in bones compartment	50
4	Lead level for blood, soft tissues and bones compartment	50
5	Lead level during lead intake in bones compartment after 1000 days	
6	Dynamics of lead in blood compartment if the daily lead input from air increases from 49 micrograms to 98 micrograms	57
7	Dynamics of lead in soft tissues compartment if the daily lead input from air increases from 49 micrograms to 98 micrograms	57
8	Dynamics of lead in bones compartment if the daily lead input from air increases from 49 micrograms to 98 micrograms	58
9	Lead level in blood compartment from day 400 after taking a massive medication	64
10	Lead level in soft tissues compartment from day 400 after taking a massive medication	64
11	Lead level in bones compartment from day 400 after taking a massive medication	65
12	Lead level in blood, soft tissues and bones compartments from day 400 after taking a massive medication	65

LIST OF FIGURES

FIGURE NO.	TITLE	PAGE
13	Lead level in blood, soft tissues and bones compartments from day 400 in non-atmospheric lead environment	72

LIST OF SYMBOLS

D	-	lead intake
$x_i(t)$	-	amount of lead in compartment i at time t
$x_i'(t)$	-	lead transfer rate in compartment i at time t
α	-	rate of lead enters the lungs
β	-	rate of lead enters the digestive system
y_{ij}	-	rate of lead transfer from compartment i to j

LIST OF APPENDICES

APPENDIX NO.	TITLE	PAGE
A	Diagrams of compartmental model and graphical results of lead transfer in human body from previous researches.	80
B	Sample of Calculation of Lead Level in Human Body in Atmospheric Lead Environment	83
C	Sample of Calculation of Lead Level in Human Body in Non-Atmospheric Lead Environment	90
D	Sample of Calculation of Lead Level in Human Body When Lead Concentration is Doubled	94
E	Sample of Calculation of Lead Level in Human Body When Lead Concentration is Decreased To 65.5%	98

CHAPTER 1

INTRODUCTION

1.1 Background of The Problem

Lead is a toxic metal and it is closely related to human life. It is used in some products that could be found around us such as paint and cosmetic (Moore et al., 1980, and Paoliello and Capitani, 2007). It is also a natural occurring chemical element contained in soil. It is estimated that the level of lead in soil is in between 50 ppm and 400 ppm (US EPA, August 3, 2012). In our current societies, the environment has been polluted with lead from the sophisticated industry such as in battery and paint manufacturing. Moreover activities such as mining, smelting and refining also resulted in the increase of substantial amount of lead level in the environment (Maja, 1996 and Niisoe *et al.*, 2011).

Due to the concern and awareness of the hazardous of lead in the environment to human health, some efforts have been taken aiming to reduce the level of human exposure to lead, such as the banned of the use leaded gasoline, closing the lead refinement and lead mining plants (US EPA, August 3, 2012, Lemos and Carvalho, 2010, and Paoliello and Capitani, 2007). Basically, human exposure to lead can happen through two major pathways, the gastrointestinal tract (GIT) and the respiratory tract (Niisoe *et al.*, 2011). The GIT intake depends on the

contamination level of food and water while the respiratory tract depends on the air concentration of contaminant (Niisoe *et al.*, 2011).

The human exposure to lead could be studied through the lead level in human body. Lead level in human body is measured by using spectrometric methods to detect the presence of lead in human body parts. The experimental data obtained from a previous study could be used to study the lead level in human body through a mathematical model. The mathematical model which is derived from a compartmental model based on previous studies reflect the real physical system of lead transfer in human body. The mathematical model can be solved using mathematical technique. This dissertation compares and analyzes the model with different perspectives.

1.2 Statement of the Problem

As mentioned earlier in the previous section, lead level in human body is studied by using a mathematical model. The mathematical model is derived from the Batschelet's three-compartmental model adopted from the Rabinowitz *et al.* (1976)'s compartmental model which reflect the kinetics of lead in human body. The parameters are estimated from the field data drawn from the study made by Rabinowitz *et al.* (1973). However, most of the solutions of the studies on lead level in human body made by previous researchers such as Batschelet *et al.* (1979) and Steyn *et al.* (2008) used numerical solutions approach to solve the mathematical model. Thus, this dissertation used an analytical approach to solve the mathematical model of lead transfer in human body and to show that results produced by analytical approach correspond almost perfectly with those of a numerical approach.

1.3 Objectives of the Study

The objectives of this study are:

- (i) to conceptualize compartmental model for lead transfer in the human body
- (ii) to formulate mathematical model based from the compartmental model for lead in the human body
- (iii) to solve the mathematical model analytically
- (iv) to discuss the result obtained by comparing with published researches

1.4 Scope of the Study

The compartmental system for lead in the human body will be interpreted into mathematical model involving differential equations in which each equation represents the lead concentration in several major parts of human body. The mathematical model was solved theoretically using the data drawn from Rabinowitz *et al.* (1973/76) studies.

1.5 Significance of the Study

The significance of the study can be seen in two different perspectives, theoretically and practically. Theoretically, the study consists of multidisciplines, the mathematical and biological disciplines at which the real biological system is interpreted in mathematical model and formula. Therefore the study is deemed to

REFERENCES

- Barreto, J. B. (1948). Factors Influencing Health And Wellbeing of Workers In The Printing Industry. *Mem. Inst. Osw. Cruz.* 46, 355-413.
- Bastos, R., Pimenta de Mello, R., and Cruz, W. O. (1949). Urinary Excretion of Coproporphyrin In Workers Handling Lead. *Mem. Inst. Osw. Cruz.* 47, 231-240.
- Batschelet, E., Brand, L., and Steiner, A. (1979). On Kinetics of Lead In The Human Body. *Journal of Mathematical Biology.* 8, 15-23.
- Bedrikow. B. (1977). Intoxicação Pelo Chumbo: Aspectos de Ocorrência e Diagnóstico Clínico. *Ver. Brás. Saude Ocup.* 5, 44-46.
- Borges, D. R. (1950). Prevenção da Intoxicação Pelo Chumbo Através da Contagem de Reticulócitos. *Rev. Paul. Méd.* 36, 39-44.
- Blumer, W. and Reich, T. (1975). Health Injuries By Leaded Gasoline. *Schweiz. Rundschau Med. Praxis.* 64, 261-265.
- Cobelli, C., Lepschy, A., Romanin-Jacur, G., and Viaro, U. (1986). An Expanded Schematic For Compartmental Systems. *Math. Biosci.* 79, 97-106.
- Dainty, J. R. (2001). Use of Stable Isotopes And Mathematical Modelling To Investigate Human Mineral Metabolism. *Nutrition Research Reviews.* 14, 295-315.
- Fonseca, E. A. (1953). Problemas de Hygiene e Segurança do Trabalho Na Indústria de Accumuladores Elétricos (Problems of Industrial Hygiene And Safety In The Acid Lead Battery Workplaces). *Arq. Fac. Hig. São Paulo.* 7, 87-100.
- Karhausen, L. (1973). Intestinal Lead Absorbtion. *In: Proceedings of The International Symposium On Environmental Health Aspects of Lead.* 1973. Luxemburg: Commission of The European Communities. 427-440.
- Kellogg, R. B. and Aronsson, G. (1978). On A Differential Equation Arising From Compartmental Analysis. *Mathematical Biosciences.* 38, 113-122.
- Lemos, V. A., and Carvalho, A. L. (2009). Determination of Cadmium And Lead In Human Biological Samples By Spectrometric Techniques: A Review. *Environ Monit Assess.* 171, 255-265.
- Lopez-Fidalgo, J., Rodriguez-Diaz, J. M., Sanchez, G., and Santos-Martin, M. T. (2005). Optimal Design For Compartmental Models With Correlated

- Observations. *Journal of Applied Statistics*. 32(10), 1075-1088.
- Maja, B. (1996). Assessment of Total Human Exposure To Lead And Cadmium. *Environmental Management And Health*. 7(4), 23-28.
- Marcus, A. H. (1983). Compartmental Models With Spatial Diffusion: Estimation For Bone-Seeking Tracers. *Mathematical Biosciences*. 64, 233-248.
- Mohler, R. R. (1974). Biological Modeling With Variable Compartmental Structure. *IEEE Transactions On Automatic Control*. AC-19(6), 922-926.
- Moore, M. R., Meredith, P. A., Watson, W. S., Sumner, D. J., Taylor, M. K., and Goldberg, A. (1980). The Percutaneous Absorption of Lead-203 In Humans From Cosmetic Preparations Containing Lead Acetate, as Assessed by Whole-Body Counting and Other Techniques. *Fd Cosmet Toxicol*. 18, 399-405.
- Nagle, R. K., Saff, E. B. and Snider, A. D. (2012). *Fundamentals of Differential Equations*. (8th ed.). Boston: Pearson Education, Inc.
- Niisoe, T., Harada, K. H., Hitomi, T., Watanabe, T., Hung, N. N., Ishikawa, H., Wang, Z., and Koizumi, A. (2011). Environmental Ecological Modeling of Human Blood Lead Levels In East Asia. *Environmental Science & Technology*. 45, 2856-2862.
- Oliveira , O. G. and Cavalcanti, T. (1950). O Problema do Chumbo Numa Indústria Gráfica [Lead Problem In A Printing Industry]. *Mem. Inst. Osw. Cruz*. 48, 125-142.
- Patterson, C. C. (1965). Contaminated And Natural Lead Environments of Man. *Arch. of Environmental Health*. 11, 344-360.
- Paoliello, M. M. B., and Capitani, E. M. (2007). Occupational And Environmental Human Lead Exposure In Brazil. *Environmental Research*. 103, 288-297.
- Popovic, L., McKinley, S. A., and Reed, M. C. (2011). A Stochastic Compartmental Model For Fast Axonal Transport. *Siam J. Appl. Math*. 71(4), 1531-1556.
- Rabinowitz, M. B., Wetherill, G. W., and Kopple, J. D. (1973). Lead Metabolism In The Normal Man : Stable Isotopes Studies. *Science (Wash. D.C.)*. 182 : 725 – 727.
- Rabinowitz, M. B., Wetherill, G. W., and Kopple, J. D. (1976). Kinetic Analysis of Lead In Healthy Humans. *The Journal of Clinical Investigation*. 58, 260-270.
- Remsing, C. C. (2006). Stability. Retrieved on May 2, 2015, from <https://www.ru.ac.za/media/rhodesuniversity/content/mathematics/documents/thirdyear/linearcontrol/AM32LC4%20Stability.pdf>

- Radke-Sharpe, N., and White, Jr., K. P. (1998). The Role of Qualitative Knowledge In The Formulation of Compartmental Models. *IEEE Transactions On Systems, Man, and Cybernetics-Part C: Applications and reviews*. 28(2), 272-275.
- Salgado, P. E. T. (1976). Diagnóstica Laboratorial da Intoxicação Pelo Chumbo (Laboratory Diagnosis of Lead Intoxication). *Rev. Bras. Saúde Ocup.* 5, 21-34.
- Spínola, A. G., Fernícola, N. A. G. G., and Mendes, R. (1980). *Intoxicação Profissional Por Chumbo*. In Mendes, R. (Ed.) *Medicina do Trabalho-Doenças Profissionais* (pp. 437-460). São Paulo: Sarvier.
- Steyn, F. E., Joubert, S. V., and Coetzee, C. E. (2008). A Discreet Compartmental Model For Lead Metabolism In The Human Body. *Buffelspoort TIME2008 Peer-Reviewed Conference Proceedings*. 22-26 September. Buffelspoort, South Africa. 174-186.
- Teisinger, J., Xintaras, C., Pfitzer, E. (1973). Prague International Lead Panel. Effects of Atmospheric Lead On Biological Systems. *Science*. 179, 197-198.
- United State Environmental Protection Agency (EPA). Retrieved on August 3, 2012 from <http://www.epa.gov/lead/pubs/learn-about-lead.html>.
- Walter, G. G. (1984). Eigenvalues And Structure of Compartmental Models. *Mathematical Biosciences*. 71, 181-199.
- Whitfield, J. B., Dy, V., McQuilty, R., Zhu, G., Heath, A. C., Montgomery, G. W., and Martin, N. (2010). Genetics Effects on Toxic and Essentials Elements in Human: Arsenic, Cadmium, Cooper, Lead, Mercury, Selenium, and Zinc in Erythrocytes. *Environmental Health Perspectives*. 118(6), 776-782
- Zill, D. G. (2005). *A First Course In Differential Equations With Modeling Applications*. (8th ed.). USA: Brooks/Cole.