

MOLECULAR ENGINEERING FOR CATALYTIC EFFICIENCY OF  
XYLANASE FROM *Aspergillus fumigatus* RT-1 AND ITS APPLICATION IN  
HYDROLYSIS OF PRETREATED KENAF

SITI INTAN ROSDIANAH BINTI DAMIS

UNIVERSITI TEKNOLOGI MALAYSIA

MOLECULAR ENGINEERING FOR CATALYTIC EFFICIENCY OF  
XYLANASE FROM *Aspergillus fumigatus* RT-1 AND ITS APPLICATION IN  
HYDROLYSIS OF PRETREATED KENAF

SITI INTAN ROSDIANAH BINTI DAMIS

A thesis submitted in fulfilment of the  
requirements for the award of the degree of  
Doctor of Philosophy (Bioprocess Engineering)

School of Chemical and Energy Engineering  
Faculty of Engineering  
Universiti Teknologi Malaysia

MARCH 2020

## ACKNOWLEDGEMENT

Foremost, I would like to express my sincere gratitude to my supervisor Prof. Dr. Rosli B. Md. Illias for the continuous support of my Ph.D. study and research, for his patience, motivation, enthusiasm, and immense knowledge. His guidance helped me in all the time of research and writing of this thesis. I could not have imagined having a better mentor for my Ph.D. study.

My sincere thanks also go to Ministry of Science, Technology and Innovation (MOSTI) for providing a scholarship of MyBrain which support my fee expenses and allowances throughout five years of my study.

Besides, I would like to thank the rest of the committees from Malaysian Genome Institute which had provided courses and guides for technical knowledge regarding my studies.

I thank my fellow labmates in Genetic Lab from 2011 to 2019: Rabiatul Adawiyah, Izawati, Rohaida, Shalyda, Abbas, Izyan, Hasma, Baidura, Ling How Lie, Faizah, Samson, Hakimi, Yeng, Amalina, Joyce, Joanne, Aishah, Kheng Onn, Aizi Mazila, Anuar, Siti Fatimah, Namirah, Noradilin, Hidayah, Hamiza, Nardiah and Nashriq, for the stimulating discussions, for the sleepless nights we were working together before deadlines, and for all the fun we have had in the last eight years.

Last but not the least, I would like to thank my family: my parents Rusnah Ibrahim and Damis Madek; my siblings: Nina, Mamat, Diana, Huda, Dinah, and Syifaa; and my dearest: Fairos, Khaleef, and Najla for endless loves and cares and supporting me spiritually and financially throughout my life.

## ABSTRACT

The lignocellulose of industrial crops consists of three main polymers: cellulose, hemicellulose, and lignin. The combination of these complex and heterogeneous polymers contributes to the recalcitrant structure of lignocellulose. Thus, it becomes a drawback for a group of hydrolytic enzymes which work synergistically to hydrolyse the lignocellulosic substrate including xylanase. Hence, this study aimed to improve the catalytic efficiency of *Aspergillus fumigatus* RT-1 xylanase (AfxynG1) on pretreated kenaf hydrolysis through protein engineering of amino acids that located near the substrate-binding site and at the N-terminal region. Molecular docking analysis revealed 5 subsites; -3, -2, -1, +1, and +2 and several of substrate-binding residues which distributed alongside the subsites. Two putative binding residues of Phe 146 and Phe 30 and a putative secondary binding site of residue Tyr 7 were determined. High-throughput and low-throughput screenings of 5000 clones from error-prone PCR library which acted as fine tuner and 414 clones from site-saturation mutagenesis library were successfully performed to screen out three improved mutants; c168t, Q192H, and Y7L. The site-directed mutagenesis was applied to construct double and triple mutants and this process resulted in only two improved mutants; c168t/Q192H and c168t/Q192H/Y7L. The triple mutant c168t/Q192H/Y7L was the most stable enzyme in high temperature 60 and 70 °C and acidic pH 3-6, while the double mutant c168t/Q192H showed to contribute to the most effective hydrolysis reaction with a 7.6-fold increase in catalytic efficiency. Mutant Y7L produced the highest sugar yield with 28 % increase in pretreated kenaf hydrolysis. Overall, these improved mutants are feasible to be used synergistically with cellulases for bioconversion of lignocellulose into reducing sugar.

## ABSTRAK

Lignoselulosa yang diperolehi daripada tanaman industri terdiri daripada tiga jenis polimer utama: selulosa, hemiselulosa dan lignin. Gabungan polimer-polimer yang kompleks dan heterogen ini menyumbang kepada struktur rekalsitran lignoselulosa. Maka, ia menjadi masalah bagi sekumpulan enzim hidrolitik yang bertindak secara sinergistik untuk menghidrolisis substrat lignoselulosa termasuk xilanase. Oleh itu, kajian ini dijalankan adalah bertujuan untuk meningkatkan kecekapan pemangkinan xilanase daripada *Aspergillus fumigatus* RT-1 (AfxynG1) ke atas hidrolisis kenaf terawat melalui kejuruteraan protein terhadap asid amino yang terletak berhampiran tapak pengikat substrat dan di kawasan N-terminal. Analisis dok molekul mendedahkan 5 tapak pengikat substrat; -3, -2, -1, +1, dan +2 dan beberapa residu pengikat substrat yang teragih sepanjang tapak pengikat substrat. Dua pengikat residu putatif iaitu Phe 146 dan Phe 30 serta satu residu tapak pengikat sekunder putatif Tyr 7 telah dikenalpasti. Penyaringan pemprosesan tinggi dan rendah terhadap 5000 klon daripada pustaka cenderung ralat PCR yang berperanan sebagai penala halus dan 414 klon daripada pustaka mutagenesis tapak penepuan berjaya menyaring tiga mutan yang ditambahbaik; c168t, Q192H, dan Y7L. Mutagenesis tapak terarah telah digunakan untuk membina mutan ganda dua dan ganda tiga yang mana proses ini menghasilkan hanya dua mutan ditambahbaik; c168t/Q192H dan c168t/Q192H/Y7L. Mutan ganda tiga c168t/Q192H/Y7L ialah enzim yang paling stabil pada keadaan suhu tinggi 60 dan 70 °C dan pH berasid 3-6 manakala mutan ganda dua c168t/Q192H telah menyumbang kepada reaksi hidrolisis paling efektif dengan 7.6 kali ganda peningkatan dalam kecekapan pemangkinan. Mutan Y7L menghasilkan gula tertinggi iaitu sebanyak 28 % peningkatan dalam hidrolisis kenaf terawat. Secara keseluruhannya, semua mutan yang ditambahbaik ini boleh digunakan secara sinergi dengan selulase untuk biopenukaran lignoselulosa kepada gula penurun.

## TABLE OF CONTENTS

	<b>TITLE</b>	<b>PAGE</b>
	<b>DECLARATION</b>	<b>iii</b>
	<b>DEDICATION</b>	<b>iv</b>
	<b>ACKNOWLEDGEMENT</b>	<b>v</b>
	<b>ABSTRACT</b>	<b>vi</b>
	<b>ABSTRAK</b>	<b>vii</b>
	<b>TABLE OF CONTENTS</b>	<b>viii</b>
	<b>LIST OF TABLES</b>	<b>xii</b>
	<b>LIST OF FIGURES</b>	<b>xiv</b>
	<b>LIST OF ABBREVIATIONS</b>	<b>xx</b>
	<b>LIST OF SYMBOLS</b>	<b>xxiii</b>
<b>CHAPTER 1</b>	<b>INTRODUCTION</b>	<b>1</b>
1.1	Background of Study	1
1.2	Problem Statement and Gap of the Study	4
1.3	Objectives of the Study	5
1.4	Scopes of the Study	6
<b>CHAPTER 2</b>	<b>LITERATURE REVIEW</b>	<b>7</b>
2.1	Industrial Biotechnology	7
2.1.1	Biorefinery	8
2.2	Industrial Crops	11
2.2.1	Kenaf as an Industrial Crops	12
2.3	Recalcitrance of Lignocellulosic Biomass	15
2.3.1	Xylan	17
2.4	Enzymatic Hydrolysis of Xylan	18
2.5	GH11 Xylanases: Structure-Function Relationship	19

2.5.1	Catalytic Mechanism	20
2.5.2	Catalytic Domain	21
2.5.3	Secondary-Binding Site	23
2.6	Biochemical Characterisation of GH11 Xylanase	25
2.6.1	Thermostability	25
2.6.2	pH Dependence	39
2.7	Locally Isolated of <i>Aspergillus fumigatus</i> RT-1 Xylanase (AfxynG1)	31
2.8	Xylanase Modification for Industrial Use	31
2.9	Protein Engineering Approaches	33
2.9.1	Error-prone PCR (epPCR)	35
2.9.2	Site-Saturation mutagenesis (SSM)	35
2.9.3	Site-Directed Mutagenesis (SDM)	37
<b>CHAPTER 3</b>	<b>MATERIALS AND METHODS</b>	<b>39</b>
3.1	Work Outline	39
3.2	Bacterial Strains and Plasmids	41
3.3	Chemicals and Biological Enzymes	41
3.4	Pretreatment of Kenaf	42
3.5	Library Screening Towards Pretreated Kenaf	42
3.5.1	High-Throughput Screening (HTS)	43
3.5.2	Low-Throughput Screening (LTS)	44
3.6	Homology Modeling and Molecular Docking of AfxynG1	45
3.6.1	Protein Sequence Analysis	45
3.6.2	Template Search and Selection	45
3.6.3	AfxynG1 3D Structure Modeling	46
3.6.4	Structure Refinement and Validation	46
3.6.5	Molecular Docking	47
3.6.6	Protein Interaction Analysis	47
3.7	Site Saturation and Site-Directed Mutagenesis	48
3.8	Insert Screening	49
3.9	SSM Library Construction	50
3.10	General DNA Manipulation Techniques	50

3.10.1	Agarose Gel DNA Electrophoresis	50
3.10.2	Plasmid DNA Isolation	51
3.10.3	Gel Extraction of PCR Amplified Gene Fragments	52
3.10.4	<i>E. coli</i> Competent Cells Preparation	52
3.10.5	Cell Transformation	53
3.10.6	DNA Quantification	53
3.11	Enzyme Production	54
3.12	Enzyme Purification by Gravity Flow Chromatography	54
3.13	General protein techniques	55
3.13.1	Sodium Dodecyl Sulphate-Polyacrylamide Gel Electrophoresis	55
3.13.2	Western Blot	56
3.13.3	Protein Concentration Assay	57
3.13.4	Xylanase Assay	58
3.14	Xylanase Characterisation	58
3.14.1	Optimum Temperature and pH	59
3.14.2	Thermal and pH Stability	59
3.14.3	Michaelis-Menten Kinetics	60
3.16	Optimisation Condition of Hydrolysis Activity Assay	60

<b>CHAPTER 4</b>	<b>SCREENING AND SELECTION OF MUTANTS FROM EPPCR AND SSM LIBRARIES AND DETERMINATION OF SUBSTRATE-BINDING RESIDUES</b>	<b>61</b>
4.1	Introduction	61
4.2	Substrate Preparation	62
4.3	Optimisation of High-Throughput Screening	63
4.3.1	Reaction Volume	64
4.3.2	Enzyme-Buffer Ratio	65
4.3.3	Incubation Time and Agitation	68
4.4	EpPCR Library Screening	73
4.5	Protein Sequence Analysis	75



4.6	Secondary Structure Analysis	79
4.7	Homology Modeling	81
	4.7.1 Template Searching	81
	4.7.2 Protein Sequence Alignment of AfxynG1 and 1TE1_B	83
	4.7.3 Model Building	84
	4.7.4 Model Evaluation	85
4.8	Energy Minimisation of AfxynG1	89
4.9	Molecular Docking	92
4.10	Substrate-Binding Site	92
	4.10.1 Thumb Region	99
	4.10.2 Novel Substrate-Binding Residues	100
4.11	Site-Saturation Mutagenesis (SSM)	102
	4.11.1 SSM Library	104
	4.11.2 Screening of SSM Mutants	105
4.12	Conclusion	106
<b>CHAPTER 5</b>	<b>BIOCHEMICAL CHARACTERISATION AND HYDROLYSIS PERFORMANCE OF THE MUTANTS</b>	<b>109</b>
5.1	Introduction	109
5.2	Construction of Multiple Mutants and Identification of the Final Mutants	109
5.3	Effect of Temperature on Enzyme Activity	111
5.4	Effect of pH on Enzyme Activity	115
5.5	Mutations Affect the Kinetic Parameters	118
5.6	Reaction Optimisation of Pretreated Kenaf Hydrolysis	123
5.7	Hydrolysis of Pretreated Kenaf	126
5.8	Conclusion	128
<b>CHAPTER 6</b>	<b>CONCLUSION AND FUTURE WORKS</b>	<b>129</b>
6.1	Conclusion	129
6.2	Future Works	130

<b>REFERENCES</b>	<b>132</b>
<b>APPENDICES</b>	<b>158</b>
<b>LIST OF PUBLICATIONS AND PAPER PRESENTED</b>	<b>177</b>

### LIST OF TABLES

<b>TABLE NO.</b>	<b>TITLE</b>	<b>PAGE</b>
Table 3.1	Primers used for site-saturation and site-directed mutagenesis	49
Table 4.1	The volume of enzyme and buffer for ratio enzyme: buffer optimisation	65
Table 4.2	Comparison of three ratios enzyme: buffer according to the ranks	66
Table 4.3	Comparison between the sampling time in different mode of incubation	69
Table 4.4	Comparison of incubator shaker and water bath at different incubation times	71
Table 4.5	Summary of optimum parameters for HTS	73
Table 4.6	Mutants selected from epPCR library screening	74
Table 4.7	ProtParam analysis result of AfxynG1	77
Table 4.8	Blastp analysis shows the highest match of AfxynG1 to the nr database	79
Table 4.9	Homologous protein hit with the highest identity and coverage according to PSI-BLAST, HHpred and Phyre2 servers	82
Table 4.10	Model evaluation of AfxynG1 by online tools	86
Table 4.11	Comparison of the AfxynG1 (non-energy minimised) and AfxynG1_EM (energy minimised) model via online tools	90
Table 4.12	Protein-ligand interaction of AfxynG1 and X2-X6 ligands involving stacking interactions, hydrogen bonds and Van der Waals forces at the applicable subsites.	93
Table 4.13	Distance between mutated residues and catalytic sites and their location.	103
Table 4.14	Summary of SSM libraries screening	106
Table 5.1	Pre-screening of mutants towards pretreated kenaf	

	hydrolysis	110
Table 5.2	Kinetic parameters of the wild-type AfxynG1 and its mutants	119
Table 5.3	Protein interaction analysis of residue 192 in the wild-type and mutant Q192H	121
Table 5.4	Protein interactions of residue 7 in the wild-type and mutant Y7L	122
Table 5.5	Fixed parameters for optimisation of pretreated kenaf hydrolysis	124
Table 5.6	Final condition for pretreated kenaf hydrolysis	125
Table 5.7	Enzyme hydrolysis of pretreated kenaf at three different temperature; 30 °C, 40 °C and 50 °C with 36 h incubation	127

## LIST OF FIGURES

FIGURE NO.	TITLE	PAGE
Figure 1.1	Ribbon representation of <i>Thermobacillus xylanilyticus</i> xylanase (Tx-Xyn) three-dimensional structure. The schematic protein is ‘color-ramped’ from the N-terminus (blue, N-ter) to the C-terminus (red, C-ter). The relevant regions of ‘thumb’, ‘palm’ and ‘fingers’ are highlighted in frames, and the ‘knuckles’ in the fingers region is indicated by an arrow (Song <i>et al.</i> , 2012).	3
Figure 2.1	Basic pathway for ethanol production from lignocellulosic biomass through sugar and syngas platform (Vohra <i>et al.</i> , 2014)	9
Figure 2.2	Four main stages involved in lignocellulosic biorefinery for ethanol production (Asgher <i>et al.</i> , 2014).	11
Figure 2.3	Kenaf plants (Ramesh, 2016).	13
Figure 2.4	Possible bioenergy pathway based on kenaf biomass (Saba <i>et al.</i> , 2015).	14
Figure 2.5	The main components and structure of lignocellulose (Isikgor and Becer, 2015).	16
Figure 2.6	Schematic diagram of xylan hydrolysis by glycoside hydrolases or carbohydrate esterases (Madadi <i>et al.</i> , 2017).	19
Figure 2.7	Retaining mechanism. The reaction occurs via double displacement where one carboxylic group acts as general acid/base, and the other one as nucleophile forming a covalent intermediate (Linares-Pastén, Aronsson and Karlsson, 2016).	21
Figure 2.8	Cartoon representation (left) and topology diagram (right) of a typical GH11 xylanase member ( Paës <i>et al.</i> , 2012a).	22
Figure 2.9	Schematic diagram of <i>Bacillus subtilis</i> (BsXynA) subsites. Subsites where the glycon region (non-reducing	

	end) of the substrate binds (-3 to -1), and subsites where the aglycon region (reducing end) binds (+1 to +3), are indicated. The bond cleavage occurs in between -1 and +1 is represented by the ‘scissor’ (Pollet <i>et al.</i> , 2010).	23
Figure 2.10	Secondary binding sites in xylanase from <i>Bacillus subtilis</i> (PDB 2QZ3) that bind the substrate ligand at the ‘knuckle’ region (Vandermarliere <i>et al.</i> , 2008).	24
Figure 2.11	Geometries of aromatic-aromatic interaction: a) edge to face, b) offset stacked and c) face to face (Waters, 2002)	27
Figure 2.12	Whole plasmid, site-saturation mutagenesis using single- round PCR. The symbol ‘*’ represents point mutation (Labrou, 2010).	36
Figure 3.1	Work outline of the study	40
Figure 3.2	Layer arrangement of the western blot	57
Figure 4.1	Pretreated kenaf using two-stage pretreatment and seized at size 0.1-0.5 mm	63
Figure 4.2	Scatter plot of hydrolysis reaction by varying the enzyme: buffer ratio from three different 96-well plates containing epPCR clones	67
Figure 4.3	Sampling data of enzyme hydrolysis at different incubation times, A) water bath B) incubator shaker	70
Figure 4.4	Scatter plot of enzyme reaction in incubator shaker and water bath at different incubation time; A) 2 h, B) 4 h, C) 6 h and D) 16 h.	72
Figure 4.5	SignalP 5.0 analysis detected an N-terminal signal peptide at the cleavage site between Ala 29 and Thr 30	76
Figure 4.6	Protein sequence of AfxynG1 in FASTA format obtained from the previous study (Abdul Wahab <i>et al.</i> , 2016). Residues in red font are the N-terminal signal peptide that had been removed.	76
Figure 4.7	InterProScan analysis categorised AfxynG1 in GH11 family containing GH11 family domain and two active sites.	78
Figure 4.8	Secondary structure prediction of AfxynG1 by PSIPRED 4.0. Yellow boxes are $\beta$ -strands, pink boxes are $\alpha$ -helices, and black lines are coils. The confidence of prediction is showed by blue colour boxes. The confidence is increasing as the colour becomes darker.	80
Figure 4.9	A) Phylogenetic tree analysis of AfxynG1, 1TE1_B and	

	2VUL_A by Clustal Omega. B) Structure of GH11 xylanase from <i>T. funiculosus</i> (1TE1_B) as visualised by Pymol. The structure is similar to a partially closed right hand consisting of palm, fingers and thumb.	82
Figure 4.10	Protein sequence alignment between xylanase from <i>A. fumigatus</i> (Afxyn) and xylanase from <i>T. funiculosus</i> (1TE1) by Clustal Omega. The extended N-terminal region of Afxyn was underlined. Arrows indicate two active sites of glutamate shared by these two xylanases.	83
Figure 4.11	The best model of AfxynG1 generated by Modeller. B1-B8 are concave antiparallel sheets while A2-A6 are convex antiparallel sheets. The main part of the model consists of fingers, thumb and palm	84
Figure 4.12	Topology structure of AfxynG1 contains 12 $\beta$ -sheets and one $\alpha$ -helix	85
Figure 4.13	Structure alignment between AfxynG1 (blue) and 1TE1_B (orange) by Pymol visualisation. The extended N-terminal region of AfxynG1 that was not covered by the template has been circled.	87
Figure 4.14	Ramachandran plot analysis of AfxynG1 model. The most favoured regions showed in red, the additional allowed regions in yellow, the generously allowed regions in cream and the disallowed regions in white. The residues in red located in the generously allowed and disallowed regions.	88
Figure 4.15	Protein sequence alignment between xylanase from <i>A. fumigatus</i> (AfxynG1) and xylanase from <i>N. patriciarum</i> (Np-Xyn) by Clustal Omega. Arrows represent the two anti-parallel $\beta$ -sheets for both protein structures. The underlined sequence indicates a potential new, additional $\beta$ -sheets at N-terminal region of AfxynG1.	89
Figure 4.16	Structure alignment between non-energy minimised AfxynG1 (green) and energy minimised AfxynG1 (red) by Pymol.	90
Figure 4.17	Ramachandran plot analysis of energy minimised AfxynG1 model. The most favoured regions showed in red, the additional allowed regions in yellow, the generously allowed regions in cream and the disallowed regions in white. The residues in red are located in the generously allowed and disallowed regions.	91

- Figure 4.18 Pymol visualisation of protein ligand interaction between AfxynG1 and A) X2, B) X3, and C) X4. Grey cartoon represents overall structure of AfxynG1. Blue sticks are substrate-binding residues and red sticks are pair of catalytic residues. Green and red sticks are ligands that are inserted into the cleft of AfxynG1. Subsites are labelled by  $-n$  (non-reducing ends) and  $+n$  (reducing ends). Hydrogen bonds between atoms are shown by yellow dotted lines. 95
- Figure 4.19 Pymol visualisation of protein ligand interaction between AfxynG1 and A) X5, and B) X6. Grey cartoon represents overall structure of AfxynG1. Blue sticks are substrate-binding residues and red sticks are pair of catalytic residues. Green and red sticks are ligands that are inserted into the cleft of AfxynG1. Subsites are labelled by  $-n$  (non-reducing ends) and  $+n$  (reducing ends). Hydrogen bonds between atoms are shown by yellow dotted lines. 97
- Figure 4.20 Schematic diagram of AfxynG1 subsites for substrate-binding at non-reducing end, NR and product release at reducing end, R. X2-X6 are xylooligosaccharides for molecular docking. E189 and E98 are catalytic residues to initiate the xylan hydrolysis between subsite -1 and +1. 98
- Figure 4.21 Thumb region of AfxynG1 is shown in red sticks. The five residues involved are Pro 138, Ser 139, Ile 140, Gln 141, and Gly 142. 99
- Figure 4.22 Multiple sequence alignment of GH11 xylanases by Clustal Omega. Red fonts represent conserved catalytic residues. Substrate-binding residues with their conserved amino acids from other xylanases are highlighted in black. Residues highlighted in red show the putative substrate-binding residues of Phe 30 and Phe 146. GH11 xylanases are from *Trichoderma reesei* (1XYN and 1ENX; Torronen and Rouvinen, 1995), *Aspergillus kawachii* (1BK1; Fushinobu *et al.*, 2002), *Streptomyces* sp. S38 (1HIX; Wouters *et al.*, 2001), *Aspergillus niger* (1UKR; Krengel and Dijkstra, 1996), *Bacillus agaradhaerens* (1H4G; Sabini *et al.*, 2001), *Bacillus circulans* (1BCX;

Wakarchuk *et al.*, 1994) and *Thermobacillus xylanilyticus* (Tx-xyl; Paës and O'Donohue, 2006). 101

- Figure 4.23 Protein-ligand interactions of putative substrate-binding residues. Green and red sticks are ligands that are inserted into the enzyme cleft. A) Phe 146 stacks onto the aromatic ring of xylose at the -1 subsite. B) Phe 30 forms a hydrogen bond with the xylose at subsite -2. 102
- Figure 4.24 Location of site-saturation mutagenesis residues of AfxynG1. Red sticks indicate catalytic residues and blue sticks show selected residues for site-saturation mutagenesis. 104
- Figure 5.1 Effects of temperature on the activity and stability of AfxynG1 and its mutants. A) Effect of temperature on the activity of AfxynG1 and its mutants. The assay was performed at different temperature ranging from 30 to 70 °C for 10 min. B) Effect of temperature on the Stability of AfxynG1 and its mutants. The purified enzymes were incubated in 50 mM Na-acetate buffer (pH 5.0) without substrate for 30 min at varied temperature (30-70 °C) and the relative activities were measured at the optimal condition for 10 min. The highest activity of each WT and mutants were set as 100 %. 113
- Figure 5.2 Different conformation of residue 192 within the wild-type and mutant Q192H. A) Side chain of Gln 192 is facing away from Asp 33 in the wild-type. B) Side chain of His 192 is facing towards Asp 33. Location of Gln 192 (red stick) in the tip of fingers is shown in the small box at the top right. 115
- Figure 5.3 Effects of pH on the activity and stability of AfxynG1 and its mutants. A) Effect of pH on the activity of AfxynG1 and its mutants. The assay was performed in different pH buffer ranging from pH 3 to 8 at 50 °C for 10 min. B) Effect of pH on the stability of AfxynG1 and mutants. The enzyme was incubated in different buffer (pH 3 to 6) at 25 °C for 30 min and the relative activities were measured at the optimal condition for 10 min. The highest activity of each WT and mutants were set as 100 %. 117
- Figure 5.4 A) Gln 192 interaction with close residues at the fingers region. Grey cartoon is xylanase structure. Green sticks represent residues involved. Black dotted line is the



distance measurement between residues. B) His 192 interaction with close residues. Cyan stick shows xylohexaose being inserted into the active site cleft. Red dotted line shows protein-ligand interaction. The small box at the top right shows the location of residue 192 at the tip of the fingers of xylanase

120

Figure 5.5 Reaction optimisation of pretreated kenaf hydrolysis involving several parameters studied: A) substrate loadings B) pH and C) enzyme loadings. At each experiment, the fixed conditions are applied except the parameters studied.

124

## LIST OF ABBREVIATIONS

GH	-	Glycoside hydrolase
N-ter	-	N-terminus
C-ter	-	C-terminus
PCR	-	polymerase chain reaction
epPCR	-	error-prone PCR
GHG	-	greenhouse gas
AFLP	-	amplified fragment length polymorphism
MARDI	-	Malaysia Agricultural Research and Development Institute
EC	-	Enzyme Commission
CBM	-	carbohydrate-binding modules
SBS	-	secondary binding site
Asn	-	asparagine
Thr	-	threonine
Ser	-	serine
Trp	-	tryptophan
Arg	-	arginine
CMC	-	carboxymethyl cellulose
StEP	-	staggered extension process
mM	-	millimolar
dNTP	-	deoxynucleoside triphosphate
Mg <sup>2+</sup>	-	ion magnesium
Mn <sup>2+</sup>	-	ion mangan
DNA	-	deoxyribonucleic acid
Kb	-	kilobase

SSM	-	site-saturation mutagenesis
DMSO	-	dimethyl sulfoxide
DNS	-	3,5-dinitrosalicylic acid
PPA	-	polyphosphoric acid
ml	-	mililiter
h	-	hour
HTS	-	high-throughput screening
LTS	-	low-throughput screening
LB	-	Luria-Bertani
IPTG	-	Isopropyl $\beta$ - d-1-thiogalactopyranoside
rpm	-	revolutions per minute
$\mu$ l	-	microliter
min	-	minutes
nm	-	nanometer
OD	-	optical density
mg/ml	-	milligram per mililiter
g	-	gram
mg	-	milligram
BLAST	-	Basic Local Alignment Search Tool
RMSD	-	root mean square deviation
PDB	-	Protein Data Bank
3D	-	three-dimensional
PIC	-	Protein Interaction Calculator
ng	-	nanogram
U	-	enzyme unit
sec	-	seconds
$\mu$ g	-	microgram
TAE	-	tris-acetate-EDTA
TSS	-	transformation and storage solution
NiSO <sub>4</sub>	-	nickel sulphate
SDS-PAGE	-	sodium dodecyl sulphate-polyacrylamide gel electrophoresis
V	-	voltan
w/v	-	weight per volume
RT	-	room temperature

BSA	-	bovine serum albumin
UV	-	ultra-violet
$K_m$	-	Michaelis-Menten constant
$k_{cat}$	-	turnover number
$k_{cat}/K_m$	-	specificity constant
OFAT	-	one factor at a time
mg/g	-	milligram per gram
Ala	-	alanine
GRAVY	-	grand average of hydrophobicity
Lys	-	lysine
MW	-	molecular weight
DOPE	-	discrete optimized protein energy
Leu	-	leucine
Gln	-	glutamine
Tyr	-	tyrosine
Glu	-	glutamate
Sec.	-	section
X2	-	xylobiose
X3	-	xylotriose
X4	-	xylotetraose
X5	-	xylopentaose
X6	-	xylohexaose
kcal/mol	-	kilo calorie per mole
Phe	-	phenylalanine
Val	-	valine
Pro	-	proline
Ile	-	isoleucine
Asp	-	aspartate
His	-	histidine
Gly	-	glycine
SDM	-	site-directed mutagenesis
wild-type	-	wild-type
Ni-NTA	-	nickel-NTA
Cys	-	cysteine

R <sub>s</sub> max	-	maximum reducing sugar
MD	-	molecular dynamics
NMR	-	nuclear magnetic resonance
HPLC	-	high-performance liquid chromatography
TLC	-	thin-layer chromatography

### LIST OF SYMBOLS

%	-	percent
°C	-	degree celcius
β	-	beta
α	-	alpha
Å	-	angstrom
°	-	degree
≥	-	more than or equal
=	-	equal
>	-	more than

# CHAPTER 1

## INTRODUCTION

### 1.1 Background of Study

According to the “green chemistry” evolution concept, efficient utilization of raw materials is applied on to minimize waste by avoiding the use of toxic and/or hazardous substances that can lead to health, safety and environmental issues (Sheldon, 2014). Therefore, to exploit these raw materials in a sustainable approach by an economical and friendly manner, then biorefinery comes into the picture (Arevalo-Gallegos *et al.*, 2017). A lignocellulose biorefinery plan manages to produce various value-added biochemical products like ethanol, ethylene, sorbitol, xylitol, furfural, etc (Iqbal, Kyazze and Keshavarz, 2013). Examples of lignocellulosic biomasses are industrial crops which have dominated 89.5% of agricultural land in Malaysia (Arshad, 2017). Besides palm oil, rubber, and cocoa, kenaf has emerged as one of the potential industrial crops since 2000 for production of fibres, polymer composite, and paper (Cheng, Haque Akanda and Nyam, 2016; Alkbir *et al.*, 2016; Ashori, 2006).

Lignocellulose is composed of three main polymers which are cellulose (40-60 % of the total dry weight), hemicellulose (20-40 %) and lignin (10-25 %). These polymers are the root cause of the lignocellulose recalcitrance according to the high crystallinity and polymerization degree of cellulose (Lee, Hamid and Zain, 2014), heterogeneity of hemicellulose (Gírio *et al.*, 2010) and highly aromatic contained lignin (Schoenherr, Ebrahimi and Czermak, 2018). Thus, the degradation of lignocellulosic biomass is very complex and requires synergistic action of

hemicellulases, cellulases and ligninolytic enzymes (Andlar *et al.*, 2018). The second most abundant polymer of hemicellulose has a random and amorphous structure comprises xylan, galactomannan, glucuronoxylan, arabinoxylan, glucomannan and xyloglucan (Isikgor and Becer, 2015; Baeyens *et al.*, 2015). Thus, complete degradation of hemicellulosic polysaccharides requires the cooperation of multiple hemicellulases including endo-xylanases,  $\beta$ -xylosidases,  $\alpha$ -L-arabinofuranosidases,  $\alpha$ -D-glucuronidases, and acetyl-xylan-esterases to break down xylan into linear xylooligosaccharides (Shallom and Shoham, 2003; Beg *et al.*, 2001; Madadi, Tu and Abbas, 2017).

Xylanase as one of the hemicellulases are grouped into glycoside hydrolase (GH) 10 and 11 based on amino acid sequence homologies and hydrophobic cluster analysis of CAZy database (Henrissat, 1991). GH11 xylanase has a lower molecular weight compared to the GH 10 family and it folds into a domain composed of two  $\beta$ -sheets (A and B) packed parallel to each other and one  $\alpha$ -helix. The protein structure is similar to a partially closed right hand consisted of ‘thumb’, ‘palm’ and ‘fingers’ regions (Havukainen *et al.*, 1996; Törrönen and Rouvinen, 1997) (Figure 1.1). These regions are involved in substrate binding and catalysis and even in certain xylanase, they get assisted by the secondary binding site located in N-terminal region (Ludwiczek *et al.*, 2007). Due to the ability to hydrolyse the  $\beta$ -D-(1,4) xylosidic linkages in xylan, GH 11 xylanases have a great commercial interest in potential industrial applications of biorefinery. However, the nature of “biomass recalcitrance” of the lignocellulose is challenging for efficient hydrolysis by xylanase that leads to the increase of the enzyme costs but low yields sugar produced (Himmel *et al.*, 2007; Visser *et al.*, 2015).




Figure 1.1 Ribbon representation of *Thermobacillus xylanilyticus* xylanase (Tx-Xyn) three-dimensional structure. The schematic protein is ‘color-ramped’ from the N-terminus (blue, N-ter) to the C-terminus (red, C-ter). The relevant regions of ‘thumb’, ‘palm’ and ‘fingers’ are highlighted in frames, and the ‘knuckles’ in the fingers region is indicated by an arrow (Song *et al.*, 2012).

To encounter the problem of inefficiency and low yields in enzymatic hydrolysis of lignocellulosic biomass, the catalytic efficiency of xylanase is needed to be increased first before performing cooperation action with cellulases (Morone and Pandey, 2014; Berlin *et al.*, 2005). As a solution, protein engineering can be applied on using one of the approaches that are rational design which is more suitable for the improvement of activity, stability or substrate specificity of the target enzyme. Plus, it encounters the problem of large clone libraries needed by random mutagenesis which causes a troublesome for high-throughput screening (Chica, Doucet and Pelletier, 2005; Martinez and Schwaneberg, 2013). However, researchers discovered that the most successful strategy is the combination of random and focused mutagenesis (Packer and Liu, 2015). For instances, the combination of N-terminal region replacement and site-directed mutagenesis at the cord of xylanase has significantly improved the specific activity (5.3-fold increase), substrate affinity and catalytic efficiency (Li *et al.*, 2017). Another study by Hoffmam *et al.*, 2016 showed that the



fusion of a carbohydrate-binding module from GH 6 resulted in 65 % increase of catalytic efficiency of GH11 xylanase and led to the 17 % increase of sugar release from pretreated sugarcane bagasse hydrolysis. Plus, error-prone PCR mutagenesis in combination with site saturation mutagenesis at H179 residue improved  $k_{cat}/K_M$  of xylanase to 3.46-fold (Wang *et al.*, 2013).

Previously, one of the potential GH11 xylanase isolated from *Aspergillus fumigatus* RT-1, afxynG1 (GenBank accession no: GQ458016) showed a great thermostability by retaining 70% of its activity after 30 minutes incubation at 70°C compared to its optimum temperature at 50°C (Abdul Wahab, Jonet and Illias, 2016). However, the sugar produced from the hydrolysis of the lignocellulosic substrate is very low. Thus, it is necessary to improve the catalytic efficiency of AfxynG1 for better hydrolytic performance. The main focus of this study is to improve the catalytic efficiency of AfxynG1 using protein engineering for producing a better yield of sugar. Error-prone PCR, site-saturation and site-directed mutagenesis approaches led to the amino acid substitutions of AfxynG1 within the N-terminal region and substrate binding site of the AfxynG1. The result showed that the mutations acquired increased catalytic efficiency ( $k_{cat}/K_M$ ) and improved thermostability and acid stability. Therefore, the production of total reducing sugar was enhanced up to 28.6 % from pretreated kenaf hydrolysis. In consequences, these mutants have a great potential in lignocellulosic biomass saccharification for large industrial applications.

## **1.2 Problem Statement and Gap of the Study**

The recalcitrance of lignocellulose structure becomes a drawback for enzymatic hydrolysis due to the complex and heterogeneous structure of cellulose, hemicellulose and, lignin. A previous study used two-step pretreatment process involves calcium hydroxide (Ca(OH)<sub>2</sub>), and peracetic acid (PPA) had successfully removing a major part of the lignin layer and maintaining most of the hemicellulose

of kenaf (Wan Azelee *et al.*, 2014). This hemicellulose part is ready to be degraded by hydrolytic enzymes for fermentable sugar production.

Multiple hydrolysis enzymes are required to function synergistically in saccharification of the lignocellulosic biomass for biorefinery industry. One of the main enzymes involved is xylanase which is still lack of study compared to the cellulases. The most studied xylanases concerning lignocellulosic degradation are involved in a mixture or synergism with other hydrolytic enzymes (Jia *et al.*, 2015; Yang *et al.*, 2015). Besides, the genetic modification for xylanase improvements are more focus on the high temperature and extreme pH tolerance (Li *et al.*, 2015; Boonyapakron *et al.*, 2017) but the catalytic enhancement is still very little. In the meantime, only one study was focus to improve the catalytic efficiency of xylanase towards natural substrate hydrolysis (Song *et al.*, 2012).

Thus, the main target of this study is to improve the catalytic efficiency of the xylanase to produce a higher yield of sugar from the lignocellulosic substrate hydrolysis. The knowledge regarding substrate binding and catalysis of the xylanase must be explored to enable the genetic modification of the suitable amino acid residues. Furthermore, the enhancement of thermostability and pH stability is required to prepare a high potential enzyme for industrial use.

### **1.3 Objectives of the Study**

The objectives of this research are stated as below;

- 1) To identify substrate binding residues that involve and important for substrate binding and catalysis of AfxynG1

- 2) To improve the catalytic efficiency of AfxynG1 for hydrolysis of pretreated kenaf and biochemically characterise the AfxynG1 mutants

#### **1.4 Scopes of the Study**

This study focuses on the improvement of catalytic efficiency of a GH11 xylanase from *A. fumigatus* RT1 (AfxynG1) using directed evolution of epPCR, site saturation and site-directed mutagenesis. The epPCR acts as a fine tuner to select potential residues for SSM following two strategies: 1) residues that shared by more than one mutant and 2) residues which near to catalytic and substrate binding site. The second strategy needs structural information using molecular docking to identify the substrate-binding residues. All of the clones from epPCR and SSM libraries undergo high-throughput and low-throughput screening towards pretreated kenaf hydrolysis to isolate the potential mutants. Multiple mutants from three single improved mutants are constructed employing site-directed mutagenesis and screened towards pretreated kenaf. All of the final mutants are expressed and partially purified for biochemically characterised and kinetic determination. The reducing sugar produced from pretreated kenaf hydrolysis for each mutant was compared.

## REFERENCES

- Abdul Wahab, M.K.H., Jonet, M.A. and Illias, R.M. (2016) 'Thermostability enhancement of xylanase *Aspergillus fumigatus* RT-1', *Journal of Molecular Catalysis B: Enzymatic*, 134, pp.154–163. Available at: <http://dx.doi.org/10.1016/j.molcatb.2016.09.020>.
- Acker, M.G. and Auld, D.S. (2014) 'Considerations for the design and reporting of enzyme assays in high-throughput screening applications', *Perspectives in Science*, 1(1–6), pp.56–73. Available at: <http://dx.doi.org/10.1016/j.pisc.2013.12.001>.
- Ahmad, S., Kamal, M.Z., Sankaranarayanan, R. and Rao, N.M. (2008) 'Thermostable *Bacillus subtilis* Lipases: In Vitro Evolution and Structural Insight', *Journal of Molecular Biology*, 381(2), pp.324–340.
- Akber Basri, M.H., Arifin, A., Junejo, N., Abdul Hamid, H. and Ahmed, K. (2014) 'Journey of kenaf in Malaysia: A Review', *Scientific Research and Essays*, 9(11), pp.458–470. Available at: <http://academicjournals.org/journal/SRE/article-abstract/B60A2D745210>.
- Alam, M. M., Siwar, C., Md Wahid, M., Molla, R. I. and Toriman, M. E. (2010) 'Socioeconomic Profile of Farmer in Malaysia: Study on Integrated Agricultural Development Area in North-West Selangor', *Agricultural Economics and Rural Development New Series*, 7(2), pp.249–265. Available at: [ftp://www.ipe.ro/RePEc/iag/iag\\_pdf/AERD1013\\_249-265.pdf](ftp://www.ipe.ro/RePEc/iag/iag_pdf/AERD1013_249-265.pdf).
- Alexopoulou, E., Li, D., Papatheohari, Y., Siqui, H., Scordia, D. and Testa, G. (2015) 'How kenaf (*Hibiscus cannabinus* L.) can achieve high yields in Europe and China', *Industrial Crops and Products*, 68, pp.131–140.
- Ali, M.H. and Imperiali, B. (2005) 'Protein oligomerization: How and why', *Bioorganic and Medicinal Chemistry*, 13(17), pp.5013–5020.
- Alkbir, M.F.M., Sapuan, S.M., Nuraini, A.A. and Ishak, M.R. (2016) 'Fibre properties and crashworthiness parameters of natural fibre-reinforced composite structure: A literature review', *Composite Structures*, 148, pp.59–73. Available at: <http://dx.doi.org/10.1016/j.compstruct.2016.01.098>.

- Altschul, S.F., Madden, T. L., Schäffer, A. A., Zhang, J., Zhang, Z., Miller, W. and Lipman, D. J. (1997) 'Gapped BLAST and PSI-BLAST: A new generation of protein database search programs', *Nucleic Acids Research*, 25(17), pp.3389–3402.
- Altschul, S. F., Gish, W., Miller, W., Myers, E. W. and Lipman, D J. (1990) 'Basic local alignment search tool', *Journal of molecular biology*, 215(3), pp.403–10. Available at:  
<http://www.sciencedirect.com/science/article/pii/S0022283605803602>.
- Álvarez, C., Reyes-Sosa, F.M. and Díez, B. (2016) 'Enzymatic hydrolysis of biomass from wood', *Microbial Biotechnology*, 9(2), pp.149–156.
- Andlar, M., Rezić, T., Marđetko, N., Kracher, D., Ludwig, R. and Šantek, B. (2018) 'Lignocellulose degradation: An overview of fungi and fungal enzymes involved in lignocellulose degradation', *Engineering in Life Sciences*, 18(11), pp.768–778.
- Andrić, P., Meyer, A.S., Jensen, P.A. and Dam-Johansen, K. (2010) 'Reactor design for minimizing product inhibition during enzymatic lignocellulose hydrolysis: I. Significance and mechanism of cellobiose and glucose inhibition on cellulolytic enzymes', *Biotechnology Advances*, 28(3), pp.308–324.
- Anjana, R., Vaishnavi, M. K., Sherlin, D., Kumar, S. P., Naveen, K., Kanth, P. S. and Sekar, K. (2012) 'Aromatic-aromatic interactions in structures of proteins and protein-DNA complexes: a study based on orientation and distance', *Bioinformatics*, 8(24), pp.1220–1224.
- Anwar, Z., Gulfraz, M. and Irshad, M. (2014) 'Agro-industrial lignocellulosic biomass a key to unlock the future bio-energy: A brief review', *Journal of Radiation Research and Applied Sciences*, 7(2), pp.163–173. Available at:  
<http://linkinghub.elsevier.com/retrieve/pii/S1687850714000119>.
- Arevalo-Gallegos, A., Ahmad, Z., Asgher, M., Parra-Saldivar, R., and Iqbal, Hafiz M.N. (2017) 'Lignocellulose: A sustainable material to produce value-added products with a zero waste approach—A review', *International Journal of Biological Macromolecules*, 99, pp.308–318. Available at:  
<http://dx.doi.org/10.1016/j.ijbiomac.2017.02.097>.
- Arkin, M. (2001) *In vitro Mutagenesis*, in: Brenner, S. (eds) *Encyclopedia of Genetics*. Academic Press, pp. 1010–1014.

- Arnold, F.H. and Moore, J.C. (1997) 'Optimizing industrial enzymes by directed evolution', *Adv Biochem Eng Biotechnol*, 58, pp.1–14.
- Arnold, F.H. (1996) 'Directed Evolution: Creating Biocatalysts for the Future', *Chemical Engineering Science*, 51(23), pp.5091–5102.
- Arshad, F.M. (2017) 'Food Policy in Malaysia', *Reference Module in Food Science*, pp.1-12.
- Asgher, M., Shahid, M., Kamal, S. and Iqbal, H.M.N. (2014) 'Recent trends and valorization of immobilization strategies and ligninolytic enzymes by industrial biotechnology', *Journal of Molecular Catalysis B: Enzymatic*, 101, pp.56–66. Available at: <http://dx.doi.org/10.1016/j.molcatb.2013.12.016>.
- Ashori, A. (2006) 'Pulp and paper from kenaf bast fibers', *Fibers and Polymers*, 7(1), pp.26–29.
- Ayadi, R., Hanana, M., Mzid, R., Hamrouni, L., Khouja, M. I. and Salhi Hanachi, A. (2016) '*Hibiscus cannabinus* L.–Kenaf: A Review Paper', *Journal of Natural Fibers*, 14(4), pp.466–484. Available at: <http://dx.doi.org/10.1080/15440478.2016.1240639>.
- Baeyens, J., Kang, Q., Appels, L., Dewil, R., Lv, Y. and Tan, T. (2015) 'Challenges and opportunities in improving the production of bio-ethanol', *Progress in Energy and Combustion Science*, 47, pp.60–88. Available at: <http://dx.doi.org/10.1016/j.peccs.2014.10.003>.
- Bai, W., Cao, Y., Liu, J., Wang, Q. and Jia, Z. (2016) 'Improvement of alkalophilicity of an alkaline xylanase Xyn11A-LC from *Bacillus* sp. SN5 by random mutation and Glu135 saturation mutagenesis. *BMC Biotechnology*, 16(1), p.77. Available at: <http://bmcbiotechnol.biomedcentral.com/articles/10.1186/s12896-016-0310-9>.
- Bailey, M. J., Biely, P. and Poutanen, K. (1992) 'Interlaboratory testing of methods for assay of xylanase activity', *Journal of Biotechnology*, 23(3), pp:257-270
- Balan, V. (2014) 'Current Challenges in Commercially Producing Biofuels from Lignocellulosic Biomass', *ISRN Biotechnology*, 2014(i), pp.1–31. Available at: <http://www.hindawi.com/journals/isrn/2014/463074/>.

- Banerjee, S., Mudliar, S., Sen, R., Giri, B., Satpute, D., Chakrabarti, T. and Pandey, R.A. (2010) 'Commercializing lignocellulosic bioethanol: technology bottlenecks and possible remedies', *Biofuels, Bioproducts and Biorefining*, (4), pp.77–93.
- Beg, M.A., Shivangi, Thakur, S.C. and Meena, L.S. (2018) 'Structural Prediction and Mutational Analysis of Rv3906c Gene of *Mycobacterium tuberculosis* H37Rv to Determine Its Essentiality in Survival', *Advances in Bioinformatics*.
- Beg, Q.K., Kapoor, M., Mahajan, L. and Hoondal, G.S. (2001) 'Microbial xylanases and their industrial applications: A review', *Applied Microbiology and Biotechnology*, 56(3–4), pp.326–338.
- Bendtsen, J.D., Nielsen, H., Von Heijne, G. and Brunak, S. (2004) 'Improved prediction of signal peptides: SignalP 3.0', *Journal of Molecular Biology*, 340(4), pp.783–795.
- Berg, J.M., Tymoczko, J.L. and Stryer, L. (2002) *Biochemistry*, (5<sup>th</sup> edn), W. H. Freeman
- Berlin, A., Gilkes, N., Kilburn, D., Bura, R., Markov, A., Skomarovsky, A., Okunev, O., Gusakov, A., Maximenko, V., Gregg, D., Sinitsyn, A. and Saddler, J. (2005) 'Evaluation of novel fungal cellulase preparations for ability to hydrolyze softwood substrates - Evidence for the role of accessory enzymes', *Enzyme and Microbial Technology*, 37(2), pp.175–184.
- Bhagavan, N.V. and Ha, C.-E. (2015) 'Three-Dimensional Structure of Proteins and Disorders of Protein Misfolding', *Essentials of Medical Biochemistry*, pp.31–51.
- Bhowmick, G. De, Sarmah, A.K. and Sen, R. (2018) 'Lignocellulosic biorefinery as a model for sustainable development of biofuels and value added products', *Bioresource Technology*, 247, pp.1144–1154. Available at: <https://doi.org/10.1016/j.biortech.2017.09.163>.
- Biely, P., Vrsanska, M., Tenkanen, M. and Kluepfel, D. (1997) 'Endo-B -1,4-xylanase families: differences in catalytic properties', *Journal of biotechnology*, 57, pp.151–166.
- Black, M. J., Whittaker, C., Hosseini, S. A., Diaz-Chavez, R., Woods, J. and Murphy, R. J. (2011) 'Life Cycle Assessment and sustainability methodologies for assessing industrial crops, processes and end products', *Industrial Crops and Products*, 34(2), pp. 1332-1339.

- Bloom, J.D. and Arnold, F.H. (2009) 'In the light of directed evolution: Pathways of adaptive protein evolution', *Proceedings of the National Academy of Sciences*, 106(Supplement\_1), pp.9995–10000. Available at: <http://www.pnas.org/cgi/doi/10.1073/pnas.0901522106>.
- Boonyapakron, K., Jaruwat, A., Liwnaree, B., Nimchua, T., Champreda, V. and Chitnumsub, P. (2017) 'Structure-based protein engineering for thermostable and alkaliphilic enhancement of endo- $\beta$ -1,4-xylanase for applications in pulp bleaching', *Journal of Biotechnology*. Available at: <http://dx.doi.org/10.1016/j.jbiotec.2017.07.035>.
- Bradford, M.M. (1976) 'A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding', *Analytical Biochemistry*, 72(1–2), pp.248–254.
- Chan, M. K., Mukund, S., Kletzin, A., Adams, M. W. W. and Rees, D. C. (1995) 'Structure of a hyperthermophilic tungstopterin enzyme, aldehyde ferredoxin oxidoreductase', *Science*, 267(5203), pp.1463–1469.
- Cheng, W.-Y., Haque Akanda, J.M. and Nyam, K.-L. (2016) 'Kenaf seed oil: A potential new source of edible oil', *Trends in Food Science & Technology*, 52, pp.57–65. Available at: <http://linkinghub.elsevier.com/retrieve/pii/S0924224415301242>.
- Cheng, Y. S., Chen, C. C., Huang, J. W., Ko, T. P., Huang, Z. and Guo, R. T. (2015) 'Improving the catalytic performance of a GH11 xylanase by rational protein engineering', *Applied Microbiology and Biotechnology*, 99(22), pp.9503–9510.
- Cheng, Z., Lu, B. R., Sameshima, K., Fu, D. X. and Chen, J. K. (2004) 'Identification and genetic relationships of kenaf (*Hibiscus cannabinus* L.) germplasm revealed by AFLP analysis', *Genetic Resources and Crop Evolution*, 51(4), pp.393–401.
- Chica, R.A., Doucet, N. and Pelletier, J.N. (2005) 'Semi-rational approaches to engineering enzyme activity: Combining the benefits of directed evolution and rational design', *Current Opinion in Biotechnology*, 16(4), pp.378–384.
- Chundawat, S.P.S., Balan, V. and Dale, B.E. (2008) 'High-throughput microplate technique for enzymatic hydrolysis of lignocellulosic biomass', *Biotechnology and Bioengineering*, 99(6), pp.1281–1294.



- Clayton, A. (2016) The Potential Role of Bioscience Industries in Small Developing Economies. In: Badal, S. Pharmacognosy: Fundamentals, Applications and Strategy. Academic Press, pp. 677–685.
- Cuyvers, S., Dornez, E., Delcour, J.A. and Courtin, C.M. (2012) ‘Occurrence and functional significance of secondary carbohydrate binding sites in glycoside hydrolases’, *Critical Reviews in Biotechnology*, 32(2), pp.93–107. Available at: <http://www.tandfonline.com/doi/full/10.3109/07388551.2011.561537>.
- Cuyvers, S., Dornez, E., Delcour, J.A. and Courtin, C.M. (2011a.) ‘The secondary substrate binding site of the *Pseudoalteromonas haloplanktis* GH8 xylanase is relevant for activity on insoluble but not soluble substrates’, *Applied Microbiology and Biotechnology*, 92(3), pp.539–549.
- Cuyvers, S., Dornez, E., Rezaei, M. N., Pollet, A., Delcour, J. A. and Courtin, C. M. (2011b) ‘Secondary substrate binding strongly affects activity and binding affinity of *Bacillus subtilis* and *Aspergillus niger* GH11 xylanases’, *FEBS Journal*, 278(7), pp.1098–1111.
- Cuyvers, S., Hendrix, J., Dornez, E., Engelborghs, Y., Delcour, J. A. and Courtin, C. M. (2011c) ‘Both substrate hydrolysis and secondary substrate binding determine xylanase mobility as assessed by FRAP’, *Journal of Physical Chemistry B*, 115(16), pp.4810–4817.
- Datta, R., Maher, M.A., Jones, C. and Brinker, R.W. (2011) ‘Ethanol-the primary renewable liquid fuel’, *Journal of Chemical Technology and Biotechnology*, 86(4), pp.473–480.
- Davies, G.J., Wilson, K.S. and Henrissat, B. (1997) ‘Nomenclature for sugar-binding subsites in glycosyl hydrolases’, *The Biochemical journal*, 321, pp.557–559.
- Decker, S. R., Brunecky, R., Tucker, M. P., Himmel, M. E. and Selig, M. J. (2009) ‘High-throughput screening techniques for biomass conversion’, *Bioenergy Research*, 2(4), pp.179–192.
- Demartini, J. and Wyman, C.E. (2013) ‘High-Throughput Pretreatment and Hydrolysis Systems for Screening Biomass Species in Aqueous Pretreatment of Plant Biomass’, *Aqueous Pretreatment of Plant Biomass for Biological and Chemical Conversion to Fuels and Chemicals*, (2), pp.471–488.
- Drummond, D.A., Iverson, B.L., Georgiou, G. and Arnold, F.H. (2005) ‘Why high-error-rate random mutagenesis libraries are enriched in functional and improved proteins’, *Journal of Molecular Biology*, 350(4), pp.806–816.

- Dubnovitsky, A.P., Kapetaniou, E.G. and Papageorgiou, A.C. (2009) 'Enzyme adaptation to alkaline pH: Atomic resolution (1.08 Å) structure of phosphoserine aminotransferase from *Bacillus alcalophilus*', *Protein Science*, 14(1), pp.97–110.
- Ehrt, S. and Schnappinger, D. (2003). *Isolation of Plasmids from E. coli by Alkaline Lysis* in Casali, N. and Preston, A. *E. coli Plasmid Vectors: Methods and Applications*, Humana Press, pp.75-78.
- Elias, M., Wieczorek, G., Rosenne, S. and Tawfik, D.S. (2014) 'The universality of enzymatic rate-temperature dependency', *Trends in Biochemical Sciences*, 39(1), pp.1–7. Available at: <http://dx.doi.org/10.1016/j.tibs.2013.11.001>.
- Erickson, B., Nelson and Winters, P. (2012) 'Perspective on opportunities in industrial biotechnology in renewable chemicals', *Biotechnology Journal*, 7(2), pp.176–185.
- Farzad, S., Mandegari, M. A., Guo, M., Haigh, K. F., Shah, N. and Görgens, J. F. (2017) 'Multi-product biorefineries from lignocelluloses: A pathway to revitalisation of the sugar industry?', *Biotechnology for Biofuels*, 10(1), pp.1–24.
- Felsenstein Joseph (1985) 'Confidence Limits on Phylogenies: an Approach Using the Bootstrap; Confidence Limits on Phylogenies: an Approach Using the Bootstrap', *Evolution*, 39(4), pp.783–791.
- Ferreira De Freitas, R. and Schapira, M. (2017) 'A systematic analysis of atomic protein-ligand interactions in the PDB', *MedChemComm*, 8(10), pp.1970–1981.
- Finch, H.J.S., Samuel, A.M. and Lane, G.P.F. (2014) *Industrial crops*. In: *Lockhart & Wiseman's Crop Husbandry Including Grassland*. Woodhead Publishing, pp. 387–395.
- Fushinobu, S., Ito, K., Konno, M., Wakagi, T. and Matsuzawa, H. (1998) 'Crystallographic and mutational analyses of an extremely acidophilic and acid-stable xylanase: biased distribution of acidic residues and importance of Asp37 for catalysis at low pH. *Protein Engineering Design and Selection*, 11(12), pp.1121–1128.

- Ganguly, A., Das, S., Bhattacharya, A., Dey, A. and Chatterjee, P. K. (2013) 'Enzymatic hydrolysis of water hyacinth biomass for the production of ethanol: Optimization of driving parameters. *Indian Journal of Experimental Biology*, 51(7), pp.556–566.
- Gasteiger, E., Hoogland, C., Gattiker, A., Duvaud, S. S., Wilkins, M. R., Appel, R. D. and Bairoch, A. S. (2005) *Protein Identification and Analysis Tools on the ExPASy Server*, in: Walker, J. M. (eds) *The Proteomics Protocols Handbook*, Totowa, NJ: Humana Press Inc., pp.571–607. Available at: <http://dx.doi.org/10.1385/1-59259-890-0:571>.
- Georis, J., Esteves, Frederic D. L., Lamotte-Brasseur, J., Bougnet, V., Giannotta, F., Frère, J-M.,
- Devreese, B. and Granier, B. (2008) 'An additional aromatic interaction improves the thermostability and thermophilicity of a mesophilic family 11 xylanase: Structural basis and molecular study', *Protein Science*, 9(3), pp.466–475. Available at: <http://doi.wiley.com/10.1110/ps.9.3.466>.
- Gírio, F M., Fonseca, C., Carvalheiro, F., Duarte, L. C. and Marques, S. (2010) 'Hemicelluloses for fuel ethanol : A review', *Bioresource Technology*, 101(13), pp.4775–4800. Available at: <http://dx.doi.org/10.1016/j.biortech.2010.01.088>.
- Goh, C.S., Tan, K.T., Lee, K.T. and Bhatia, S. (2010) 'Bio-ethanol from lignocellulose: Status, perspectives and challenges in Malaysia', *Bioresource Technology*, 101(13), pp.4834–4841. Available at: <http://dx.doi.org/10.1016/j.biortech.2009.08.080>.
- Gruber, K., Klintschar, G., Hayn, M., Schlacher, A., Steiner, W. and Kratky, C. (1998) 'Thermophilic Xylanase from *Thermomyces lanuginosus* : High-Resolution X-ray Structure and Modeling Studies', *Biochemistry*, 37(39), pp.13475–13485. Available at: <http://pubs.acs.org/doi/abs/10.1021/bi980864l>.
- Hakulinen, N., Turunen, O., Jänis, J., Leisola, M. and Rouvinen, J. (2003) 'Three-dimensional structures of thermophilic B-1,4-xylanases from *Chaetomium thermophilum* and *Nonomuraea flexuosa* Comparison of twelve xylanases in relation to their thermal stability', *European Journal of Biochemistry*, 270(7), pp.1399–1412.

- Hanahan, D., Jessee, J. and Bloom, F.R. (1991) 'Bacterial Genetic Systems', *Methods in Enzymology*, 204(1970), pp.63–113. Available at: <http://www.sciencedirect.com/science/article/pii/007668799104006A>.
- Haro, P., Perales, A.L.V., Arjona, R. and Ollero, P. (2013) 'Thermochemical biorefineries with multiproduction using a platform chemical', *Biofuels, Bioproducts and Biorefining*, 8, pp.155–170.
- Hasunuma, T., Okazaki, F., Okai, N., Hara, K. Y., Ishii, J. and Kondo, A. (2012) 'A review of enzymes and microbes for lignocellulosic biorefinery and the possibility of their application to consolidated bioprocessing technology', *Bioresource Technology*. Available at: <http://dx.doi.org/10.1016/j.biortech.2012.10.047>.
- Havukainen, R., Törrönen, A., Laitinen, T. and Rouvinen, J. (1996.) 'Covalent binding of three epoxyalkyl xylosides to the active site of endo-1,4-xylanase II from *Trichoderma reesei*', *Biochemistry*, 35(29), pp.9617–9624.
- Henrissat, B. (1991) 'A Classification of Glycosyl Hydrolases Based on Sequence Similarities Amino Acid', *Journal of Biochemistry*, 280, pp.309–316.
- Hess, B., Kutzner, C., Van Der Spoel, D. and Lindahl, E. (2008) 'GRGMACS 4: Algorithms for highly efficient, load-balanced, and scalable molecular simulation', *Journal of Chemical Theory and Computation*, 4(3), pp.435–447.
- Himmel, M. E., Ding, S.-Y., Johnson, D. K., Adney, W. S., Nimlos, M. R., Brady, J. W. (2007) 'Biomass recalcitrance: Engineering plants and enzymes for biofuels production', *Science*, 315(5813), pp.804–807. Available at: <http://www.sciencemag.org/cgi/doi/10.1126/science.1137016>.
- Hoffman, Z. B., Zanzorini, L. M., Cota, J., Diogo, J. A., Almeida, G. B., Damásio, A. R. L., Squina, F., Murakami, M. M. T. and Ruller, R. (2016) 'Xylan-specific carbohydrate-binding module belonging to family 6 enhances the catalytic performance of a GH11 endo-xylanase', *New Biotechnology*, 33(4), pp.467–472.
- Iqbal, H.M.N., Kyazze, G. and Keshavarz, T. (2013) 'Advances in the valorization of lignocellulosic materials by biotechnology: An overview', *BioResources*, 8(2), pp.3157–3176.

- Irfan, M., Gonzalez, C. F., Raza, S., Rafiq, M., Hasan, F., Khan, S. and Shah, A. A. (2018) 'Improvement in thermostability of xylanase from *Geobacillus thermodenitrificans* C5 by site directed mutagenesis', *Enzyme and Microbial Technology*, 111(January), pp.38–47.
- Irfan, M., Ibrahim, H., Ozer, A., Tuncel, M., Osman, A., Hasan, F. and Ali, A. (2016) 'C-Terminal proline-rich sequence broadens the optimal temperature and pH ranges of recombinant xylanase from *Geobacillus thermodenitrificans* C5', *Enzyme and Microbial Technology*, 91, pp.34–41. Available at: <http://dx.doi.org/10.1016/j.enzmictec.2016.05.012>.
- Isikgor, F.H. and Becer, C.R. (2015) 'Lignocellulosic biomass: a sustainable platform for the production of bio-based chemicals and polymers', *Polymer Chemistry*, 6(25), pp.4497–4559. Available at: <http://xlink.rsc.org/?DOI=C5PY00263J>.
- Jeya, M., Thiagarajan, S., Lee, J.K. and Gunasekaran, P. (2009) 'Cloning and expression of GH11 xylanase gene from *Aspergillus fumigatus* MKU1 in *Pichia pastoris*', *Journal of Bioscience and Bioengineering*, 108(1), pp.24–29. Available at: <http://dx.doi.org/10.1016/j.jbiosc.2009.02.003>.
- Jia, L., Goncalves, G. A. L., Takasugi, Y., Mori, Y., Noda, S., Tanaka, T., Ichinose, H. and Kamiya, N. (2015) 'Effect of pretreatment methods on the synergism of cellulase and xylanase during the hydrolysis of bagasse', *Bioresource Technology*, 185, pp.158–164. Available at: <http://dx.doi.org/10.1016/j.biortech.2015.02.041>.
- Jommuengbout, P., Pinitglang, S., Kyu, K.L. and Ratanakhanokchai, K. (2009) 'Substrate-binding site of family 11 xylanase from *Bacillus firmus* K-1 by molecular docking', *Bioscience, biotechnology, and biochemistry*, 73(4), pp.833–839. Available at: [http://www.jstage.jst.go.jp/article/bbb/73/4/73\\_833/\\_article%5Cnpapers2://publication/uuid/18F11BC9-7FF1-417C-9D8E-16B7628AC31A](http://www.jstage.jst.go.jp/article/bbb/73/4/73_833/_article%5Cnpapers2://publication/uuid/18F11BC9-7FF1-417C-9D8E-16B7628AC31A).
- Jonet, M. A., Mahadi, N. M., Murad, A. M. A., Rabu, A., Bakar, F. D., Abu Rahim, R. A., Low, K. O. and Illias, R. M. (2012) 'Optimization of a heterologous signal peptide by site-directed mutagenesis for improved secretion of recombinant proteins in *Escherichia coli*', *Journal of Molecular Microbiology and Biotechnology*, 22(1), pp.48–58.

- Joshi, M. D., Sidhu, G., Pot, I., Brayer, G. D., Withers, S. G. and McIntosh, L. P. (2000) 'Hydrogen bonding and catalysis: A novel explanation for how a single amino acid substitution can change the pH optimum of a glycosidase', *Journal of Molecular Biology*, 299(1), pp.255–279.
- Karnik, A., Karnik, R. and Grefen, C. (2013) 'SDM-Assist software to design site-directed mutagenesis primers introducing "silent" restriction sites', *BMC bioinformatics*, 14(1), p.105. Available at: <http://www.biomedcentral.com/1471-2105/14/105>.
- Kelly, L.A., Mezulis, S., Yates, C., Wass, M. and Sternberg, M. (2015) 'The Phyre2 web portal for protein modelling, prediction, and analysis', *Nature Protocols*, 10(6), pp.845–858. Available at: <http://dx.doi.org/10.1038/nprot.2015-053>.
- Khor, B. Y., Tye, G. J., Lim, T.S., Noordin, R. and Choong, Y.S. (2014) 'The structure and dynamics of BmR1 protein from *Brugia malayi*: In silico approaches', *International Journal of Molecular Sciences*, 15(6), pp.11082–11099.
- Khow, O. and Suntrarachun, S. (2012) 'Strategies for production of active eukaryotic proteins in bacterial expression system', *Asian Pacific Journal of Tropical Biomedicine*, 2(2), pp.159–162.
- Kim, T., Mullaney, E. J., Porres, J. M., Roneker, K. R., Crowe, S., Rice, S., Ko, T., Ullah, A. H. J., Daly, C. B., Welch, R. and Lei, X. G. (2006) 'Shifting the pH profile of *Aspergillus niger* PhyA phytase to match the stomach pH enhances its effectiveness as an animal feed additive', *Applied and Environmental Microbiology*, 72(6), pp.4397–4403.
- Krengel, U. and Dijkstra, B.W. (1996) 'Three-dimensional structure of endo-B-1,4-xylanase 11 from *Aspergillus niger*: Molecular basis for its low pH optimum. *Journal of Molecular Biology*, 263(1), pp.70–78.
- Kumar, V., Dangi, A.K. and Shukla, P. (2018) 'Engineering Thermostable Microbial Xylanases Toward its Industrial Applications', *Molecular Biotechnology*, 60(3), pp.226–235. Available at: <https://doi.org/10.1007/s12033-018-0059-6>.
- Kumar, L., Dutt, D., Tapas, S. and Kumar, P. (2013) 'Purification, bio-chemical characterization, homology modeling and active site binding mode interactions of thermo-alkali-tolerant  $\beta$ -1,4 endoxylanase from *Coprinus cinereus* LK-D-NCIM-1369', *Biocatalysis and Agricultural Biotechnology*, 2(3), pp.267–277. Available at: <http://dx.doi.org/10.1016/j.bcab.2013.04.004>.

- Kumar, S., Tsai, C.-J. and Nussinov, R. (2000) 'Factors enhancing protein thermostability', *Protein Engineering, Design and Selection*, 13(3), pp.179–191.
- Labrou, N.E. (2010) 'Random mutagenesis methods for in vitro directed enzyme evolution', *Current protein & peptide science*, 11(1), pp.91–100.
- Laemmli, U.K. (1970) 'Cleavage of structural proteins during the assembly of the head of bacteriophage T4', *Nature*, 227(5259), pp.680–685.
- Lee, C.W., Wang, H.J., Hwang, J.K. and Tseng, C.P. (2014) 'Protein thermal stability enhancement by designing salt bridges: A combined computational and experimental study', *PLoS ONE*, 9(11).
- Lee, H. V, Hamid, S.B. and Zain, S.K. (2014) 'Conversion of lignocellulosic biomass to nanocellulose: structure and Chemical Process', *The Scientific World Journal*, 2014(July), pp.1–14.
- de Lemos E. F., Gouders, T., Lamotte-Brasseur, J., Rigali, S. and Frere, J. M. (2005) 'Improving the alkalophilic performances of the Xyl1 xylanase from *Streptomyces* sp. S38: Structural comparison and mutational analysis. In: *Protein Science*, 14, pp. 292–302.
- Leys, S., Pauly, A., Delcour, J.A. and Courtin, C.M. (2016) 'Modification of the secondary binding site of xylanases illustrates the impact of substrate selectivity on bread making', *Journal of Agricultural and Food Chemistry*, 64(26), pp.5400–5409.
- Li, F., Xie, J., Zhang, X. and Zhao, L. (2015) 'Improvement of the optimum pH of *Aspergillus niger* xylanase towards an alkaline pH by site-directed mutagenesis', *Journal of Microbiology and Biotechnology*, 25(1), pp.11–17.
- Li, Q., Sun, B., Jia, H., Hou, J., Yang, R., Xiong, K., Xu, Y. and Li, X. (2017) 'Engineering a xylanase from *Streptomyces rochei* L10904 by mutation to improve its catalytic characteristics', *International Journal of Biological Macromolecules*, 101, pp.366–372. Available at: <http://dx.doi.org/10.1016/j.ijbiomac.2017.03.135>.
- Li, W., Xu, S., Zhang, B., Zhu, Y., Hua, Y., Kong, X., Sun, L. and Hong, J. (2017) 'Directed evolution to improve the catalytic efficiency of urate oxidase from *Bacillus subtilis*', *PLoS ONE*, 12(5).

- Linares-Pastén, J., Aronsson, A. and Karlsson, E. (2016) ‘Structural considerations on the use of endo-xylanases for the production of prebiotic xylooligosaccharides from biomass’, *Current Protein & Peptide Science*, 17(999), pp.1–1. Available at:  
<http://www.eurekaselect.com/openurl/content.php?genre=article&doi=10.2174/1389203717666160923155209>.
- Liu, Y., Huang, L., Jia, L., Gui, S., Fu, Y., Zheng, D., Guo, W. and Lu, F. (2012) ‘Improvement of the acid stability of *Bacillus licheniformis* alpha amylase by site-directed mutagenesis’, *Process Biochemistry*, 58, pp.174–180.
- Ludwiczek, M.L., Heller, M., Kantner, T. and McIntosh, L.P. (2007), ‘A secondary xylan-binding site enhances the catalytic activity of a single-domain family 11 glycoside hydrolase’, *Journal of Molecular Biology*, 373(2), pp.337–354.
- Madadi, M., Tu, Y. and Abbas, A. (2017) ‘Recent status on enzymatic saccharification of lignocellulosic biomass for bioethanol production’, *Electronic Journal of Biology*, 13(2), pp.135–143.
- Mamo, G., Thunnissen, M., Hatti-Kaul, R. and Mattiasson, B. (2009) ‘An alkaline active xylanase: Insights into mechanisms of high pH catalytic adaptation’, *Biochimie*, 91(9), pp.1187–1196. Available at:  
<http://dx.doi.org/10.1016/j.biochi.2009.06.017>.
- Mansfeld, J., Vriend, G., Dijkstra, B. W., Veltman, O. R., Van Den Burg, B., Venema, G., Ulbrich-Hofmann, R. and Eijsink, V. G. H. (1997) ‘Extreme stabilization of a thermolysin-like protease by an engineered disulfide bond’, *Journal of Biological Chemistry*, 272(17), pp.11152–11156.
- Martinez, R. and Schwaneberg, U. (2013) ‘A roadmap to directed enzyme evolution and screening systems for biotechnological applications’, *Biological Research*, 46(4), pp.395–405.
- McCarter, J.D. and Stephen Withers, G. (1994) ‘Mechanisms of enzymatic glycoside hydrolysis’, *Current Opinion in Structural Biology*, 4(6), pp.885–892.
- McLachlan, M., Sullivan, R.P. and Zhao, H. (2009) ‘Directed enzyme evolution and high throughput screening’, *Biocatalysis for the Pharmaceutical Industry: Discovery , Development and Manufacturing*, pp.1–20. Available at:  
<http://scs.illinois.edu/~zhaogrp/publications/HZ72.pdf>.



- Messaoudi, A., Belguith, H. and Ben Hamida, J. (2013) 'Homology modeling and virtual screening approaches to identify potent inhibitors of VEB-1  $\beta$ -lactamase', *Theoretical Biology and Medical Modelling*, 10(1), pp.1–10.
- Meyer, A.S., Rosgaard, L. and Sørensen, H.R. (2009) 'The minimal enzyme cocktail concept for biomass processing', *Journal of Cereal Science*, 50(3), pp.337–344.
- Michaux, C., Pouyez, J., Mayard, A., Vandurm, P., Housen, I. and Wouters, J. (2010) 'Structural insights into the acidophilic pH adaptation of a novel endo-1,4- $\beta$ -xylanase from *Scytalidium acidophilum*', *Biochimie*, 92(10), pp:1407-1415.
- Morley, K.L. and Kazlauskas, R.J. (2005) 'Improving enzyme properties: When are closer mutations better?', *Trends in Biotechnology*, 23(5), pp.231–237.
- Morone, A. and Pandey, R. A. (2014) 'Lignocellulosic biobutanol production : Gridlocks and potential remedies', *Renewable and Sustainable Energy Reviews*, 37, pp. 21-35.
- Motta, F.L., Andrade, C.C.P. and Santana, M.H.A. (2013) *A Review of Xylanase Production by the Fermentation of Xylan: Classification, Characterization and Applications*, in: Chandel, A.K. and Silva, S.S. (eds.) *Sustainable Degradation of Lignocellulosic Biomass*. IntechOpen, pp. 251–275.
- Morris, G.M., Goodsell, D.S., Huey, R. and Olson, A.J. (1996) 'Distributed automated docking of flexible ligands to proteins: Parallel applications of AutoDock 2.4', *Journal of Computer-Aided Molecular Design*, 10(4), pp.293–304. Available at: <http://link.springer.com/10.1007/BF00124499>.
- Muilu, J., To, A., Pera, M. and Rouvinen, J. (1998) 'Functional Conformational Changes of Endo-1, 4-xylanase II From *Trichoderma reesei*', *Proteins: Structure, Function and Genetics*, 31(March 1997), pp.434–444.
- Mussatto, S. I., Dragone, G., Fernandes, M., Milagres, A. M. F. and Roberto, I. C. (2008) 'The effect of agitation speed, enzyme loading and substrate concentration on enzymatic hydrolysis of cellulose from brewer's spent grain', *Cellulose*, 15(5), pp: 711-721.

- Navarro, D., Couturier, M., da Silva, G. G. D., Berrin, J-G., Rouau, X., Asther, M. and Bignon, C. (2010) 'Automated assay for screening the enzymatic release of reducing sugars from micronized biomass', *Microbial Cell Factories*, 9, p.58. Available at: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2919459&tool=pmcentrez&rendertype=abstract>.
- Nielsen, J.E. and McCammon, J.A. (2003) 'Calculating pKa values in enzyme active sites', *Protein Science*, 12(9), pp.1894–1901. Available at: <http://doi.wiley.com/10.1110/ps.03114903>.
- Nisius, L. and Grzesiek, S. (2012) 'Key stabilizing elements of protein structure identified through pressure and temperature perturbation of its hydrogen bond network', *Nature Chemistry*, 4(9), pp.711–717. Available at: <http://dx.doi.org/10.1038/nchem.1396>.
- Novák, P. and Havlíček, V. (2016) *Protein Extraction and Precipitation*, in: Ciborowski, P. and Silberring, J. (2<sup>nd</sup> edn) *Proteomic Profiling and Analytical Chemistry: The Crossroads*, Elsevier, pp. 52–62.
- Octave, S. and Thomas, D. (2009) 'Biorefinery: Toward an industrial metabolism', *Biochimie*, 91(6), pp.659–664.
- Ohkuri, T. and Yamagishi, A. (2003) 'Increased thermal stability against irreversible inactivation of 3-isopropylmalate dehydrogenase induced by decreased van der Waals volume at the subunit interface', *Protein Engineering Design and Selection*, 16(8), pp.615–621.
- Packer, M.S. and Liu, D.R. (2015) 'Methods for the directed evolution of proteins', *Nature Reviews Genetics*, 16(7), pp.379–394. Available at: <http://www.nature.com/doi/10.1038/nrg3927>.
- Paës, G., Berrin, J.G. and Beaugrand, J. (2012a) 'GH11 xylanases: Structure/function/properties relationships and applications', *Biotechnology Advances*, 30(3), pp.564–592.
- Paës, G., Cortés, J., Siméon, T., O'Donohue, M. J. and Tran, V. (2012b) 'Thumb-Loops Up for Catalysis: a Structure/Function Investigation of a Functional Loop Movement in a GH11 Xylanase', *Computational and Structural Biotechnology Journal*, 1(2), pp: e201207001.

- Paës, G., Tran, V., Takahashi, M., Boukari, I. and O'Donohue, M. J. (2007) 'New insights into the role of the thumb-like loop in GH-11 xylanases', *Protein Engineering, Design and Selection*, 20(1), pp.15–23.
- Paës, G. and O'Donohue, M.J. (2006) 'Engineering increased thermostability in the thermostable GH-11 xylanase from *Thermobacillus xylanilyticus*', *Journal of Biotechnology*, 125(3), pp.338–350.
- Payan, F., Leone, P., Porciero, S., Furniss, C., Tahir, T., Williamson, G., Durand, A., Manzanares, P., Gilbert, H. J., Juge, N. and Roussel, A. (2004) 'The dual nature of the wheat xylanase protein inhibitor XIP-I: Structural basis for the inhibition of family 10 and family 11 xylanases', *Journal of Biological Chemistry*, 279(34), pp.36029–36037.
- Pérez-Pérez, J.M., Candela, H. and Micol, J.L. (2009) 'Understanding synergy in genetic interactions', *Trends in Genetics*, 25(8), pp.368–376.
- Petersen, T.N., Brunak, S., von Heijne, G. and Nielsen, H. (2011) 'SignalP 4.0: discriminating signal peptides from transmembrane regions', *Nature Methods*, 8(10), pp.785–786. Available at: <http://www.nature.com/articles/nmeth.1701>.
- Phitsuwan, P., Sakka, K. and Ratanakhanokchai, K. (2013) 'Improvement of lignocellulosic biomass in planta: A review of feedstocks , biomass recalcitrance , and strategic manipulation of ideal plants designed for ethanol production and processability', *Biomass and Bioenergy*, pp.1–16. Available at: <http://dx.doi.org/10.1016/j.biombioe.2013.08.027>.
- Piovesan, D., Minervini, G. and Tosatto, S.C.E. (2016) 'The RING 2.0 web server for high quality residue interaction networks', *Nucleic acids research*, 44(W1), pp.W367–W374.
- Plaza Del Pino, I.M., Ibarra-Molero, B. and Sanchez-Ruiz, J.M. (2000) 'Lower kinetic limit to protein thermal stability: A proposal regarding protein stability in vivo and its relation with misfolding diseases', *Proteins: Structure, Function and Genetics*, 40(1), pp.58–70.
- Pokhrel, S., Joo, J.C. and Yoo, Y.J. (2013) 'Shifting the optimum pH of *Bacillus circulans* xylanase towards acidic side by introducing arginine', *Biotechnology and Bioprocess Engineering*, 18(1), pp.35–42.
- Polizeli, M. L T M., Rizzatti, A. C S., Monti, R., Terenzi, H. F., Jorge, J. A. and Amorim, D. S. (2005) 'Xylanases from fungi: Properties and industrial applications', *Applied Microbiology and Biotechnology*, 67(5), pp.577–591.

- Pollard, Thomas, D., Lippincott-Schwartz, J., Earnshaw, William, C. and Johnson, Graham, T. (2017) *Biophysical Principles* in: Pollard, Thomas, D., Lippincott-Schwartz, J., Earnshaw, William, C. and Johnson, Graham, T., (eds.) *Cell Biology*. Elsevier, pp. 53–62.
- Pollet, A., Lagaert, S., Eneyskaya, E., Kulminskaya, A., Delcour, J. A. and Courtin, C. M. (2010) ‘Mutagenesis and subsite mapping underpin the importance for substrate specificity of the aglycon subsites of glycoside hydrolase family 11 xylanases’, *Biochimica et Biophysica Acta - Proteins and Proteomics*, 1804(4), pp.977–985. Available at: <http://dx.doi.org/10.1016/j.bbapap.2010.01.009>.
- Prajapati, A. S., Pawar, V. A., Panchal, K. J., Sudhir, A. P., Dave, B. R., Patel, D. H. and Subramanian, R. B. (2018) ‘Effects of substrate binding site residue substitutions of xynA from *Bacillus amyloliquefaciens* on substrate specificity’, *BMC Biotechnology*, 18(1), pp.1–10.
- Rajesh Kumar, P., Eswaramoorthy, S., Vithayathil, P.J. and Viswamitra, M.A. (2000) ‘The tertiary structure at 1.59 Å resolution and the proposed amino acid sequence of a family-11 xylanase from the thermophilic fungus *Paecilomyces varioti* Bainier’, *Journal of Molecular Biology*, 295(3), pp.581–593.
- Rajpal, G. and Arvan, P. (2013) *Disulfide Bond Formation*, in Kastin, A. J. (2<sup>nd</sup> edn) *Handbook of Biologically Active Peptides*, Elsevier Inc., pp. 1721-1729.
- Ramesh, M. (2016) ‘Kenaf (*Hibiscus cannabinus* L.) fibre based bio-materials: A review on processing and properties’, *Progress in Materials Science*, 78–79, pp.1–92. Available at: <http://dx.doi.org/10.1016/j.pmatsci.2015.11.001>.
- Reetz, M.T., Höbenreich, H., Soni, P. and Fernández, L. (2008) ‘A genetic selection system for evolving enantioselectivity of enzymes’, *Chemical Communications*, (43), p.5502. Available at: <http://xlink.rsc.org/?DOI=b814538e>.
- Renthal, R. (2008) ‘Buried water molecules in helical transmembrane proteins’, *Protein Science*, 17(2), pp.293–298.
- Rice, A.J., Truong, L., Johnson, M.E. and Lee, H. (2013) ‘A colorimetric assay optimization for high-throughput screening of dihydroorotase by detecting ureido groups’, *Analytical Biochemistry*, 441(1), pp.87–94. Available at: <http://dx.doi.org/10.1016/j.ab.2013.05.035>.
- Robinson, P.K. (2015) ‘Enzymes: principles and biotechnological applications’, *Essays In Biochemistry*, 59(0), pp.75–75.

- Roche, J., Caro, J. A., Norberto, D. R., Barthe, P., Roumestand, C., Schlessman, J. L., Garcia, A. E., García-Moreno, B. E. and Royer, C. A. (2012) Cavities determine the pressure unfolding of proteins. *Proceedings of the National Academy of Sciences of the United States of America*, 109(18), pp.6945–6950.
- Rother, K., Hildebrand, P. W., Goede, A., Gruening, B. and Preissner, R. (2009) ‘Voronoi: Analyzing packing in protein structures’, *Nucleic Acids Research*, 37(SUPPL. 1), pp.2008–2010.
- Rowell, R. and Stout, H. (1998) *Jute and kenaf*, in: Lewin, M. (3<sup>rd</sup> edn) *Handbook of Fiber Chemistry*, Marcel Dekker, Inc., pp. 465–504.
- Rubin-Pitel, S.B. and Zhao, H. (2006) ‘Recent advances in biocatalysis by directed enzyme evolution’, *Combinatorial chemistry & high throughput screening*, 9(4), pp.247–257. Available at: <http://scs.illinois.edu/~zhaogrp/publications/HZ43.pdf>.
- Saba, N., Jawaid, M., Hakeem, K. R., Paridah, M. T., Khalina, A. and Alothman, O. Y. (2015) ‘Potential of bioenergy production from industrial kenaf (*Hibiscus cannabinus* L.) based on Malaysian perspective’, *Renewable and Sustainable Energy Reviews*, 42, pp.446–459. Available at: <http://dx.doi.org/10.1016/j.rser.2014.10.029>.
- Sabini, E., Wilson, K. S., Danielsen, S., Schülein, M. and Davies, G. J. (2001) ‘Oligosaccharide binding to family 11 xylanases: Both covalent intermediate and mutant product complexes display 2,5B conformations at the active centre’, *Acta Crystallographica Section D: Biological Crystallography*, 57(9), pp.1344–1347.
- Salleh, A.B. (2012) ‘The Role of Arg157Ser in Improving the Compactness and Stability of ARM Lipase’, *Journal of Computer Science & Systems Biology*, 05(02), pp.39–46.
- Sambrook, J. F. and Russell, D. W. (2001) *Molecular Cloning: A Laboratory Manual* (3<sup>rd</sup> edn) Cold Spring Harbor Laboratory Press.
- Sapag, A., Wouters, J., Lambert, C., De Ioannes, P., Eyzaguirre, J., Depiereux, E. (2002) ‘The endoxylanases from family 11: Computer analysis of protein sequences reveals important structural and phylogenetic relationships’, *Journal of Biotechnology*, 95(2), pp.109–131.

- Schäfer, K., Magnusson, U., Scheffel, F., Schiefner, A., Sandgren, M. O. J., Diederichs, K., Welte, W., Hülsmann, A., Schneider, E. and Mowbray, S. L. (2004) 'X-ray structures of the maltose-maltodextrin-binding protein of the thermoacidophilic bacterium *Alicyclobacillus acidocaldarius* provide insight into acid stability of proteins', *Journal of Molecular Biology*, 335(1), pp.261–274.
- Schmidt, A., Gübitz, G.M. and Kratky, C. (1999) 'Xylan binding subsite mapping in the xylanase from *Penicillium simplicissimum* using xylooligosaccharides as cryo-protectant', *Biochemistry*, 38(8), pp.2403–2412.
- Schoenherr, S., Ebrahimi, M. and Czermak, P. (2018) 'Lignin degradation processes and the purification of valuable products', *Lignin: trends and applications*, pp. 29–63.
- Schwede, T., Diemand, A., Guex, N. and Peitsch, M.C. (2000) 'Protein structure computing in the genomic era', *Research in Microbiology*, 151(2), pp.107–112.
- Scopes, R.K., 2006. Enzyme Activity and Assays. *Encyclopedia of Life Sciences*, pp.1–6.
- Seeliger, D. and De Groot, B.L. (2010) 'Ligand docking and binding site analysis with PyMOL and Autodock/Vina', *Journal of Computer-Aided Molecular Design*, 24(5), pp.417–422.
- Shallom, D. and Shoham, Y. (2003) 'Microbial hemicellulases', *Current Opinion in Microbiology*, 6(3), pp.219–228.
- Sharma, M. and Kumar, A. (2013) 'Xylanases : An Overview', *British Biotechnology Journal*, 3(1), pp.1–28.
- Shaw, K. L., Grimsley, G. R., Yakovlev, G. I., Makarov, A. A. and Pace, C. N. (2001) 'The effect of net charge on the solubility, activity, and stability of ribonuclease Sa', *Protein Science*, 10(6), pp.1206–1215.
- Sheldon, R.A. (2014) 'Green and sustainable manufacture of chemicals from biomass: State of the art', *Green Chemistry*, 16(3), pp.950–963.
- Shrivastava, A., Kumar, J., Akhter, M., Alan, M. M. and Shaqiquzaman, M. (2016) 'In-Silico Assessment of Various PDB Entries of Pfldh Enzyme for their Use in SBDD', *Chemical Informatics*, 2(1), pp.1–9.

- Siloto, R.M.P. and Weselake, R.J. (2012) 'Site saturation mutagenesis: Methods and applications in protein engineering', *Biocatalysis and Agricultural Biotechnology*, 1(3), pp.181–189. Available at: <http://dx.doi.org/10.1016/j.bcab.2012.03.010>.
- Singh, R.S. (2014) *Industrial Biotechnology: An Overview*. In: Singh, R. S., Pandey, A. and Larroche, C. (eds) *Advances in Industrial Biotechnology*. India: I.K. International Publishing House Pvt. Ltd., pp. 1–35.
- Söding, J., Biegert, A. and Lupas, A.N. (2005) 'The HHpred interactive server for protein homology detection and structure prediction;', *Nucleic Acids Research*, 33(SUPPL. 2), pp.244–248.
- Song, L., Dumon, C., Siguier, B., André, I., Eneyskaya, E., Kulminkaya, A., Bozonnet, S. and O'Donohue, M. J. (2014) 'Impact of an N-terminal extension on the stability and activity of the GH11 xylanase from *Thermobacillus xylanilyticus*', *Journal of Biotechnology*, 174(1), pp.64–72. Available at: <http://dx.doi.org/10.1016/j.jbiotec.2014.01.004>.
- Song, L., Siguier, B., Dumon, C., Bozonnet, S., O'Donohue, M.J. (2012) 'Engineering better biomass-degrading ability into a GH11 xylanase using a directed strategy', *Biotechnology for biofuels*, 5(1), p.3. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22244361> <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC3299623>.
- Song, L., Laguerre, S., Dumon, C., Bozonnet, S. and 'Donohue, M. J. (2010) 'A high-throughput screening system for the evaluation of biomass-hydrolyzing glycoside hydrolases', *Bioresource Technology*, 101(21), pp.8237–8243. Available at: <http://dx.doi.org/10.1016/j.biortech.2010.05.097>.
- Soo Yee, T. (2010) *Cloning and biochemical characterization of recombinant xylanase from Aspergillus fumigatus RT-1*. MSc Thesis, Universiti Teknologi Malaysia.
- de Souza, A. R., de Araújo, G. C., Zanphorlin, L. M., Ruller, R., Franco, F. C., Torres, F. A.G., Mertens, J. A., Bowman, M. J., Gomes, E. and Da Silva, R. (2016) 'Engineering increased thermostability in the GH-10 endo-1, 4- $\beta$ -xylanase from *Thermoascus aurantiacus* CBMAI 756', *International Journal of Biological Macromolecules*, 93, pp.20–26. Available at: <http://dx.doi.org/10.1016/j.ijbiomac.2016.08.056>.

- Steiner, K. and Schwab, H. (2012) 'Recent Advances in Rational Approaches for Enzyme engineering', *Computational and Structural Biotechnology*, 3(September).
- Takano, K., Yamagata, Y. and Yutani, K. (2003) 'Buried water molecules contribute to the conformational stability of a protein', *Protein Engineering*, 16(1), pp.5–9.
- Talebnia, F., Karakashev, D. and Angelidaki, I. (2010) 'Bioresource Technology Production of bioethanol from wheat straw : An overview on pretreatment , hydrolysis and fermentation', *Bioresource Technology*, 101(13), pp.4744–4753.
- Talley, K. and Alexov, E. (2010) 'On the pH-optimum of activity and stability of proteins', *Proteins*, 78(12), pp.2699–2706. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3624763/pdf/nihms412728.pdf>.
- Tanakai, Y., Tsumoto, K., Yasutake, Y., Umetsu, M., Yao, M., Fukada, H., Tanaka, I. and Kumagai, I. (2004) 'How oligomerization contributes to the thermostability of an archaeon protein: Protein L-isoaspartyl-O-methyltransferase from *Sulfolobus tokodaii*', *Journal of Biological Chemistry*, 279(31), pp.32957–32967.
- Tang, W.L. and Zhao, H. (2009) 'Industrial biotechnology: Tools and applications', *Biotechnology Journal*, 4(12), pp.1725–1739.
- Tian, L., Liu, S., Wang, S. and Wang, L. (2016) 'Ligand-binding specificity and promiscuity of the main lignocellulolytic enzyme families as revealed by active-site architecture analysis', *Scientific Reports*, 6(1), p.23605. Available at: <http://www.nature.com/articles/srep23605>.
- Thompson, J.D., Higgins, D.G. and Gibson, T.J. (1994) 'CLUSTAL W: Improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice', *Nucleic Acids Research*, 22(22), pp.4673–4680.
- Tina, K.G., Bhadra, R. and Srinivasan, N. (2007) 'PIC: Protein Interactions Calculator', *Nucleic Acids Research*, 35(SUPPL.2), pp.473–476.



- Torpenholt, S., De Maria, L., Olsson, M. H. M., Christensen, L. H., Skjøt, M., Westh, P., Jensen, J. H. and Lo Leggio, L. (2015) 'Effect of mutations on the thermostability of *Aspergillus aculeatus*  $\beta$ -1,4-galactanase', *Computational and Structural Biotechnology Journal*, 13, pp.256–264. Available at: <http://dx.doi.org/10.1016/j.csbj.2015.03.010>.
- Törrönen, A. and Rouvinen, J. (1997) 'Structural and functional properties of low molecular weight endo-1,4-beta-xylanases', *Journal of biotechnology*, 57(1–3), pp.137–149.
- Torronen, A. and Rouvinen, J. (1995) 'Structural Comparison of Two Major endo-1,4-Xylanases from *Trichoderma reesei*', *Biochemistry*, 34(3), pp.847–856. Available at: <http://pubs.acs.org/doi/abs/10.1021/bi00003a019>.
- Törrönen, A., Harkki, A. and Rouvinen, J. (1994) 'Three-dimensional structure of endo-1,4-beta-xylanase II from *Trichoderma reesei*: two conformational states in the active site', *The EMBO journal*, 13(11), pp.2493–501. Available at: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=395120&tool=pmcentrez&rendertype=abstract>.
- Tu, T., Luo, H., Meng, K., Cheng, Y., Ma, R., Shi, P., Huang, H., Bai, Y., Wang, Y., Zhang, L. and Yao, B. (2015) 'Improvement in thermostability of an *Achaetomium* sp. strain Xz8 endopolygalacturonase via the optimization of charge-charge interactions', *Applied and Environmental Microbiology*, 81(19), pp.6938–6944.
- Tye, Y.Y., Lee, K.T., Wan Abdullah, W.N. and Leh, C.P. (2016) 'The world availability of non-wood lignocellulosic biomass for the production of cellulosic ethanol and potential pretreatments for the enhancement of enzymatic saccharification', *Renewable and Sustainable Energy Reviews*, 60, pp.155–172. Available at: <http://dx.doi.org/10.1016/j.rser.2016.01.072>.
- Vajda, S. and Kozakov, D. (2010) 'Convergence and combination of methods in protein-protein docking', *Curr Opin Struct Biol.*, 71(2), pp.233–236.
- Van Dyk, J.S. and Pletschke, B.I. (2012) 'A review of lignocellulose bioconversion using enzymatic hydrolysis and synergistic cooperation between enzymes-Factors affecting enzymes, conversion and synergy', *Biotechnology Advances*, 30(6), pp.1458–1480. Available at: <http://dx.doi.org/10.1016/j.biotechadv.2012.03.002>.

- Vandermarliere, E., Bourgois, T. M., Rombouts, S., van Campenhout, S., Volckaert, G., Strelkov, S. V., Delcour, J. A., Rabijns, A. and Courtin, C. M. (2008) 'Crystallographic analysis shows substrate binding at the -3 to +1 active-site subsites and at the surface of glycoside hydrolase family 11 endo-1,4- $\beta$ -xylanases', *Biochemical Journal*, 410(1), pp.71–79. Available at: <http://biochemj.org/lookup/doi/10.1042/BJ20071128>.
- Vardakou, M., Dumon, C., Murray, J. W., Christakopoulos, P., Weiner, D. P., Juge, N., Lewis, R. J., Gilbert, H. J. and Flint, J. E. (2008) 'Understanding the Structural Basis for Substrate and Inhibitor Recognition in Eukaryotic GH11 Xylanases', *Journal of Molecular Biology*, 375(5), pp.1293–1305.
- Vaskan, P., Pachón, E.R. and Gnansounou, E. (2017) *Life Cycle Assessment of Sugar Crops and Starch-Based Integrated Biorefineries*, in Gnansounou, E. and Pandey, A. (1<sup>st</sup> edn) *Life-Cycle Assessment of Biorefineries*, Elsevier, pp. 97-146.
- Verardi, A., Ricca, E., De Bari, I. and Calabrò, V. (2012) *Hydrolysis of lignocellulosic biomass: current status of processes and technologies and future perspectives*, in: Lima, M. A. P. (1<sup>st</sup> edn) *Bioethanol*. InTech, pp: 95-122.
- Vieille, C. and Zeikus, G.J. (2001) 'Hyperthermophilic Enzymes: Sources, Uses, and Molecular Mechanisms for Thermostability', *Microbiology and Molecular Biology Reviews*, 65(1), pp.1–43.
- Villeret, V., Clantin, B., Tricot, C., Legrain, C., Roovers, M., Stalon, V., Glansdorff, N. and Van Beeumen, J. (1998) 'The crystal structure of *Pyrococcus furiosus* ornithine carbamoyltransferase reveals a key role for oligomerization in enzyme stability at extremely high temperatures', *Proceedings of the National Academy of Sciences of the United States of America*, 95(6), pp.2801–2806.
- Visser, E., Leal, T., de Almeida, M. and Guimarães, V. (2015) 'Increased enzymatic hydrolysis of sugarcane bagasse from enzyme recycling', *Biotechnology for Biofuels*, 8(1), p.5. Available at: <http://www.biotechnologyforbiofuels.com/content/8/1/5>.
- Vohra, M., Manwar, J., Manmode, R., Padgilwar, S. and Patil, S. (2014) 'Bioethanol production: Feedstock and current technologies', *Journal of Environmental Chemical Engineering*, 2(1), pp.573–584. Available at: <http://dx.doi.org/10.1016/j.jece.2013.10.013>.

- Vyas, N., Vyas, M.N. and Quioco, F. (1991a) 'Comparison of the Periplasmic Receptors for L-Arabinose, D-glucose/D-galactose and D-ribose', *The journal of biological chemistry*, 266, pp.5226–5237.
- Vyas, N.K. (1991b) 'Atomic features of protein-carbohydrate interactions', *Current Opinion in Structural Biology*, 1, pp.732–740.
- Wakarchuk, W. W., Campbell, R. L., Sung, W. L., Davoodi, J. and Yaguchi, M. (1994) 'Mutational and crystallographic analyses of the active site residues of the *Bacillus circulans* xylanase. *Protein Science*, 3(3), pp.467–475.
- Wan Azelee, N. I., Jahim, J. M., Ismail, A. H., Fuzi, S. F. Z. M., Rahman, R. A. and Illias, R. M. (2016) 'High xylooligosaccharides (XOS) production from pretreated kenaf stem by enzyme mixture hydrolysis', *Industrial Crops and Products*, 81, pp.11–19.
- Wan Azelee, N.I., Md Jahim, J., Rabu, A., Abdul Murad, A. M., Abu Bakar, F. D. and Illias, R. M. (2014) 'Efficient removal of lignin with the maintenance of hemicellulose from kenaf by two-stage pretreatment process', *Carbohydrate Polymers*, 99, pp.447–453. Available at: <http://dx.doi.org/10.1016/j.carbpol.2013.08.043>.
- Wang, Y., Feng, S., Zhan, T., Huang, Z., Wu, G. and Liu, Z. (2013) 'Improving catalytic efficiency of endo- $\beta$ -1, 4-xylanase from *Geobacillus stearothermophilus* by directed evolution and H179 saturation mutagenesis', *Journal of Biotechnology*, 168(4), pp.341–347. Available at: <http://dx.doi.org/10.1016/j.jbiotec.2013.09.014>.
- Waters, M.L. (2002) 'Aromatic interactions in model systems', *Current Opinion in Chemical Biology*, 6(6), pp.736–741.
- Webb, B. and Sali, A. (2017) 'Comparative Protein Structure Modeling Using MODELLER', *Curr Protoc Bioinformatics*, 54(ii), pp.1–55.
- Webber, C.L.I. and Bledsoe, V.K. (2002) 'Kenaf Yield Components and Plant Composition', *Trends in new crops and new uses*, pp.348–357.
- Wen, F., Mclachlan, M. and Zhao, H. (2008) 'Directed Evolution: Novel and improved enzymes', *Web Encyclopedia of Chemical Biology*, pp.1–7.
- Wilson, P.N., Wade, J.C. and Leones, J.P. (1995) 'The economics of commercializing new industrial crops', *Agribusiness*, 11(1), pp.45–55.
- Wouters, J., Georis, J., Engher, D., Vandenhoute, J., Dusart, J., Frere, J. M., Depiereux, E. and Charlier, P. (2001) 'Crystallographic analysis of family 11 endo- $\beta$ -1,4-

- xylanase Xyl1 from *Streptomyces* sp. S38. *Acta Crystallographica Section D Biological Crystallography*, 57(12), pp.1813–1819. Available at: <http://scripts.iucr.org/cgi-bin/paper?S0907444901015153>.
- Wu, D., Guo, X., Lu, J., Sun, X., Li, F., Chen, Y. and Xiao, D. (2013) ‘A rapid and efficient one-step site-directed deletion, insertion, and substitution mutagenesis protocol’, *Analytical Biochemistry*, 434(2), pp.254–258. Available at: <http://dx.doi.org/10.1016/j.ab.2012.11.028>.
- Yang, H., Li, J., Du, G. and Liu, L. (2017) ‘Microbial Production and Molecular Engineering of Industrial Enzymes: Challenges and Strategies. In: *Biotechnology of Microbial Enzymes: Production, Biocatalysis and Industrial Applications*. Academic Press, pp. 151–165.
- Yang, M., Zhang, J., Kuittinen, S., Vepsäläinen, J., Soininen, P. and Keinänen, M. (2015) ‘Enhanced sugar production from pretreated barley straw by additive xylanase and surfactants in enzymatic hydrolysis for acetone – butanol – ethanol fermentation’, *Bioresource Technology*, 189, pp.131–137. Available at: <http://dx.doi.org/10.1016/j.biortech.2015.04.008>.
- Yi, X., Shi, Y., Xu, H., Li, W., Xie, J., Yu, R., Zhu, J., Cao, Y. and Qiao, D. (2010) ‘Hyperexpression of two *Aspergillus niger* xylanase genes in *Escherichia coli* and characterization of the gene products’, *Brazilian Journal of Microbiology*, 41(3), pp.778–786.
- Yuan, S-F., Wu, T-H., Lee, H-L., Hsieh, H-Y., Lin, W-L., Yang, B., Chang, C-K., Li, Q., Gao, J., Huang, C-H., Ho, M-C., Guo, R-T. and Liang, P-H. (2015) ‘Biochemical Characterization and Structural Analysis of a Bifunctional Cellulase/Xylanase from *Clostridium thermocellum*’, *Journal of Biological Chemistry*, 290(9), pp.5739–5748. Available at: <http://www.jbc.org/lookup/doi/10.1074/jbc.M114.604454>.
- Zabed, H., Sahu, J N., Suely, A., Boyce, A N. and Faruq, G. (2016) ‘Bioethanol production from renewable sources : Current perspectives and technological progress’, *Renewable and Sustainable Energy Reviews*, (December), pp.1–27. Available at: <http://dx.doi.org/10.1016/j.rser.2016.12.076>.
- Zdobnov, E.M. and Apweiler, R. (2001) ‘InterProScan - An integration platform for the signature-recognition methods in InterPro’, *Bioinformatics*, 17(9), pp.847–848.

- Zhang, B-Z., Zhang, X., An, X-P., Ran, D-L., Zhou, Y-S., Lu, J. and Tong, Y-G. (2009) 'An easy-to-use site-directed mutagenesis method with a designed restriction site for convenient and reliable mutant screening', *Journal of Zhejiang University SCIENCE B*, 10(6), pp.479–482. Available at: <http://www.springerlink.com/index/10.1631/jzus.B0820367>.
- Zheng, H., Liu, Y., Sun, M., Han, Y., Wang, J., Sun, J and Lu, F. (2014) 'Improvement of alkali stability and thermostability of *Paenibacillus campinasensis* Family-11 xylanase by directed evolution and site-directed mutagenesis', *Journal of Industrial Microbiology and Biotechnology*, 41(1), pp.153–162.

## APPENDICES

### APPENDIX A

#### Medium, solutions and buffers preparation

##### Appendix A1: Medium Preparation

1.	Luria Bertani (LB) medium	100 ml
	Bacto-tryptone	1 g
	Bacto-yeast extracts	0.5 g
	NaCl	0.5 g

To make LB agar, the same ingredient used with addition 1.5 g agar before autoclaving.

2.	TSS Reagent	40 ml
	Bacto-tryptone	0.4 g
	Bacto-yeast extract	0.2 g
	NaCl	0.2 g
	Polyethylene glycol (PEG)	4.0 g
	Dinethyl sulfoxide	2.0 ml
	MgCl <sub>2</sub>	0.194 g

## Appendix A2: Autoinduction medium

### 1. Luria-Bertani for Autoinduction (LBA) medium

Composition: 1% w/v tryptone, 0.5% yeast extract, 0.1% NaCl

For 1 liter LBA medium

- 1) The following ingredients were dissolved in 950 ml distilled water: 10 g tryptone, 5 g yeast extract, 1 g NaCl .
- 2) Distilled water was added to bring the volume to 1 liter
- 3) The solution was autoclaved for 15 min at 121 °C
- 4) The solution has cooled to ~ 55 °C before used.

### 2. 20X NPS stock solution

Composition: 0.5 M (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 1 M KH<sub>2</sub>PO<sub>4</sub>, 1 M Na<sub>2</sub>HPO<sub>4</sub>.

For 1 liter 20 X NPS stock solution:

- 1) The following ingredients in sequence in beaker; stir until all were dissolved in 900 ml distilled water:

Component	1 liter	Conc. at 1 x
dH <sub>2</sub> O	900 ml	
(NH <sub>4</sub> ) <sub>2</sub> SO <sub>4</sub>	66	25 mM
KH <sub>2</sub> PO <sub>4</sub>	136	50 mM
Na <sub>2</sub> HPO <sub>4</sub>	142	50 mM

- 2) Distilled water was added to bring the volume to 1 liter
- 3) The solution was autoclaved for 15 min at 121 °C
- 4) The solution has to be cooled to ~ 55 °C before used.

### 3. 50 X 5052 Stock solution

Composition (5052= 0.5% glycerol, 0.05% glucose, 0.2% α-lactose)

For 1 liter 50 X 5052 stock solution:

- 1) The following ingredients in sequence in beaker; stir until all were dissolved in 900 ml distilled water:

Component	1 liter
Glycerol (weight in beaker)	250 g
H <sub>2</sub> O	730 ml
Glucose	25 g
α-lactose	100 g

4. 1000X trace metals mixture stock solution  
(100 ml in ~50 mM HCl)

- 1) All metal stock solutions were prepared in miliQ H<sub>2</sub>O, except for FeCl<sub>3</sub>, which is dissolved in ~0.1M HCl, as noted in the table below. Combine the metal solutions as in the table below:

Component	Volume	MW	1 X concentration
H <sub>2</sub> O	36 ml	-	-
0.1 M FeCl <sub>3</sub> .6H <sub>2</sub> O <i>yellow</i> (dissolved in ~0.1M HCl)	50 ml	270.30	50 μM Fe
1 M CaCl <sub>2</sub>	2 ml	110.99	20 μM Ca
1M MnCl <sub>2</sub> .4H <sub>2</sub> O	1 ml	197.91	10 μM
1 M ZnSO <sub>4</sub> .7H <sub>2</sub> O	1 ml	287.56	10 μM Zn
0.2 M CoCl <sub>2</sub> .6H <sub>2</sub> O <i>pink</i>	1 ml	237.95	2 μM Co
0.1 M CuCl <sub>2</sub> .2H <sub>2</sub> O <i>blue</i>	2 ml	170.486	2 μM Cu
0.2 M NiCl <sub>2</sub> .6H <sub>2</sub> O <i>green</i>	1 ml	237.72	2 μM Ni
0.1 M Na <sub>2</sub> MoO <sub>4</sub> .2H <sub>2</sub> O	2 ml	241.98	2 μM Mo
0.1 M Na <sub>2</sub> SeO <sub>3</sub> .5H <sub>2</sub> O	2 ml	263.03	2 μM se
0.1 M H <sub>2</sub> BO <sub>3</sub>	2 ml	61.83	2 μM H <sub>3</sub> BO <sub>3</sub>

- 2) The stock solutions of the individual metals were autoclaved, except for 0.1 M FeCl<sub>2</sub> in 1/100 volume of concentrated HCl.
- 3) For Na<sub>2</sub>SeO<sub>3</sub>, a brief precipitate appeared upon addition, which redissolved rapidly.
- 4) The stock were stored at room temperature.



5. MgSO<sub>4</sub> stock solution

Composition: 1 M MgSO<sub>4</sub>

For 100 ml of 1M MgSO<sub>4</sub> stock solution:

- 1) The MgSO<sub>4</sub>.7H<sub>2</sub>O powder was weighed to 24.65 g and dissolved in 100 ml distilled water.
- 2) The solution was filtered using a sterile 0.2 µm nylon syringe filter and kept in a sterile vial.
- 3) The solution was stored in room temperature prior to use.

6. LBA-5052 medium

- 1) Contains glucose to suppress expression while growing to high density, lactose to induce expression when glucose and glycerol exhausted. NB metals should be added before addition of NPS to avoid precipitation.
- 2) The following ingredients were dissolved in ~ 928 ml LBA:

<b>Component</b>	<b>1 liter</b>
LBA	~ 928 ml
1 M MgSO <sub>4</sub>	1 ml
1000 X metals (use 0.1 x)	100 µl
50 x 5052	20 ml
20 x NPS	50 ml
Ampicillin (100 mg/ml)	1 ml

### **Appendix A3: Antibiotic and solutions for enzyme expression**

#### 1. Ampicillin stock solution

Composition: 100 mg/ml ampicillin.

For 10 ml ampicillin stock solution:

- 1) The ampicillin powder was weighed to 1 g and dissolved in 10 ml distilled water.
- 2) The solution was filtered using a sterile 0.2  $\mu\text{m}$  nylon syringe filter and kept in a sterile vial.
- 3) The solution was stored in  $-20\text{ }^{\circ}\text{C}$  prior to use.

#### 2. IPTG stock solution

Composition: 0.5 M IPTG

For 10 ml IPTG solution:

- 1) The IPTG powder was weighed to 1.19 g and dissolved in 100 ml distilled water.
- 2) The solution was filtered using a sterile 0.2  $\mu\text{m}$  nylon syringe filter and kept in a sterile vial.

The solution was stored in  $-20\text{ }^{\circ}\text{C}$  and thawed prior to use

## Appendix A4: Solutions for Molecular Works

- |    |                                |          |
|----|--------------------------------|----------|
| 1. | 50X TAE Electrophoresis Buffer | 2 L      |
|    | (2 M Tris, 50 mM EDTA)         |          |
|    | Tris base                      | 484 g    |
|    | Glacial acetic acid            | 114.2 ml |
|    | 0.5 M EDTA, pH 8.0             | 200 ml   |

To make 1X TAE 20 L, add 400 ml 50X Buffer into 19.6 L of distilled water.
  
- |    |               |        |
|----|---------------|--------|
| 2. | 10X TE Buffer | 100 ml |
|    | Tris base     | 100 mM |
|    | EDTA, pH 8.0  | 10 mM  |

To make 1X TE Buffer, dilute 1 ml of 10X TE Buffer with 9 ml of distilled water.
  
- |    |                 |         |
|----|-----------------|---------|
| 3. | Tris-Cl, pH 8.0 | 500 ml  |
|    | Tris            | 60.57 g |

Add 350 ml distilled water and adjust pH to 8.0 with concentrated HCl (approximately 21 ml, but start with less). Let it cool to room temperature and make final adjustment to the pH. Autoclavable.
  
- |    |                    |         |
|----|--------------------|---------|
| 4. | 0.5 M EDTA, pH 8.0 | 500 ml  |
|    | EDTA               | 93.05 g |

Dissolve in 350 ml distilled water. Place on a magnetic stirrer and stir vigorously. Adjust the pH to 8.0 by adding approximately 10 g NaOH pellets. The disodium salt of EDTA will not go into solution until the solution is adjusted to approximately pH 8.0. Bring to 500 ml total volume with distilled water. Filter and sterilize by autoclaving.

## Appendix A5: Working Solutions for SDS-PAGE

1.	Acrylamide Mix Stock Solution	100 ml
	30% (w/v) Acrylamide	
	0.8% (w/v) Bis-acrylamide	
2.	10% SDS	100 ml
	Sodium dodecyl sulfate	10 g
	Distilled water	100 ml
	Store at room temperature.	
3.	10% Ammonium persulfate	5 ml
	Ammonium persulfate	0.5 g
	Distilled water	5 ml
	Stable for months in a capped tube in refrigerator.	
4.	10% Separating Gel	5 ml
	Distilled water	1.9 ml
	30% Acrylamide mix	1.7 ml
	1.5 M Tris-HCl, pH 8.8	1.3 ml
	10% SDS	0.05 ml
	10% Ammonium persulfate	0.05 ml
	TEMED	0.002 ml

5.	5% Stacking Gel	5 ml
	Distilled water	3.4 ml
	30% Acrylamide mix	0.83 ml
	1 M Tris-HCl, pH 6.8	0.63 ml
	10% SDS	0.05 ml
	10% Ammonium persulfate	0.05 ml
	TEMED	0.005 ml

6.	Tris-glycine Electrophoresis Buffer	1 L
	Tris	3 g
	Glycine	14.4 g
	SDS	1 g

Use distilled water to make 1 liter solution. pH should be approximately 8.3.

7.	5X Sample Buffer	10 ml
	1 M Tris-HCl, pH 6.8	0.6 ml
	50% Glycerol	5 ml
	10% SDS	2 ml
	2-mercaptoethanol	0.5 ml
	1% Bromophenol blue	1 ml
	Distilled water	0.9 ml

Stable for weeks in the refrigerator or for months at -20°C.

8.	Staining Solution (Solution A)	1 L
	Methanol	450 ml
	Distilled water	450 ml
	Glacial acetic acid	100 ml
	Coomassie Blue R-250	1 g

9.	Destaining Solution (Solution B)	1 L
	Distilled water	800 ml
	Methanol	100 ml
	Glacial acetic acid	100 ml

### **Appendix A6: Working Solutions for Western Blot**

1.	Blocking Solution	20 ml
	5% skimmed milk	1 g
	Add with 1X TBST until 20 ml	

2.	10X TBS	1 L
	Tris-base	24.2 g
	NaCl	80 g
	Adjust to pH 7.5	

3.	1X TBSTT (Ab 1:2000)	
	1X TBS	200 ml
	Tween-20	100 $\mu$ l

## Appendix A7: pH Buffers

### 1. Glycine-HCl (pH 3)

0.2 M glycine 12.5 ml

0.2 M HCl 2.85 ml

Top up to 50 ml with ddH<sub>2</sub>O.

### 2. Sodium acetate (pH 4)

0.2 M acetic acid 164 ml

0.2 M sodium acetate 36 ml

Top up to 200 ml with ddH<sub>2</sub>O.

### 3. Sodium acetate (pH 5)

0.2 M acetic acid 59 ml

0.2 M sodium acetate 141 ml

Top up to 200 ml with ddH<sub>2</sub>O

### 4. Sorensen phosphate (pH 6)

0.2 M NaH<sub>2</sub>PO<sub>4</sub> 44 ml

0.2 M Na<sub>2</sub>HPO<sub>4</sub> 6.2 ml

Top up to 50 ml with ddH<sub>2</sub>O

### 5. Sorensen phosphate (pH 7)

0.2 M NaH<sub>2</sub>PO<sub>4</sub> 39 ml

0.2 M Na<sub>2</sub>HPO<sub>4</sub> 61 ml

Top up to 100 ml with ddH<sub>2</sub>O

6. Sorensen phosphate (pH 8)

0.2 M NaH <sub>2</sub> PO <sub>4</sub>	5.3 ml
0.2 M Na <sub>2</sub> HPO <sub>4</sub>	94.7 ml
Top up to 100 ml with ddH <sub>2</sub> O	

7. Glycine -NaOH (pH 9)

0.2 M NaOH	4.4 ml
0.2 M glycine	25 ml
Top up to 50 ml with ddH <sub>2</sub> O	

**Appendix A8: DNS Assay Buffer**

Dinitrosalicylic acid (DNS) solution	1 L
3,5-dinitrosalicylic acid	10 g
Sodium potassium tartrate tetrahydrate	300 g
2N NaOH	200 ml

Add 3,5-dinitrosalicylic to 500 ml of reagent grade water. Add slowly sodium potassium tartrate tetrahydrate and 2 N NaOH. Dilute to a final volume of 1 L with reagent grade water. Protect from carbon dioxide and store no longer than 2 weeks.



## **Appendix A9: Protein Purification buffer**

### 1. Buffer A

Composition: 20 mM sodium phosphate buffer (pH 7.4), 0.5 M NaCl, 30 mM imidazole

### 2. Buffer B

Composition: 20 mM sodium phosphate buffer (pH 7.4), 0.5 M NaCl, 300 mM imidazole

### 3. Stripping buffer

Composition: 20 mM sodium phosphate buffer (pH 7.4), 0.5 M NaCl, 50 mM EDTA

### 4. Phenylmethylsulfonyl fluoride (PMSF)

Composition: 10 mM PMSF.

For 10 ml 10 mM PMSF:

- 1) The PMSF powder was weighed to 17.4 mg and dissolved in 10 ml isopropanol
- 2) The solution was filtered using a sterile 0.2  $\mu$ m nylon syringe filter and kept in a sterile vial.
- 3) The solution was aliquoted of appropriate volume and stored in -20 °C prior to use.

Note: The half-life of a 20  $\mu$ M aqueous solution of PMSF is about 35 minutes at pH 8.

## APPENDIX B

### XYLOSE STANDARD

Plotting graph for xylose standard


- 1) Prepare 1 mg/ml xylose stock in 50 mM sodium acetate buffer (pH 5). Prepare different concentrations of xylose as stated below:

Concentration (mg/ml)	Volume of xylose( $\mu$ l)	Volume of buffer( $\mu$ l)
0	0	750
0.05	37.5	712.5
0.1	75	675
0.15	112.5	637.5
0.2	150	600
0.25	187.5	562.5
0.3	225	525
0.35	262.5	487.5
0.4	300	450
0.45	337.5	412.5
0.5	375	375

- 2) Run the standard xylose as DNS assay of xylanase in triplicate.
- 3) Record the readings of OD at 540 nm and convert the xylose concentration to  $\mu$ mol as stated below:

Xylose concentration (mg/ml)	Xylose in $\mu\text{mol}$	Readings of at OD 540
0.00	0.00	0.00
0.05	0.25	0.07
0.10	0.50	0.27
0.15	0.75	0.52
0.20	1.00	0.72
0.25	1.25	0.98
0.30	1.50	1.17
0.35	1.75	1.38
0.40	2.00	1.59
0.45	2.25	1.73
0.50	2.50	2.01

4) Plotting the graph of xylose ( $\mu\text{mol}$ ) vs OD at 540 nm.



5) Use the equation from the graph for calculation of xylanase enzyme activity as 1U of enzyme is defined as the amount of enzyme releasing 1  $\mu\text{mol}$  xylose per min.

## APPENDIX C

### BSA STANDARD

Plotting graph for xylose standard


- 1) Prepare 2.5 µg/ml BSA stock in 50 mM sodium acetate buffer (pH 5.0) and prepare a set concentration of BSA as stated below:

BSA concentration (µg/ml)	Volume of BSA (µl)	Volume of buffer (µl)
0.001	0.2	99.8
0.002	0.4	99.6
0.003	0.6	99.4
0.004	0.8	99.2
0.005	1.0	99.0
0.006	1.2	98.8
0.007	1.4	98.6
0.008	1.6	98.4
0.009	1.8	98.2
0.01	2.0	98.0

- 2) 100 µl BSA at varied concentration is mixed with 100 µl Bradford reagent at RT in the 96-flat bottom plate. The mixture is equally mixed by pipetting. The triplicate mixture is prepared for each of BSA concentration.
- 3) The colorimetric readings are measured by multiwell plate reader immediately at 595 nm absorbance.
- 4) Record the readings as stated in the table below:

BSA concentration ( $\mu\text{g/ml}$ )	OD readings at 595 nm
0.001	0.0983
0.002	0.1190
0.003	0.1537
0.004	0.1770
0.005	0.1830
0.006	0.1850
0.007	0.1985
0.008	0.2160
0.009	0.2230
0.01	0.2447

5) Plotting the graph of BSA concentration ( $\mu\text{g/ml}$ ) vs OD at 595 nm.




6) Use the equation from the graph for calculation of concentration of xylanase

## APPENDIX D

### KINETIC PARAMETER MICHAELIS-MENTEN

The derivation of the Michaelis–Menten equation assumes that a slow, product forming reaction follows the rapid, reversible formation of an enzyme–substrate complex:



Where, E is the enzyme, S is the substrate and P is the product. The Michaelis–Menten equation is then derived by using the steady-state approximation for the ES complex: specifically the concentration of the enzyme–substrate complex is assumed to change much more slowly than the concentration of the substrate, so the rate equation takes the form.

$$\frac{d[P]}{dt} = \frac{V_{\max}[S]}{K_m + [S]}, \quad \text{Equation E12}$$


Where,  $K_m$  is the Michaelis constant and  $V_{\max}$  is the maximum velocity of the reaction achieved when the enzyme active sites in the sample are all complexed with substrate all the time, and [P] is the concentration of product at any given time during the time course. The relationship between the  $K_m$  and the unitary rate constants in the reaction scheme is:

$$K_m = \frac{k_{-1} + k_2}{k_1} \quad \text{Equation E13}$$

Taking the reciprocal of both sides of the Michaelis–Menten equation gives the Lineweaver–Burk equation that is often used to graphically analyse enzyme kinetic data. The equation is:


$$\frac{1}{V} = \frac{K_m}{V_{\max}[S]} + \frac{1}{V_{\max}}. \quad \text{Equation E14}$$

This relationship was used to estimate  $V_{\max}$  and  $K_m$  values.



**Figure D1:** Plot of rate of reaction versus the substrate concentration.

The Michaelis-Menten constant ( $K_m$ ) and maximum velocity of substrate hydrolysis ( $V_{\max}$ ) were determined from the Lineweaver-Burk plot



**Figure D2:** The Lineweaver-Burk plot and kinetic parameters determination.



## LIST OF PUBLICATIONS AND PAPER PRESENTED

1. **Damis, S.I.R.**, Abd. Murad, A.M., Abu Bakar, F.D., Rashid, S.A., Jaafar, N.R., and Illias, R.M. (2019). Protein engineering of GH11 xylanase from *Aspergillus fumigatus* RT-1 for catalytic efficiency improvement on kenaf biomass hydrolysis. *Enzyme and Microbial Technology*. 131 [109383].
2. **Damis, S.I.R.**, Illias, R.M. (2016). Substrate-binding site recognition of family 11 xylanase from *Aspergillus fumigatus* by molecular docking. *International Conference on Bioinformatics and Computational Biology*. Feb 2-3, 2016. Kuala Lumpur.
3. Noor, Y.M., Samsulrizal, N.H., Jema'on, N.A., Low, K.O., Ramli, A.N., Alias, N.I., **Damis, S.I.R.**, Fuzi, S.F., Isa, M.N., Murad, A.M., Raih, M.F., Bakar, F.D., Najimudin, N., Mahadi, N.M., and Illias RM. (2014). A comparative genomic analysis of the alkalitolerant soil bacterium *Bacillus lehensis* G1. *Gene*. 545(2):253-61.