

BEE-BASED SENSOR USING *APIS MELLIFERA* FOR DETECTION OF
ANDROGRAPHIS PANICULATA VOLATILE COMPOUNDS

KERK WEN CHIANN

A thesis submitted in fulfilment of the
requirement for the award of the degree of
Doctor of Philosophy (*Bioprocess Engineering*)

School of Chemical and Energy Engineering
Faculty of Engineering
Universiti Teknologi Malaysia

DECEMBER 2019

ACKNOWLEDGEMENT

In preparing this thesis, I was in contact with many people, researchers, academicians, and practitioners. They have contributed towards my understanding and thoughts. I wish to express my sincere appreciation to my main thesis supervisor, Associate Professor Dr. Chua Lee Suan, for encouragement, guidance, critics and friendship.

My fellow postgraduate students should also be recognised for their support. My sincere appreciation also extends to all my colleagues and others who have aided at various occasions. Their views and tips are useful indeed. Unfortunately, it is not possible to list all of them in this limited space.

I am grateful to all my family members especially my father, mother and husband who never give up supporting me from the first day of my study until now.

ABSTRACT

Honey bees can be trained to be sniffer bees due to the presence of high number of odorant receptors (170 odorant receptors) in their heads. This unique characteristic enables them to detect scent down to part per trillion level. In this study, localized honey bees (*Apis mellifera*) were trained by using the method of classical Pavlovian conditioning. The proboscis extension reflex (PER) of sniffer bees towards the target odor was observed and recorded. The phytochemical profile of *Andrographis paniculata* (*A. paniculata*) was constructed using headspace solid-phase microextraction coupled with gas chromatography integrated with mass spectrometer method. The volatile marker compounds were identified. The sniffing capacity of the sniffer bees was determined by varying the heating temperature from 50-120 °C, the weight of plant material from 20-100 mg and the percentage (20-100%) of the target herbal sample in the mixture of *A. paniculata* and *Clinacanthus nutans* (*C. nutans*). *C. nutans* is an herbal plant which is morphologically similar to *A. paniculata* and it also belongs to the Acanthaceae family. The efficiency, accuracy and sensitivity of sniffer bees were analyzed and validated statistically. The success rate of sniffer bees for heating temperatures was approximately 90 %. The success rate for minimum weight of plant sample, 20 mg was 50 %. The success rate percentage of target herbal sample increased when the percentage of *A. paniculata* was proportionally increased. Compounds such as caryophyllene, β -elemene, 3,3-dimethylhexane, apiol, 6,10,14-trimethyl-2-pentadecanone and dihydroactinidiolide were detected in the gaseous mixture. The kinetics of volatile marker compounds released from the plant samples were studied to predict the concentrations of the volatile marker compounds for sniffer bee detection at 85 °C. Second-order and two-site kinetic models were selected because of the kinetic data of these volatile marker compounds fitted well to these models ($R^2 > 0.9$).

ABSTRAK

Lebah madu boleh dilatih menjadi lebah pengesan kerana kehadiran reseptor bau yang banyak (170 reseptor bau) dalam kepala mereka. Ciri unik ini membolehkan mereka mengesan bau sampai ke tahap part per trillion. Dalam kajian ini, lebah madu (*Apis mellifera*) telah dilatih dengan kaedah classical Pavlovian conditioning. Tindakbalas pengeluaran lidah (PER) lebah pengesan terhadap bau sasaran telah diperhatikan dan dicatatkan. Profil fitokimia hempedu bumi (*A. paniculata*) telah dibangunkan melalui kaedah pengekstrakan mikro fasa pepejal ruang tutupan yang digandingkan dengan kromatografi gas dan spektrometer jisim. Sebatian-sebatian penanda yang mudah meruap telah dikenalpasti. Keupayaan lebah pengesan untuk mengesan bau telah ditentukan dengan mengubah suhu pemanasan dari 50 °C ke 120 °C, jisim bahan tumbuhan dari 20-100 mg dan peratusan (20-100 %) sampel herba sasaran dalam campuran hempedu bumi (*A. paniculata*) dan belalai gajah (*C. nutans*). Belalai gajah merupakan tumbuhan herba yang mempunyai kesamaan dengan hempedu bumi dari segi morfologi dan ia juga digolongkan kepada keluarga Acanthaceae. Kecekapan, ketepatan dan kepekaan lebah pengesan telah dianalisa dan disahkan secara statistik. Kadar kejayaan untuk pelbagai suhu pemanasan adalah sekitar 90 %. Kadar kejayaan bagi jisim minimum tumbuhan 20 mg adalah sekitar 50 %. Peratusan kadar kejayaan untuk mengesan sampel herba sasaran meningkat apabila peratusan hempedu bumi dalam campuran meningkat secara berkadar. Cariofillene, β -elemene, 3,3-dimetilheksana, apiol, 6,10,14-trimetil-2-pentadekanona dan dihidroaktinidiolida adalah sebatian yang dikesan dalam campuran gas. Kinetik sebatian-sebatian penanda mudah meruap yang dikeluarkan daripada sampel tumbuhan telah dikaji untuk meramal kepekatan sebatian-sebatian penanda yang mudah meruap bagi pengesanan lebah pengesan pada 85 °C. Model kinetic tertib kedua dan model dwi tapak telah dipilih kerana data kinetik dari sebatian-sebatian penanda yang mudah meruap bersesuaian dengan model-model tersebut ($R^2 > 0.9$).

TABLE OF CONTENTS

	TITLE	PAGE
	DECLARATION	iii
	DEDICATION	iv
	ACKNOWLEDGEMENT	v
	ABSTRACT	vi
	ABSTRAK	vii
	TABLE OF CONTENTS	ix
	LIST OF TABLES	xii
	LIST OF FIGURES	xv
	LIST OF SYMBOLS	xvi
	LIST OF ABBREVIATIONS	xvii
CHAPTER 1	INTRODUCTION	1
1.1	Research Background	1
1.2	Research Problem	3
1.3	Research Objectives	4
1.4	Research Significance	5
1.5	Research Scopes	6
CHAPTER 2	LITERATURE REVIEW	7
2.1	Olfactory and Learning in Honey Bees (<i>Apis mellifera</i>)	7
2.2	Food Foraging Behavior of Worker Bees	10
2.2.1	Influence of Floral Odors to Honey Bees in Food Search	12
2.3	Recognition of Worker Bee in Beehive	13
2.4	Rational of Honey Bee as Sniffer Bee	14
2.4.1	Mushroom Bodies for Olfactory Learning	14
2.4.2	Odorant Receptors in Antennae	15

2.4.3	Proboscis Extension for Food	17
2.5	Advantages of Sniffer Bee	17
2.6	Limitation of Sniffer Bee	19
2.7	Classical Conditioning for Bee Training	19
2.8	Simulation of Proboscis Extension Reflex (PER)	21
2.9	Factors Affecting Success Rate of Learning in Honey Bees	25
2.9.1	Concentration of Odorant	25
2.9.2	Sampling Time	26
2.9.3	Heating Temperature for Sample	26
2.10	Insect Based Sniffers	27
2.11	Utilization of Sniffer Bee	29
2.12	Herbal Plant Detection by Sniffer Bee	30
2.13	HS-SPME/GC-MS Analysis of Volatile Compounds from <i>Andrographis paniculata</i>	33
2.14	Other Methods for Herbal Plant Identification	38
2.15	Kinetic Modelling of Volatile Marker Compounds Emission	41
2.16	Future Perspective of Sniffer Bees for Herbal Plant Detection	45
CHAPTER 3	MATERIAL AND METHODOLOGY	49
3.1	Introduction to Overall Study	49
3.2	Plant Materials and Honey Bees	52
3.3	Collection and Storage of Honey Bees	52
3.4	Bee Training by Classical Pavlovian Conditioning	53
3.5	Thermal Treatment on Andrographolide	55
3.6	Exposure to Vapor Released from Thermal-treated <i>A. paniculata</i>	55
3.7	Validation of Bee Sniffing Capacity	55
3.7.1	Efficiency of Sniffer Bees	56
3.7.2	Selectivity of Sniffer Bees	57
3.7.3	Sensitivity of Sniffer Bees	57

3.8	Headspace Solid Phase Micro-extraction Coupled with Gas Chromatography-Mass Spectrometry (HS-SPME/GC-MS)	58
3.9	Kinetic Model Development	59
3.10	Validation of the Kinetic Models	60
CHAPTER 4	RESULTS AND DISCUSSION	63
4.1	Sniffer Bees for <i>Andrographis paniculata</i> Detection	63
4.2	Effects of Heat-treated <i>Andrographis paniculata</i> on Sniffing Success Rate	66
4.3	Sensitivity of Sniffer Bees	68
4.4	Selectivity of Sniffer Bees	69
4.5	Persistency of Sniffer Bees	71
4.6	Volatile Phytochemicals Extracted by HS-SPME	73
4.7	Volatile Marker Compounds in <i>Andrographis paniculata</i>	78
4.8	Effect of Heating Duration on Phytochemical Profile	81
4.9	Kinetics of Volatile Marker Compounds Released Under Thermal Treatment	84
CHAPTER 5	CONCLUSION AND RECOMMENDATION	91
5.1	Conclusion	91
5.2	Recommendation	93
REFERENCES		95
APPENDIX A		113

LIST OF TABLES

TABLE NO.	TITLE	PAGE
Table 2.1	Traditional uses of <i>A. paniculata</i>	32
Table 2.2	GC-MS analysis of the components in ethanolic extracts of <i>Andrographis paniculata</i>	36
Table 3.1	Equations for kinetic study of volatile marker compounds	59
Table 4.1	Detected compounds in heated <i>Andrographis paniculata</i>	73
Table 4.2	Equations for kinetic study of volatile marker compounds	82
Table 4.3	Kinetics of volatile marker compounds	83
Table 4.4	<i>h</i> values for volatile marker compounds	84
Table 4.5	Parameters values for volatile marker compounds	85
Table 4.6	R ² and RMSE values of volatile marker compounds to kinetic models	86

LIST OF FIGURES

FIGURE NO.	TITLE	PAGE
Figure 2.1	Honey bee (<i>Apis mellifera</i>)	11
Figure 2.2	Stinger of honey bee	15
Figure 2.3	Restrained honey bee	22
Figure 2.4	Bee holder designed by Inscentinel Ltd	22
Figure 2.5	Classical conditioning of honey bee	25
Figure 2.6	<i>Andrographis paniculata</i>	33
Figure 2.7	GC-MS chromatogram for ethanolic extract of <i>Andrographis paniculata</i>	38
Figure 2.8	Andrographolide	38
Figure 3.1	Flow chart of the study	50
Figure 3.2	Transparent plastic box	53
Figure 3.3	Honey bee in bee holder	53
Figure 3.4	Classical Pavlovian conditioning training for honey bees	54
Figure 4.1	Efficiency of sniffer bees for <i>Andrographis paniculata</i> detection at 85°C	63
Figure 4.2	Sniffing capacity of sniffer bees at different sample heating temperatures	65
Figure 4.3	Chromatograms of <i>Andrographis paniculata</i> at different heating times	66
Figure 4.4	Success rate of sniffer bee production at different sample masses	67
Figure 4.5	Success rate of sniffer bees responding to the scent of <i>Andrographis paniculata</i> in different percentages of a mixture	69
Figure 4.6	Success rate of the same batch of sniffer bees in sniffing <i>Andrographis paniculata</i> for consecutive 5 days	70

Figure 4.7	Relationship of β -elemene (\blacklozenge); 3,3-dimethylhexane (\blacksquare); caryophyllene (\blacktriangle); 5,6,7,7a-tetrahydro-4,4,7a-trimethyl-2(4H)- benzofuranone (\times); apiol ($+$) and 6,10,14-trimethyl-2-pentadecanone (\bullet) concentration with sample heating temperature	77
Figure 4.8	Chromatograms of <i>Andrographis paniculata</i> (a) and <i>Clinacanthus nutans</i> (b)	78
Figure 4.9	Chromatograms of <i>Andrographis paniculata</i> at different heating times	80
Figure 4.10	Relationship of 5,6,7,7 α -tetrahydro-4,4,7 α -trimethyl-2(4H)- benzofuranone (\blacklozenge); apiol (\blacksquare); 6,10,14-trimethyl-2-pentadecanone (\blacktriangle); caryophyllene (\bullet); 3,3-dimethylhexane (\times) and β -elemene (\times) contents with heating duration	81
Figure 4.11	Kinetics of dihydroactinidiolide (\bullet), caryophyllene (\blacksquare) and β -elemene (\blacklozenge) where blue lines are predicted values by second-order model	84
Figure 4.12	Kinetics of 6,10,14-trimethyl-2-pentadecanone (\bullet), apiol (\blacksquare) and 3,3-dimethylhexane (\blacktriangle) where blue lines are predicted values by second-order model	85

LIST OF SYMBOLS

n	-	Number of honey bees positively respond to target odorant
N	-	Total number of honey bees in a training
N_a	-	Number of sniffer bees correctly recognize target odorant
N_A	-	Total number of sniffer bees in a training
N_t	-	Number of sniffer bees positively respond to target odorant
N_T	-	Total number of sniffer bees in a training

LIST OF ABBREVIATIONS

AL	-	Antennal lobe
CR	-	Conditioned response
CS	-	Conditioned stimulus
eLTM	-	Early long-term memory
GC	-	Gas chromatography
HPLC	-	High performance liquid chromatography
HS	-	Headspace
KCs	-	Kenyon cells
LH	-	Lateral horn
lLTM	-	Late long-term memory
LN _s	-	Local interneurons
LTM	-	Long-term memory
MB _s	-	Mushroom bodies
MS	-	Mass spectrometer
MTM	-	Mid-term memory
OBPs	-	Odorant-binding proteins
OR _s	-	Olfactory receptors
ORCs	-	Olfactory receptor cells
ORN _s	-	Olfactory receptor neurons
OSN _s	-	Olfactory sensory neurons
PBPs	-	Pheromone binding proteins
PER	-	Proboscis Extension Reflex
PN _s	-	Projection neurons
R ²	-	Coefficient of determination
RMSE	-	Root mean square error

SPME	-	Solid phase micro-extraction
STM	-	Short-term memory
TB	-	Tuberculosis
TNT	-	Trinitrotoluene
UR	-	Unconditioned response
US	-	Unconditioned stimulus

CHAPTER 1

INTRODUCTION

1.1 Research Background

Honey bee possesses several key criteria to be an excellent sniffer. Honey bees (*Apis mellifera*) are subset of bees in the genus *Apis*. Honey bees have 170 odorant receptors and a pair of mushroom bodies (Robertson and Wanner, 2006, Farris et al., 1999). Therefore, they can differentiate odors better than fruit flies and mosquitoes which only have 62 and 79 odor receptors respectively (Clyne et al., 1999, Vosshall et al., 1999, Hill et al., 2002). Besides their outstanding ability to distinguish smells, their sensitivity to odors is also excellent. Honey bees can be trained to detect scents down to parts per trillion level (Bromenshenk et al., 2003). The mushroom bodies assist honey bees to keep their memory about the scent, and thus they are able to remember the scent which can last for several days and even for their entire life (Menzel, 1999, Schröter and Menzel, 2003). The life spans of honey bees are different; workers live 3-6 weeks during spring and summer, but about 4 months during winter. For bee queen, her lifespan is around 2-3 years, but less than 1 year in commercial hives. There is not much information about the life span of drones (Page and Peng, 2001).

Due to their extraordinary abilities, honey bees can be used to replace sniffer dogs. Defense Advanced Research Projects Agency (DARPA), an agency of United States Department of Defense had funded a project in 2004 by utilizing honey bees for explosives detection. It was found that they could detect trinitrotoluene (TNT) at the

level of part per trillion (ppt). Besides that, Inscentinel Ltd., a company which specializes in the development of insect olfaction technologies for the detection of trace chemicals discovered that honey bees were able to detect 2,4-dinitrotoluene (DNT) down to at least 78 ppt. In this case, honey bees are likely to be superior than sniffer dogs. In 2008, the company has invented Vasor136 (biosensor) which can house 36 bees to detect 6 different chemicals simultaneously. According to Suckling and Sagar (2011), honey bees could be trained to detect the scent of *Mycobacterium tuberculosis* through human breath. A person who is suffering from Tuberculosis (TB) can be determined in a faster and more sensitive manner using sniffer bees. In agricultural industry, honey bees were also used to detect the presence of Mediterranean fruit fly (*Ceratitiscapitata Wiedemann*) larvae in Valencia oranges at early stage of maturity (Chamberlain et al., 2012). Behavioral studies have demonstrated that bees are able to recognize learned odors within <200ms (Krofczik et al., 2009). This characteristic makes them manage to response fast and immediately while exposed to target odorant. Therefore, sniffer bees have been getting actively applied in medical, security and agricultural industries for the last few years. This is the first study to report the use of sniffer bees for herbal plant recognition.

In this research, honey bees are trained by classical Pavlovian conditioning method to detect *Andrographis paniculata* (Hempedu Bumi). *A. paniculata* is an erect herb that widely distributed in South East Asia. The whole plant is traditionally used as an anti-inflammatory and antipyretic folk medicine for the treatment of fever, cold, laryngitis, diarrhoea, and inflammation (Sheeja et al., 2006, Suebsasana et al., 2009). The extract of *A.paniculata* and its major *ent*-labdane diterpenoids have been shown to display antiviral (Wiart et al., 2005), bacteriostatic (Mishra et al., 2009), immunostimulatory (Ajaya Kumar et al., 2004), as well as hepatoprotective and hepatostimulating activities (Kapil et al., 1993) in scientific studies.

In the training of honey bees, their antenna will be touched by cotton bud dipped with sucrose solution when the bees are exposed to the odor of thermally treated *A. paniculata*. The bees will show Proboscis Extension Reflex (PER), which is sticking out their tongues and sugar solution is given to them immediately as a reward.

By repeating this procedure for several times, honey bees will associate the odor of *A. paniculata* with sugar reward. They will automatically stick out their tongues whenever they are exposed to the scent of *A. paniculata* even after a couple of days after training. Honey bees which are successfully trained (sniffer bees) can be further studied and developed into bee-sensor for the application in herbal and food industries in the future.

The kinetics of volatile marker compounds was studied to estimate the shortest time for sniffer bees to detect the volatile marker compounds during the heating process of *A. paniculata* at the optimum temperature. This kinetic model is very important to relate the concentration of volatile marker compounds to the Proboscis Extension Reflex (PER) of bees for better performance of sniffer bees in carrying out their sniffing task.

1.2 Research Problem

The identification of *A. paniculata* is usually carried out by solvent extraction and followed by high performance liquid chromatographic analysis. This conventional technique is time-consuming and environmental unfriendly since lot of solvent is required for plant extraction. The detection of *A. paniculata* based on gas phase sample is relatively seldom studied by researchers. This could be the difficulty in gas sample collection and data reproducibility. The plant sample is heated to generate volatile compounds which are then collected for gas chromatography injection. Indeed, this gas phase technique is simpler, cost effective and time-saving if sample lost could be reduced to a minimum level. Nevertheless, this gas chromatography method must be validated before use. *Andrographis paniculata* is becoming popular due to its significant medical value. It is manufactured into pharmaceutical products for medical purpose. Therefore, the purity of raw material becomes the largest concern when

comes to the formulation of medical products and a rapid screening of the raw material is a must to prevent adulteration.

Honey bees have been proven by many researchers to be excellent sniffer bees. The performance of sniffer bees is likely to be comparable or better than sniffer dogs (Gazit and Terkel, 2003, Bromenshenk et al., 2003, Angle et al., 2016). It was found that honey bees can be trained to detect explosives, drugs, landmines, infestation of food by insects and disease through the breath of Tuberculosis (TB) patients. However, there is still no report showing its usage and effectiveness in sniffing herbal plants. Therefore, the sensitivity and accuracy of sniffing capacity of sniffer bees to recognize *A. paniculata* is not yet explored and quantified.

1.3 Research Objectives

This research is conducted to determine the sniffing capacity of sniffer bees for *A. paniculata* detection. Honey bees are trained by Pavlovian conditioning method to detect and recognize the volatile marker compounds released from thermally-treated *A. paniculata*. The reliability, sensitivity and persistency of sniffer bees are validated to prove that the utilization of sniffer bees in herbal plant detection is feasible and effective.

There are three objectives for this study. The objectives are:

- i. To determine the sniffing capacity of honey bees in terms of reliability, sensitivity and persistency for the detection of volatile marker compounds from thermal-treated *A. paniculata* (Hempedu Bumi).
- ii. To construct the volatile phytochemical profile of *A. paniculata*.
- iii. To determine the kinetics of the release of key volatile marker compounds from thermally-treated *A. paniculata*.

1.4 Research Scopes

- i. To determine the efficiency, accuracy and sensitivity limit of sniffer bees for *A. paniculata* detection by introducing other herb in the same family of Acanthaceae, namely *Clinacanthus nutans* (Sabah snake grass) as sample adulterant.
- ii. To validate the persistency characteristic of sniffer bees by keeping the same batch of sniffer bees for 5 days.
- iii. To construct the volatile phytochemical profile of *A. paniculata* at heating temperature varied from 50-120°C and detected by gas chromatography integrated with mass spectrometer.
- iv. To identify the relationship of volatile marker compounds concentration and heating temperature with the performance of sniffer bees.
- v. To determine the kinetics of the release of volatile marker compounds and predict the shortest sample heating time for sufficient amount of volatile marker compounds release at the optimized heating temperature (85°C).

1.5 Research Significance

The prominent significance of the study is the use of honey bees to perform task that is difficult and less accurate to be carried out by mankind. The sniffing

capacity of honey bees can be scientifically proven for the detection or differentiation of herbal plant. This non-destructive exploration uses honey bees as biological detector to detect the target odor. The sensitivity capacity of *A. mellifera* could be established for detection of *A. paniculata*.

This research generated fundamental data for the development of bee-based sensor. This sensor is very useful for herbal and/or food industries for quality control purposes. When raw herbal materials are used for research purpose, bee-based sensor would be a great tool to ensure the quality of raw herbal materials can be ensured so that unnecessary errors can be prevented during researches. The advantages of bee-based sensor are due to its unique characteristics such as simple operation, time-saving and cost effective. This is because a bee hive usually contains up to 30,000 to 70,000 of worker bees (Davidson et al., 2016) and a bee hive usually costs around RM 2000, which means that lot of sniffer bees could be produced. This is potentially cost effective compared to the cost of an analytical instrument with its yearly maintenance expenses usually up to several thousand Ringgit Malaysia. Furthermore, laboratory testing using gas chromatography and/or liquid chromatography require time consuming sample treatment protocols before injection. Analytical technique using liquid chromatography may require hours for *A. paniculata* determination and confirmation.

Previous studies on *A. paniculata* were mostly focused on non-volatile compounds, especially andrographolide, and then detected by high performance liquid chromatography. This study suggests another reliable alternative for *A. paniculata* detection based on its phytochemical profile constructed by volatile compounds. Therefore, the volatile phytochemical profile of *A. paniculata* could be used for quality assurance of herbal related industries.

REFERENCES

- A. Rescorla, R. & Wagner, A. (1972). *A Theory of Pavlovian Conditioning: The Effectiveness of Reinforcement and Non-Reinforcement*.
- Ábalos, M., Bayona, J. M. & Pawliszyn, J. (2000). Development of a Headspace Solid-Phase Microextraction Procedure for the Determination of Free Volatile Fatty Acids in Waste Waters. *Journal of Chromatography A*, 873, 107-115.
- Ache, B. W. & Young, J. M. (2005). Olfaction: Diverse Species, Conserved Principles. *Neuron*, 48, 417-430.
- Ai, H. (2013). Sensors and Sensory Processing for Airborne Vibrations in Silk Moths and Honeybees. *Sensors*, 13, 9344.
- Ajaya Kumar, R., Sridevi, K., Vijaya Kumar, N., Nanduri, S. & Rajagopal, S. (2004). Anticancer and Immunostimulatory Compounds from *Andrographis paniculata*. *Journal Of Ethnopharmacology*, 92, 291-295.
- Akbar, S. (2011). *Andrographis paniculata*: A Review of Pharmacological Activities and Clinical Effects. *Alternative Medicine Review* 6, 16, 66-77.
- Alqarni, A. (2006). *Tolerance of Summer Temperature in Imported and Indigenous Honeybee Apis mellifera L. Races in Central Saudi Arabia*.
- Angle, C., Waggoner, L. P., Ferrando, A., Haney, P. & Passler, T. (2016). Canine Detection of the Volatilome: A Review of Implications for Pathogen and Disease Detection. *Frontiers in Veterinary Science*, 3, 47.
- Anton, S. & Homberg, U. (1999). Antennal Lobe Structure. *In*: Hansson, B. S. (ed.) *Insect Olfaction*. Berlin, Heidelberg: Springer Berlin Heidelberg.
- Arora, G., Cormier, F. & Lee, B. (1995). Analysis of Odor-Active Volatiles in Cheddar Cheese Headspace by Multidimensional GC/MS/Sniffing. *Journal of Agricultural and Food Chemistry*, 43, 748-752.
- Atmowidjojo, A. H., Wheeler, D. E., Erickson, E. H. & Cohen, A. C. (1997). Temperature Tolerance and Water Balance in Feral and Domestic Honey Bees, *Apis mellifera L. Comparative Biochemistry and Physiology Part A: Physiology*, 118, 1399-1403.

- Bargmann, C. I. (2006). Comparative Chemosensation from Receptors to Ecology. *Nature*, 444, 295.
- Barker, R. J. & Lehner, Y. (1974). Acceptance and Sustenance Value of Naturally Occurring Sugars Fed to Newly Emerged Adult Workers of Honey Bees (*Apis mellifera* L.). *Journal Of Experimental Zoology*, 187, 277-285.
- Barron, A., Schulz, D. & Robinson, G. (2002). Octopamine Modulates Responsiveness to Foraging-Related Stimuli in Honey Bees (*Apis mellifera*). *Journal of Comparative Physiology A*, 188, 603-610.
- Bhagavan, S. & Smith, B. H. (1997). Olfactory Conditioning in the Honey Bee, *Apis mellifera*: Effects of Odor Intensity. *Physiology & Behavior*, 61, 107-117.
- Bhatnagar, S., Santapau, H., Desa, J., Maniar, A., Ghadially, N., Solomon, M., Yellore, S. & Rao, T. (1961). Biological Activity of Indian Medicinal Plants. Part I. Antibacterial, Anti-tubercular and Antifungal Action. *Indian Journal of Medical Research*, 49, 799-813.
- Bicchi, C., Drigo, S. & Rubiolo, P. (2000). Influence of Fibre Coating in Headspace Solid-Phase Microextraction–Gas Chromatographic Analysis of Aromatic And Medicinal Plants. *Journal of Chromatography A*, 892, 469-485.
- Bilukha, O. O., Laurence, H., Danee, L., Subedi, K. P. & Becknell, K. (2011). Injuries and Deaths Due to Victim-Activated Improvised Explosive Devices, Landmines and Other Explosive Remnants of War in Nepal. *Injury Prevention*.
- Bitterman, M., Menzel, R., Fietz, A. & Schäfer, S. (1983). Classical Conditioning of Proboscis Extension in Honeybees (*Apis mellifera*). *Journal of Comparative Psychology*, 97, 107.
- Bitterman, M. E. (1996). Comparative Analysis of Learning in Honeybees. *Animal Learning & Behavior*, 24, 123-141.
- Blazyte-Cereskiene, L., Vaitkevičienė, G., Venskutonytė, S. & Buda, V. (2010). *Honey Bee Foraging in Spring Oilseed Rape Crops Under High Ambient Temperature Conditions*.
- Brattoli, M., De Gennaro, G., De Pinto, V., Demarinis Loiotile, A., Lovascio, S. & Penza, M. (2011). Odour Detection Methods: Olfactometry and Chemical Sensors. *Sensors*, 11, 5290.
- Braun, G. & Bicker, G. (1992). Habituation of an Appetitive Reflex in the Honeybee. *Journal of Neurophysiology*, 67, 588-598.

- Breer, H. (1997). Molecular Mechanisms of Pheromone Reception in Insect Antennae. *In: Cardé, R. T. & Minks, A. K. (eds.) Insect Pheromone Research: New Directions*. Boston, MA: Springer US.
- Breer, H., Boekhoff, I. & Tareilus, E. (1990). Rapid Kinetics of Second Messenger Formation in Olfactory Transduction. *Nature*, 345, 65-68.
- Brill, M. F., Rosenbaum, T., Reus, I., Kleineidam, C. J., Nawrot, M. P. & Rössler, W. (2013). Parallel Processing via a Dual Olfactory Pathway in the Honeybee. *The Journal of Neuroscience*, 33, 2443-2456.
- Bromenshenk, J. J., Henderson, C. B., Seccomb, R. A., Rice, S. D., Etter, R. T., Bender, S. F., Rodacy, P. J., Shaw, J. A., Seldomridge, N. L. & Spangler, L. H. (2003). Can Honey Bees Assist in Area Reduction and Landmine Detection? *J. Mine Action*, 7.
- Buchmann, S. (2010). *Honey Bees: Letters from the Hive*, Random House Children's Books.
- Buck, L. & Axel, R. (1991). A Novel Multigene Family May Encode Odorant Receptors: A Molecular Basis for Odor Recognition. *Cell*, 65, 175-187.
- Burdock, G. A. (2016). *Fenaroli's Handbook of Flavor Ingredients, Sixth Edition*, CRC Press.
- Calogirou, A., Larsen, B. R. & Kotzias, D. (1999). Gas-Phase Terpene Oxidation Products: A Review. *Atmospheric Environment*, 33, 1423-1439.
- Camhi, J. & Johnson, E. (1999). High-Frequency Steering Maneuvers Mediated by Tactile Cues: Antennal Wall-Following in the Cockroach. *Journal of Experimental Biology*, 202, 631-643.
- Campbell, A. L., Naik, R. R., Sowards, L. & Stone, M. O. (2002). Biological Infrared Imaging and Sensing. *Micron*, 33, 211-225.
- Carrillo, J. D., Salazar, C., Moreta, C. & Tena, M. T. (2007). Determination of Phthalates in Wine by Headspace Solid-Phase Microextraction Followed by Gas Chromatography–Mass Spectrometry: Fibre Comparison and Selection. *Journal of Chromatography A*, 1164, 248-261.
- Chamberlain, K., Briens, M., Jacobs, J. H., Clark, S. J. & Pickett, J. A. (2012). Use of Honey Bees (*Apis mellifera L.*) to Detect the Presence of Mediterranean Fruit Fly (*Ceratitidis Capitata Wiedemann*) Larvae in Valencia Oranges. *Journal of the Science of Food and Agriculture*, 92, 2050-2054.

- Chang, H.-M., But, P. P. & Yao, S.-C. (1986). *Pharmacology and Applications of Chinese Materia Medica*, World Scientific.
- Chao, W.-W. & LIN, B.-F. (2010). Isolation and Identification of Bioactive Compounds in *Andrographis paniculata* (Chuanxinlian). *Chinese Medicine*, 5, 17.
- Chapman, R. B., E. (2017). *Chemoreception* [Online]. Encyclopædia Britannica, inc. Available: <https://www.britannica.com/science/chemoreception/Interaction-between-taste-and-smell> [Accessed May 25, 2017 2017].
- Chapman, R. F. (1998). *The insects: Structure and Function*, Cambridge University Press.
- Checmarev, G., Casales, M. R. & Yeannes, M. I. (2013). Analysis of Applicability of Peleg Model to the Cooking-Infusion of Mackerel (*Scomber Japonicus*) Slices. *Food Science and Technology*, 33, 685-689.
- Cheng, A.-X., Lou, Y.-G., Mao, Y.-B., Lu, S., Wang, L.-J. & Chen, X.-Y. (2007). Plant Terpenoids: Biosynthesis and Ecological Functions. *Journal of Integrative Plant Biology*, 49, 179-186.
- Chittka, L., Thomson, J. D. & Waser, N. M. (1999). Flower Constancy, Insect Psychology, and Plant Evolution. *Naturwissenschaften*, 86, 361-377.
- Clifford, M. R. & Riffell, J. A. (2013). Mixture and Odorant Processing in the Olfactory Systems of Insects: A Comparative Perspective. *Journal of Comparative Physiology A*, 199, 911-928.
- Clyne, P. J., Warr, C. G., Freeman, M. R., Lessing, D., Kim, J. & Carlson, J. R. (1999). A Novel Family of Divergent Seven-Transmembrane Proteins: Candidate Odorant Receptors in *Drosophila*. *Neuron*, 22, 327-338.
- Cooper, P. D., Schaffer, W. M. & Buchmann, S. L. (1985). Temperature Regulation of Honey Bees (*Apis mellifera*) Foraging in the Sonoran Desert. *Journal of Experimental Biology*, 114, 1-15.
- Corbet, S. A. (2003). Nectar Sugar Content: Estimating Standing Crop and Secretion Rate in the Field. *Apidologie*, 34, 1-10.
- Corzo, O. & Bracho, N. (2006). Application of Peleg Model to Study Mass Transfer During Osmotic Dehydration of Sardine Sheets. *Journal of Food Engineering*, 75, 535-541.
- Crini, G. & Badot, P. M. (2010). *Sorption Processes and Pollution: Conventional and Non-conventional Sorbents for Pollutant Removal from Wastewaters*, Presses Universitaires de Franche-Comté.

- Davidson, M., Davidson, J. & Books, M. C. (2016). *The GIANT (Yet Tiny) Book on Insects*.
- Davoli, E., Gangai, M. L., Morselli, L. & Tonelli, D. (2003). Characterisation of Odorants Emissions from Landfills by SPME and GC/MS. *Chemosphere*, 51, 357-368.
- Dobson, H. E. (1993). Floral Volatiles in Insect Biology. *Insect-Plant Interactions Vol. V.*, 47-81.
- Doong, R.-A. & Liao, P.-L. (2001). Determination of Organochlorine Pesticides and Their Metabolites in Soil Samples Using Headspace Solid-Phase Microextraction. *Journal of Chromatography A*, 918, 177-188.
- Dreller, C., Page Jr., R. E. & Fondrk, M. K. (1999). Regulation of Pollen Foraging in Honeybee Colonies: Effects of Young Brood, Stored Pollen, and Empty Space. *Behavioral Ecology and Sociobiology*, 45, 227-233.
- Dreller, C. & Tarpay, D. R. (2000). Perception of the Pollen Need by Foragers in a Honeybee Colony. *Animal Behaviour*, 59, 91-96.
- Dudai, Y. (2004). *The Neurobiology of Consolidations, Or, How Stable is the Engram?*
- Dudareva, N. & Pichersky, E. (2000). Biochemical and Molecular Genetic Aspects of Floral Scents. *Plant physiology*, 122, 627-634.
- Dudareva, N., Pichersky, E. & Gershenzon, J. (2004). Biochemistry of Plant Volatiles. *Plant Physiology*, 135, 1893-1902.
- Eisenhardt, D. (2014). Molecular Mechanisms Underlying Formation of Long-Term Reward Memories and Extinction Memories in the Honeybee (*Apis mellifera*). *Learning & Memory*, 21, 534-542.
- Erber, J., Masuhr, T. H. & Menzel, R. (1980). Localization of Short-Term Memory in the Brain of the Bee, *Apis mellifera*. *Physiological Entomology*, 5, 343-358.
- Ezquerro, Ó., Ortiz, G., Pons, B. & Tena, M. A. T. (2004). Determination of Benzene, Toluene, Ethylbenzene and Xylenes in Soils by Multiple Headspace Solid-Phase Microextraction. *Journal of Chromatography A*, 1035, 17-22.
- Fahrbach, S. E., Farris, S. M., Sullivan, J. P. & Robinson, G. (2003). Limits on Volume Changes in the Mushroom Bodies of the Honey Bee Brain. *Journal of neurobiology*, 57, 141-151.
- Farris, S. M., Robinson, G. E., Davis, R. L. & Fahrbach, S. E. (1999). Larval and Pupal Development of the Mushroom Bodies in the Honey Bee, *Apis mellifera*. *The Journal of Comparative Neurology*, 414, 97-113.

- Frisch, K. V. (1919). Über den Geruchsinn der Biene und seine blutenbiologische Bedeutung.
- Galizia, C. G. & Menzel, R. (2001). The Role of Glomeruli in the Neural Representation of Odours: Results from Optical Recording Studies. *Journal of Insect Physiology*, 47, 115-130.
- Gazit, I. & Terkel, J. (2003). Explosives Detection by Sniffer Dogs Following Strenuous Physical Activity. *Applied Animal Behaviour Science*, 81, 149-161.
- Geiger, H. J. (2000). The Impact of War on Human Rights. Washington, D.C., American Public Health Association [APHA], 2000.
- Getchell, T. V., Margolis, F. L. & Getchell, M. L. (1984). Perireceptor and Receptor Events in Vertebrate Olfaction. *Progress in Neurobiology*, 23, 317-345.
- Ghelardini, C., Galeotti, N., Di Cesare Mannelli, L., Mazzanti, G. & Bartolini, A. (2001). Local Anaesthetic Activity of B-Caryophyllene. *Il Farmaco*, 56, 387-389.
- Grant, V. (1950). The Flower Constancy of Bees. *The Botanical Review*, 16, 379-398.
- Gronenberg, W. (2001). Subdivisions of Hymenopteran Mushroom Body Calyces by Their Afferent Supply. *The Journal of Comparative Neurology*, 435, 474-489.
- Gruntman, E. & Turner, G. C. (2013). Integration of the Olfactory Code Across Dendritic Claws of Single Mushroom Body Neurons. *Nat Neurosci*, 16, 1821-1829.
- Guillot, S., Peytavi, L., Bureau, S., Boulanger, R., Lepoutre, J.-P., Crouzet, J. & Schorr-Galindo, S. (2006). Aroma Characterization of Various Apricot Varieties Using Headspace–Solid Phase Microextraction Combined with Gas Chromatography–Mass Spectrometry and Gas Chromatography–Olfactometry. *Food Chemistry*, 96, 147-155.
- Hallem, E. A., Ho, M. G. & Carlson, J. R. (2004). The Molecular Basis of Odor Coding in the *Drosophila* Antenna. *Cell*, 117, 965-979.
- Hammer, M. & Menzel, R. (1995). Learning and Memory in the Honeybee. *Journal of Neuroscience*, 15, 1617-1630.
- Hammond, G. & Blankenship, M. (2009). *Apis mellifera* [Online]. Available: http://animaldiversity.org/accounts/Apis_mellifera/ [Accessed May 03 2017].
- Handa, S. & Sharma, A. (1990). Hepatoprotective Activity of Andrographolide from *Andrographis paniculata* Against Carbontetrachloride. *The Indian journal of medical research*, 92, 276-283.

- Hansson, B. S. & Anton, S. (2000). Function and Morphology of the Antennal Lobe: New Developments. *Annual Review of Entomology*, 45, 203-231.
- Heisenberg, M. (1998). What Do the Mushroom Bodies Do for the Insect Brain? An Introduction. *Learning & Memory*, 5, 1-10.
- Hill, C. A., Fox, A. N., Pitts, R. J., Kent, L. B., Tan, P. L., Chrystal, M. A., Cravchik, A., Collins, F. H., Robertson, H. M. & Zwiebel, L. J. (2002). G Protein-Coupled Receptors in *Anopheles gambiae*. *Science*, 298, 176-178.
- Hodgson, M. (2006). Sniffer Bees: New Flying Squads in War Against Terror. *The Independent*.
- Hossain, M. A., Roy, B., Ahmed, K., Chowdhury, A. S. & Rashid, M. (2007). Antidiabetic Activity of *Andrographis paniculata*. *Dhaka University Journal of Pharmaceutical Sciences*, 6, 15-20.
- Ibáñez, E., López-Sebastián, S., Ramos, E., Tabera, J. & Reglero, G. (1998). Analysis of Volatile Fruit Components by Headspace Solid-Phase Microextraction. *Food Chemistry*, 63, 281-286.
- Jabbari, E., Kim, D. H., Lee, L. P., Ghaemmaghami, A. & Khademhosseini, A. (2014). *Handbook of Biomimetics and Bioinspiration: Biologically-Driven Engineering of Materials, Processes, Devices, and Systems(In 3 Volumes)*, World Scientific Publishing Company.
- Jackson, J. C. & Robert, D. (2006). Nonlinear Auditory Mechanism Enhances Female Sounds for Male Mosquitoes. *Proceedings of the National Academy of Sciences*, 103, 16734-16739.
- Jacquin-Joly, E. & Merlin, C. (2004). Insect Olfactory Receptors: Contributions of Molecular Biology to Chemical Ecology. *Journal of chemical ecology*, 30, 2359-2397.
- Jarukamjorn, K. & Nemoto, N. (2008). Pharmacological Aspects of *Andrographis paniculata* on Health and Its Major Diterpenoid Constituent Andrographolide. *Journal of Health Science*, 54, 370-381.
- Jones, J. C., Helliwell, P., Beekman, M., Maleszka, R. & Oldroyd, B. P. (2005). The Effects of Rearing Temperature on Developmental Stability and Learning and Memory in the Honey Bee, *Apis mellifera*. *Journal of Comparative Physiology A*, 191, 1121-1129.
- Kalaiselvan, A., Gokulakrishnan, K. & Anand, T. (2012). Gas Chromatography-Mass Spectrum Analysis of Bioactive Components of the Ethanol Extract of

- Andrographis paniculata*. *Journal of Pharmaceutical and Biomedical Sciences*, 20.
- Kalaivani, C. S., Sathish, S. S., Janakiraman, N. & Johnson, M. (2012). GC-MS Studies on *Andrographis paniculata* (Burm.F.) Wall. Ex Nees - A Medicinally Important Plant. *International Journal of Medicinal and Aromatic Plants*, 2, 69-74.
- Kapil, A., Koul, I. B., Banerjee, S. K. & Gupta, B. D. (1993). Antihepatotoxic Effects of Major Diterpenoid Constituents of *Andrographis paniculata*. *Biochemical Pharmacology*, 46, 182-185.
- Karacabey, E., Bayindirli, L., Artik, N. & Mazza, G. (2013). Modeling Solid-Liquid Extraction Kinetics of Trans-Resveratrol and Trans-E-Viniferin from Grape Cane. *Journal of Food Process Engineering*, 36, 103-112.
- Kataria, S. (2011). Gas Chromatography-Mass Spectrometry: Applications. *International Journal of Pharmaceutical & Biological Archives*, 2, 1544-1560.
- Kato, A. & Touhara, K. (2009). Mammalian Olfactory Receptors: Pharmacology, G Protein Coupling and Desensitization. *Cellular and Molecular Life Sciences*, 66, 3743.
- Kee, T., Sanda, P., Gupta, N., Stopfer, M. & Bazhenov, M. (2015). Feed-Forward Versus Feedback Inhibition in a Basic Olfactory Circuit. *PLoS computational biology*, 11, e1004531.
- Keil, T. A. (1999). Morphology and Development of the Peripheral Olfactory Organs. *Insect olfaction*. Springer.
- Kendra, C. (2013). *What Is a Conditioned Response?* [Online].
- Kenyon, F. (1896). The Brain of the Bee. A Preliminary Contribution to the Morphology of the Nervous System of the Arthropoda. *Journal of Comparative Neurology*, 6, 133-210.
- Khoyi, M. R. & Hesari, J. (2007). Osmotic Dehydration Kinetics of Apricot Using Sucrose Solution. *Journal of Food Engineering*, 78, 1355-1360.
- King, T. L., Horine, F. M., Daly, K. C. & Smith, B. H. (2003). Explosives Detection with Hard-Wired Moths. Proceedings of the 20th IEEE Instrumentation Technology Conference (Cat. No.03CH37412), 20-22 May 2003. 1669-1672 vol.2.
- Knierim, J. J. (2014). Chapter 19 - Information Processing in Neural Networks. *From Molecules to Networks (Third Edition)*. Boston: Academic Press.

- Kolb, B. & Ettre, L. S. (2006). *Static Headspace-Gas Chromatography: Theory and Practice*, John Wiley & Sons.
- Krieger, J. & Breer, H. (1999). Olfactory Reception in Invertebrates. *Science*, 286, 720-723.
- Krieger, J., Mameli, M. & Breer, H. (1997). Elements of the Olfactory Signaling Pathways in Insect Antennae. *Invertebrate Neuroscience*, 3, 137-144.
- Kuang, X., Shankar, T. J., Bi, X. T., Sokhansanj, S., Lim, C. J. & Melin, S. (2008). Characterization and Kinetics Study of Off-Gas Emissions from Stored Wood Pellets. *Annals of occupational hygiene*, 52, 675-683.
- Kumar, R. A., Sridevi, K., Kumar, N. V., Nanduri, S. & Rajagopal, S. (2004). Anticancer and Immunostimulatory Compounds from *Andrographis paniculata*. *Journal of ethnopharmacology*, 92, 291-295.
- Lambropoulou, D. A. & Albanis, T. A. (2001). Optimization of Headspace Solid-Phase Microextraction Conditions for the Determination of Organophosphorus Insecticides in Natural Waters. *Journal of Chromatography A*, 922, 243-255.
- Laue, M., Steinbrecht, R. & Ziegelberger, G. (1994). Immunocytochemical Localization of General Odorant-Binding Protein in Olfactory Sensilla of the Silkworm *Antheraea polyphemus*. *Naturwissenschaften*, 81, 178-180.
- Laurent, G. (1999). A Systems Perspective on Early Olfactory Coding. *Science*, 286, 723-728.
- Laurent, G. (2002). Olfactory Network Dynamics and the Coding of Multidimensional Signals. *Nat Rev Neurosci*, 3, 884-895.
- Layton, J. (2006). *How Can You Train Honeybees to Sniff for Bombs?* [Online]. Available: <http://science.howstuffworks.com/bomb-sniffing-bees.htm> [Accessed 9 May 2017].
- Leitch, O., Anderson, A., Paul Kirkbride, K. & Lennard, C. (2013). Biological Organisms as Volatile Compound Detectors: A Review. *Forensic Science International*, 232, 92-103.
- Letzkus, P., Ribi, W. A., Wood, J. T., Zhu, H., Zhang, S.-W. & Srinivasan, M. V. (2006). Lateralization of Olfaction in the Honeybee *Apis mellifera*. *Current Biology*, 16, 1471-1476.
- Lin, H., Yan, Y., Zhao, T., Peng, L., Zou, H., Li, J., Yang, X., Xiong, Y., Wang, M. & Wu, H. (2013). Rapid Discrimination of Apiaceae Plants by Electronic Nose

- Coupled with Multivariate Statistical Analyses. *Journal of Pharmaceutical and Biomedical Analysis*, 84, 1-4.
- Llompart, M., Li, K. & Fingas, M. (1999). Headspace Solid Phase Microextraction (Hsspme) for the Determination of Volatile and Semivolatile Pollutants in Soils. *Talanta*, 48, 451-459.
- Loo, S. K. & Bitterman, M. E. (1992). Learning in Honeybees (*Apis mellifera*) As a Function of Sucrose Concentration. *Journal of Comparative Psychology*, 106, 29-36.
- Ma, L. & Selim, H. M. (1994). Predicting Atrazine Adsorption-Desorption in Soils: A Modified Second-Order Kinetic Model. *Water Resources Research*, 30, 447-456.
- Malun, D., Plath, N., Giurfa, M., Moseleit, A. D. & Müller, U. (2002). Hydroxyurea-Induced Partial Mushroom Body Ablation in the Honeybee *Apis mellifera*: Volumetric Analysis and Quantitative Protein Determination. *Journal of Neurobiology*, 50, 31-44.
- Maskan, M. (2001). Kinetics of Colour Change of Kiwifruits During Hot Air and Microwave Drying. *Journal of Food Engineering*, 48, 169-175.
- Mccabe-Sellers, B. J. & Beattie, S. E. (2004). Food Safety: Emerging Trends in Foodborne Illness Surveillance and Prevention. *Journal of the American Dietetic Association*, 104, 1708-1717.
- Mcdonald, S. T., Bolliet, D. A. & Hayes, J. E. (2016). *Chemesthesis: Chemical Touch in Food and Eating*, Wiley.
- Meiners, T., Wäckers, F. & Lewis, W. J. (2002). The Effect of Molecular Structure on Olfactory Discrimination by the Parasitoid *Microplitis croceipes*. *Chemical senses*, 27, 811-816.
- Menzel, R. (1985). *Learning in Honey Bees in an Ecological and Behavioral Context*, Sunderland, Sinauer Associates.
- Menzel, R. (1999). Memory Dynamics in the Honeybee. *Journal of Comparative Physiology A: Neuroethology, Sensory, Neural, and Behavioral Physiology*, 185, 323-340.
- Menzel, R. & Bitterman, M. E. (1983). Learning by Honeybees in an Unnatural Situation. In: Huber, F. & Markl, H. (eds.) *Neuroethology and Behavioral Physiology: Roots and Growing Points*. Berlin, Heidelberg: Springer Berlin Heidelberg.

- Menzel, R. & Erber, J. (1978). Learning and Memory in Bees. *Sci. Am*, 239, 102-110.
- Menzel, R., Erber, J. & Masuhr, T. (1974). Learning and Memory in the Honeybee. *Experimental analysis of insect behaviour*. Springer.
- Menzel, R., Greggers, U. & Hammer, M. (1993). Functional Organization of Appetitive Learning and Memory in a Generalist Pollinator, the Honey Bee. *Insect learning*. Springer.
- Mishra, U. S., Mishra, A., Kumari, R., Murthy, P. N. & Naik, B. S. (2009). Antibacterial Activity of Ethanol Extract of *Andrographis paniculata*. *Indian Journal of Pharmaceutical Sciences*, 71, 436-438.
- Mizunami, M., Weibrecht, J. M. & Strausfeld, N. J. (1993). A New Role for the Insect Mushroom Bodies: Place Memory and Motor Control. Proceedings Of The Workshop on Locomotion Control in Legged Invertebrates on Biological Neural Networks in Invertebrate Neuroethology and Robotics. Academic Press Professional, Inc., 199-225.
- Mobbs, P. (1982). The Brain of the Honeybee *Apis mellifera*. I. The Connections and Spatial Organization of the Mushroom Bodies. *Philosophical Transactions of the Royal Society of London B: Biological Sciences*, 298, 309-354.
- Mobbs, P. (1984). Neural Networks in the Mushroom Bodies of the Honeybee. *Journal of insect physiology*, 30, 43-58.
- Mombaerts, P. (2004). Genes and Ligands for Odorant, Vomeronasal and Taste Receptors. *Nat Rev Neurosci*, 5, 263-278.
- Mott, M. (2004). Bees, Giant African Rats Used to Sniff Landmines. *National Geographic News*, 10.
- Müller, D., Abel, R., Brandt, R., Zöckler, M. & Menzel, R. (2002). Differential Parallel Processing of Olfactory Information in the Honeybee, *Apis mellifera* L. *Journal of Comparative Physiology A*, 188, 359-370.
- Müller, U. (2000). Prolonged Activation of camp-Dependent Protein Kinase During Conditioning Induces Long-Term Memory in Honeybees. *Neuron*, 27, 159-168.
- Müller, U. (2002). Learning in Honeybees: From Molecules to Behaviour1. *Zoology*, 105, 313-320.
- Müller, U. (2012). The Molecular Biology of Learning and Memory – Memory Phases and Signaling Cascades. In: Galizia, C. G., Eisenhardt, D. & Giurfa, M. (eds.) *Honeybee Neurobiology and Behavior: A Tribute to Randolph Menzel*. Dordrecht: Springer Netherlands.

- Namasivayam, C. & Kavitha, D. (2002). Removal of Congo Red from Water by Adsorption onto Activated Carbon Prepared from Coir Pith, an Agricultural Solid Waste. *Dyes and Pigments*, 54, 47-58.
- Okpokwasili, G. C. & Nweke, C. O. (2005). Microbial Growth and Substrate Utilization Kinetics. *African Journal of Biotechnology*, 5, 305-517.
- Olsen, S. R. & Wilson, R. I. (2008). Lateral Presynaptic Inhibition Mediates Gain Control in an Olfactory Circuit. *Nature*, 452, 956.
- Olson, D., Rains, G., Meiners, T., Takasu, K., Tertuliano, M., Tumlinson, J., Wäckers, F. & Lewis, W. J. (2003). Parasitic Wasps Learn and Report Diverse Chemicals with Unique Conditionable Behaviors. *Chemical Senses*, 28, 545-549.
- Page, R. E. & Peng, C. Y. S. (2001). Aging and Development in Social Insects with Emphasis on the Honey Bee, *Apis mellifera L.* *Experimental Gerontology*, 36, 695-711.
- Palaniswamy, U. R. (2005). Effect of Light Intensity on the Pigment Composition and Oxalic Acid Concentrations in Kalamegh (*Andrographis paniculata*) Leaf. International Society for Horticultural Science (ISHS), Leuven, Belgium, 109-114.
- Pankiw, T., Page Jr, R. E. & Kim Fondrk, M. (1998). Brood Pheromone Stimulates Pollen Foraging in Honey Bees (*Apis mellifera*). *Behavioral Ecology and Sociobiology*, 44, 193-198.
- Parvataneni, R. & Lakshmi, R. (2010). *Antimicrobial Activity of the Chloroform Extracts of the Root and the Stem of Andrographis paniculata Nees.*
- Pavlov, I. P. (1927). *Conditional Reflexes: An Investigation of the Physiological Activity of the Cerebral Cortex*, Oxford, England, Oxford Univ. Press.
- Peleg, M. (1988). An Empirical Model for the Description of Moisture Sorption Curves. *Journal of Food Science*, 53, 1216-1217.
- Pelletier, Y. & Mcleod, C. D. (1994). Obstacle Perception by Insect Antennae During Terrestrial Locomotion. *Physiological Entomology*, 19, 360-362.
- Pelosi, P. (1994). Odorant-Binding Proteins. *Critical Reviews in Biochemistry and Molecular Biology*, 29, 199-228.
- Pelosi, P. (1996). Perireceptor Events in Olfaction. *J. Neurobiol.*, 30, 3-19.
- Pelosi, P. (2001). The Role of Perireceptor Events in Vertebrate Olfaction. *Cellular and Molecular Life Sciences CMLS*, 58, 503-509.

- Pelz, C., Gerber, B. & Menzel, R. (1997). Odorant Intensity as a Determinant for Olfactory Conditioning in Honeybees: Roles in Discrimination, Overshadowing and Memory Consolidation. *The Journal of Experimental Biology*, 200, 837-847.
- Perez-Orive, J., Mazor, O., Turner, G. C., Cassenaer, S., Wilson, R. I. & Laurent, G. (2002). Oscillations and Sparsening of Odor Representations in the Mushroom Body. *Science*, 297, 359-365.
- Pham-Delegue, M., Etievant, P., Guichard, E. & Masson, C. (1989). Sunflower Volatiles Involved in Honeybee Discrimination Among Genotypes and Flowering Stages. *Journal of Chemical Ecology*, 15, 329-343.
- Purves, D., Fitzpatrick, D., Katz, L. C., Lamantia, A. S., Mcnamara, J. O., Williams, S. M. & Augustine, G. J. (2001). *Neuroscience*, Sinauer Associates.
- Rains, G. C., Tomberlin, J., D'alessandro, M. & Lewis, W. J. (2004). Limits of Volatile Chemical Detection of a Parasitoid Wasp, *Microplitis croceipes*, and an Electronic Nose: A Comparative Study. *Transactions of the ASAE*, 47, 2145.
- Rains, G. C., Tomberlin, J. K. & Kulasiri, D. (2008). Using Insect Sniffing Devices for Detection. *Trends in biotechnology*, 26, 288-294.
- Regueiro, J., Becerril, E., Garcia-Jares, C. & Llompart, M. (2009). Trace Analysis of Parabens, Triclosan and Related Chlorophenols in Water by Headspace Solid-Phase Microextraction with in Situ Derivatization and Gas Chromatography–Tandem Mass Spectrometry. *Journal of Chromatography A*, 1216, 4693-4702.
- Resh, V. H. & Cardé, R. T. (2009). *Encyclopedia of Insects*, Elsevier Science.
- Robertson, H. M. & Wanner, K. W. (2006). The Chemoreceptor Superfamily in the Honey Bee, *Apis mellifera*: Expansion of the Odorant, but not Gustatory, Receptor Family. *Genome Research*, 16, 1395-1403.
- Rocha, S., Martins, V. M. R., Barros, A., Delgadillo, I. & Coimbra, M. A. (2001). Headspace Solid Phase Microextraction (Spme) Analysis of Flavor Compounds in Wines. Effect of the Matrix Volatile Composition in the Relative Response Factors in a Wine Model. *Journal of Agricultural and Food Chemistry*, 5142-5151.
- Rodacy, P. J., Bender, S., Bromenshenk, J., Henderson, C. & Bender, G. (2002). Training and Deployment of Honeybees to Detect Explosives and Other Agents of Harm. *AeroSense SPIE*, 8.

- Rothschild, M. & Hinton, H. E. (1968). Holding Organs on the Antennae of Male Fleas. *Proceedings of the Royal Entomological Society of London. Series A, General Entomology*, 43, 105-107.
- Ruiz, J., Cava, R., Ventanas, J. & Jensen, M. T. (1998). Headspace Solid Phase Microextraction for the Analysis of Volatiles in a Meat Product: Dry-Cured Iberian Ham. *Journal of Agricultural and Food Chemistry*, 46, 4688-4694.
- Rybak, J. (2012). The Digital Honey Bee Brain Atlas. *Honeybee Neurobiology and Behavior*. Springer.
- Sala, C., Mestres, M., Martí, M. P., Busto, O. & Guasch, J. (2002). Headspace Solid-Phase Microextraction Analysis of 3-Alkyl-2-Methoxypyrazines in Wines. *Journal of Chromatography A*, 953, 1-6.
- Sammataro, D. & Avitabile, A. (1998). *The Beekeeper's Handbook*, Comstock Pub. Associates.
- Sandoz, J.-C. (2011). Behavioral and Neurophysiological Study of Olfactory Perception and Learning in Honeybees. *Frontiers in Systems Neuroscience*, 5.
- Sato, K. & Touhara, K. (2008). Insect Olfaction: Receptors, Signal Transduction, and Behavior. *Chemosensory systems in mammals, fishes, and insects*. Springer.
- Schreiner, K. (2002). Landmine Detection Research Pushes Forward, Despite Challenges. *IEEE Intelligent Systems*, 17, 4-7.
- Scheiner, R., Erber, J. & Page Jr., R. E. (1999). Tactile Learning and the Individual Evaluation of the Reward in Honey Bees (*Apis mellifera* L.). *Journal of Comparative Physiology A*, 185, 1-10.
- Scheiner, R., Page, R. E. & Erber, J. (2004). Sucrose Responsiveness and Behavioral Plasticity in Honey Bees (*Apis mellifera*). *Apidologie*, 35, 133-142.
- Schiestl, F. P., Ayasse, M., Paulus, H. F., Erdmann, D. & Francke, W. (1997). Variation of Floral Scent Emission and Postpollination Changes in Individual Flowers of *Ophrys sphegodes* subsp. *sphogodes*. *Journal of Chemical Ecology*, 23, 2881-2895.
- Schott, M., Klein, B. & Vilcinskas, A. (2015). Detection of Illicit Drugs by Trained Honeybees (*Apis mellifera*). *PLOS ONE*, 10, e0128528.
- Schröter, U. & Menzel, R. (2003). A New Ascending Sensory Tract to the Calyces of the Honeybee Mushroom Body, the Subesophageal-Calycal Tract. *The Journal of Comparative Neurology*, 465, 168-178.

- Seeley, T. D. (1982). Adaptive Significance of the Age Polyethism Schedule in Honeybee Colonies. *Behavioral Ecology and Sociobiology*, 11, 287-293.
- Shabila, N. P., Taha, H. I. & Al-Hadithi, T. S. (2010). Landmine Injuries at the Emergency Management Center in Erbil, Iraq. *Conflict and Health*, 4, 15.
- Sheeja, K., Shihab, P. K. & Kuttan, G. (2006). Antioxidant and Anti-Inflammatory Activities of the Plant *Andrographis paniculata* Nees. *Immunopharmacology and Immunotoxicology*, 28, 129-140.
- Shimoda, M. & Shibamoto, T. (1990). Isolation and Identification of Headspace Volatiles from Brewed Coffee with an On-Column GC/MS Method. *Journal of Agricultural and Food Chemistry*, 38, 802-804.
- Steffen, A. & Pawliszyn, J. (1996). Analysis of Flavor Volatiles Using Headspace Solid-Phase Microextraction. *Journal of Agricultural and Food Chemistry*, 44, 2187-2193.
- Steinbrecht, R., Ozaki, M. & Ziegelberger, G. (1992). Immunocytochemical Localization of Pheromone-Binding Protein in Moth Antennae. *Cell and Tissue Research*, 270, 287-302.
- Stocker, R. F. (1994). The Organization of the Chemosensory System in *Drosophila melanogaster*: A Review. *Cell and Tissue Research*, 275, 3-26.
- Stopfer, M. (2014). Central Processing in the Mushroom Bodies. *Current Opinion in Insect Science*, 6, 99-103.
- Stopfer, M., Jayaraman, V. & Laurent, G. (2003). Intensity Versus Identity Coding in an Olfactory System. *Neuron*, 39, 991-1004.
- Strausfeld, N. J. (2002). Organization of the Honey Bee Mushroom Body: Representation of the Calyx Within the Vertical and Gamma Lobes. *The Journal of Comparative Neurology*, 450, 4-33.
- Suckling, D. & L Sagar, R. (2011). *Honeybees Apis mellifera Can Detect the Scent of Mycobacterium tuberculosis*.
- Suebsasana, S., Pongnaratorn, P., Sattayasai, J., Arkaravichien, T., Tiamkao, S. & Aromdee, C. (2009). Analgesic, Antipyretic, Anti-Inflammatory and Toxic Effects of Andrographolide Derivatives in Experimental Animals. *Archives of Pharmacal Research*, 32, 1191-1200.
- Sule, A., Ahmed, Q. U., Latip, J., Samah, O. A., Omar, M. N., Umar, A. & Dogarai, B. B. S. (2012). Antifungal Activity of *Andrographis paniculata* Extracts and

- Active Principles Against Skin Pathogenic Fungal Strains in Vitro. *Pharmaceutical Biology*, 50, 850-856.
- Sule, A., Ahmed, Q. U., Samah, O. A. & Omar, M. N. (2011). Bacteriostatic and Bactericidal Activities of *Andrographis paniculata* Extracts on Skin Disease Causing Pathogenic Bacteria. *Journal of Medicinal Plants Research*, 5, 7-14.
- Surrency, A. B., Graitcer, P. L. & Henderson, A. K. (2007). Key Factors for Civilian Injuries and Deaths from Exploding Landmines and Ordnance. *Injury Prevention*, 13, 197-201.
- Taylor-McCabe, K. J., Wingo, R. M. & Haarmann, T. K. (2008a). Honey Bees (*Apis mellifera*) as Explosives Detectors: Exploring Proboscis Extension Reflex Conditioned Response to Trinitrotolulene (TNT). Los Alamos National Laboratory (LANL).
- Taylor-McCabe, K. J., Wingo, R. M., Haarmann, T. K. & Maurer, S. E. (2008b). Is Olfactory-Based Associative Learning in Honey Bees Linked to Antennae Morphology? : Los Alamos National Laboratory (LANL).
- Tertuliano, M., Olson, D., Rains, G. & Lewis, W. (2004). Influence of Handling and Conditioning Protocol on Learning and Memory of *Microplitis croceipes*. *Entomologia experimentalis et applicata*, 110, 165-172.
- Tertuliano, M., Tomberlin, J. K., Jurjevic, Z., Wilson, D., Rains, G. C. & Lewis, W. J. (2005). The Ability of Conditioned *Microplitis croceipes* (Hymenoptera: Braconidae) to Distinguish Between Odors of Aflatoxigenic and Non-Aflatoxigenic Fungal Strains. *CHEMOECOLOGY*, 15, 89-95.
- Thakur, A. K., Chatterjee, S. S. & Kumar, V. (2015). Adaptogenic Potential of Andrographolide: An Active Principle of the King of Bitters (*Andrographis paniculata*). *Journal of Traditional and Complementary Medicine*, 5, 42-50.
- Tichy, H. & Kallina, W. (2010). Insect Hygroreceptor Responses to Continuous Changes in Humidity and Air Pressure. *Journal of Neurophysiology*, 103, 3274-3286.
- Tollsten, L. & Bergström, G. (1988). Headspace Volatiles of Whole Plants and Macerated Plant Parts of Brassica and Sinapis. *Phytochemistry*, 27, 4013-4018.
- Tomberlin, J. K., Tertuliano, M., Rains, G. & Lewis, W. J. (2005). Conditioned *Microplitis croceipes cresson* (Hymenoptera: Braconidae) Detect and Respond to 2, 4-DNT: Development of a Biological Sensor. *Journal of Forensic Science*, 50, JFS2005014-4.

- Triplehorn, C. A. J., Borror, N. F., Triplehorn, D. J. C. A. & Johnson, N. F. (2005). *Borror and DeLong's Introduction to the Study of Insects*.
- Tully, T. & Quinn, W. (1985). Classical Conditioning and Retention in Normal and Mutant *Drosophila melanogaster*. *Journal of Comparative Physiology A*, 157, 263-277.
- Uchida, N., Takahashi, Y. K., Tanifuji, M. & Mori, K. (2000). Odor Maps in the Mammalian Olfactory Bulb: Domain Organization and Odorant Structural Features. *Nature neuroscience*, 3, 1035-1043.
- Vasantha, S., Subramaniyan, A., Madhumathi, V. & Abubacker, M. N. (2013). Gas Chromatography-Mass Spectrometry (GC-MS) Analysis of Ethanolic Leaf Extract of *Andrographis paniculata* Nees. *International Journal of Medicine and Biosciences*, 2, 16-20.
- Viegas, J. (2004). Sniffer Moths Detect Whiff of Explosives. *ABC Science*.
- Vogt, R. G., Rybczynski, R. & Lerner, M. R. (1991). Molecular Cloning and Sequencing of General Odorant-Binding Proteins GOBP1 and GOBP2 from the Tobacco Hawk Moth *Manduca sexta*: Comparisons with Other Insect OBPs and Their Signal Peptides. *Journal of Neuroscience*, 11, 2972-2984.
- Von Frisch, K. (1967). The Dance Language and Orientation of Bees.
- Voskresenskaya, A. (1957). The Role of Mushroom Bodies (*Corpora pedunculata*) of the Supraesophageal Ganglion in the Conditioned Reflexes of the Honey Bee. *Doklady Akademii Nauk Sssr*, 112, 964-967.
- Vosshall, L. B., Amrein, H., Morozov, P. S., Rzhetsky, A. & Axel, R. (1999). A Spatial Map of Olfactory Receptor Expression in the *Drosophila* Antenna. *Cell*, 96, 725-736.
- Vosshall, L. B., Wong, A. M. & Axel, R. (2000). An Olfactory Sensory Map in the Fly Brain. *Cell*, 102, 147-159.
- Weingart, G., Kluger, B., Forneck, A., Krska, R. & Schuhmacher, R. (2012). Establishment and Application of a Metabolomics Workflow for Identification and Profiling of Volatiles from Leaves of *Vitis vinifera* by HS-SPME-GC-MS. *Phytochemical Analysis*, 23, 345-358.
- Whiton, R. S. & Zoecklein, B. W. (2000). Optimization of Headspace Solid-Phase Microextraction for Analysis of Wine Aroma Compounds. *American Journal of Enology and Viticulture*, 51, 379-382.

- Wiar, C., Kumar, K., Yusof, M. Y., Hamimah, H., Fauzi, Z. M. & Sulaiman, M. (2005). Antiviral Properties of Ent-Labdene Diterpenes of *Andrographis paniculata* Nees, Inhibitors of Herpes Simplex Virus Type 1. *Phytotherapy Research*, 19, 1069-1070.
- Wilson, A. D. & Baietto, M. (2009). Applications and Advances in Electronic-Nose Technologies. *Sensors (Basel, Switzerland)*, 9, 5099-5148.
- Wilson, D. A. & Stevenson, R. J. (2006). *Learning to Smell: Olfactory Perception from Neurobiology to Behavior*, Johns Hopkins University Press.
- Winston, M. L. (1991). *The Biology of the Honey Bee*, Harvard University Press.
- Wright, G. A., Carlton, M. & Smith, B. H. (2009). A Honeybee's Ability to Learn, Recognize, and Discriminate Odors Depends upon Odor Sampling Time and Concentration. *Behavioral neuroscience*, 123, 36-43.
- Wright, G. A. & Smith, B. H. (2004). Different Thresholds for Detection and Discrimination of Odors in the Honey bee (*Apis mellifera*). *Chemical Senses*, 29, 127-135.
- Wüstenberg, D., Gerber, B. & Menzel, R. (1998). Long- but not Medium-Term Retention of Olfactory Memories in Honeybees Is Impaired by Actinomycin D and Anisomycin. *European Journal of Neuroscience*, 10, 2742-2745.
- Yao, X.-J., Wainberg, M. A. & Parniak, M. A. (1992). Mechanism of Inhibition of Hiv-1 Infection in Vitro by Purified Extract of *Prunella vulgaris*. *Virology*, 187, 56-62.
- Zhang, Z. & Pawliszyn, J. (1993). Headspace Solid-Phase Microextraction. *Analytical chemistry*, 65, 1843-1852.
- Zhang, Z., Yang, M. J. & Pawliszyn, J. (1994). Solid-Phase Microextraction. A Solvent-Free Alternative for Sample Preparation. *Analytical Chemistry*, 66, 844A-853A.

APPENDIX A

RESULTS OF KINETIC STUDIES OF VOLATILE MARKER COMPOUNDS

A.1 Kinetic Study of 5,6,7,7 α -tetrahydro-4,4,7 α -trimethyl-2(4H)-benzofuranone

$$\text{Equation: } C_t = \frac{t}{\left(\frac{1}{h}\right) + \left(\frac{t}{C_e}\right)}$$

Time	Observed	Predicted	(y-mean(y)) ²	(y _p -y _o) ²
10	133.5345	135.5798	10156.1677	4.1831
15	181.8200	170.3824	2755.4514	130.8171
20	194.0304	195.4706	1622.6317	2.0739
25	207.4390	214.4135	722.1744	48.6427
40	246.1763	250.8828	140.7523	22.1515
50	267.2822	265.9618	1087.0134	1.7435
60	270.4553	277.0636	1306.3126	43.6693
80	306.0444	292.3158	5145.4836	188.4731
110	302.02900	306.1029	4585.5442	16.5966

Time: Sample heating time

Observed: Observed peak area

Predicted: Predicted peak area

[y-mean(y)]²: [Observed peak area-mean value of observed peak area]²

(y_p-y_o)²: (Predicted peak area-Observed peak area)²

By using Solver function in Microsoft Excel, the values of parameters can be obtained.


Parameter	Value
<i>h</i>	22.1252
<i>C_e</i>	350.1413

$$R^2 = 1 - \frac{\sum(y_p - y_o)^2}{\sum[y - \text{mean}(y)]^2}$$

$$R^2 = 0.9833$$

$$RMSE = \sqrt{\frac{\sum(y_p - y_o)^2}{\text{no. of data}}}$$

$$RMSE = 7.1364$$



A.2 Kinetic Study of Caryophyllene

$$\text{Equation: } C_t = \frac{t}{\left(\frac{1}{h}\right) + \left(\frac{t}{C_e}\right)}$$

Time	Observed	Predicted	[y-mean(y)] ²	(y _p -y _o) ²
10	2050.2065	2060.7405	81268.4535	110.9644
30	2399.2107	2349.6338	4086.7797	2457.8643
40	2356.7056	2391.5424	458.9366	1213.6016
50	2433.4385	2417.4128	9634.5446	256.8219
80	2436.8525	2457.2853	10316.4194	417.4964
90	2339.4956	2464.8138	17.7481	15704.6587
110	2255.1301	2475.8472	6424.4447	48716.0277

Time: Sample heating time

Observed: Observed peak area

Predicted: Predicted peak area

[y-mean(y)]²: [Observed peak area-mean value of observed peak area]²

(y_p-y_o)²: (Predicted peak area-Observed peak area)²

By using Solver function in Microsoft Excel, the values of parameters can be obtained.


Parameter	Value
<i>h</i>	1117.3641
<i>C_e</i>	2526.7449

$$R^2 = 1 - \frac{\sum(y_p - y_o)^2}{\sum[y - \text{mean}(y)]^2}$$

$$R^2 = 0.9579$$

$$RMSE = \sqrt{\frac{\sum(y_p - y_o)^2}{\text{no. of data}}}$$

$$RMSE = 25.2325$$



A.3 Kinetic Study of β -elemene

$$\text{Equation: } C_t = \frac{t}{\left(\frac{1}{h}\right) + \left(\frac{t}{C_e}\right)}$$

Time	Observed	Predicted	$[y-\text{mean}(y)]^2$	$(y_p-y_o)^2$
10	642.5048	659.8107	39622.8778	299.4957
30	853.4323	827.0659	140.9569	695.1819
50	892.4230	871.2358	2587.0700	448.8955
60	885.9645	883.0254	1971.7879	8.6384
80	918.1769	898.2188	5870.1890	398.3233
110	856.8569	911.0443	234.0046	2936.2673

Time: Sample heating time

Observed: Observed peak area

Predicted: Predicted peak area

$[y-\text{mean}(y)]^2$: [Observed peak area-mean value of observed peak area]²

$(y_p-y_o)^2$: (Predicted peak area-Observed peak area)²

By using Solver function in Microsoft Excel, the values of parameters can be obtained.


Parameter	Value
h	217.5146
C_e	947.1069

$$R^2 = 1 - \frac{\sum(y_p - y_o)^2}{\sum[y - \text{mean}(y)]^2}$$

$$R^2 = 0.9051$$

$$RMSE = \sqrt{\frac{\sum(y_p - y_o)^2}{\text{no. of data}}}$$

$$RMSE = 28.2454$$



A.4 Kinetic Study of 6,10,14-trimethyl-2-pentadecanone

$$\text{Equation: } C_t = C_e [1 - [F e^{-k_1 t}] - [(1 - F) e^{-k_2 t}]]$$

Time	Observed	Predicted	[y-mean(y)] ²	(y _p -y _o) ²
10	61.5333	61.5426	3158.0863	8.45112E-05
15	118.1757	118.1565	0.1984	0.00037
20	128.2121	127.4745	109.8700	0.5440
25	125.8151	129.0082	65.3657	10.1956
40	130.6017	129.3090	165.6752	1.6712
50	128.6495	129.3103	119.2317	0.4366
60	131.1240	129.3103	179.3938	3.2895

Time: Sample heating time

Observed: Observed peak area

Predicted: Predicted peak area

[y-mean(y)]²: [Observed peak area-mean value of observed peak area]²

(y_p-y_o)²: (Predicted peak area-Observed peak area)²

By using Solver function in Microsoft Excel, the values of parameters can be obtained.


Parameter	Value
C_e	129.31032
F	-18.34602
k_1	135.48671
k_2	0.3608615

$$R^2 = 1 - \frac{\sum(y_p - y_o)^2}{\sum[y - \text{mean}(y)]^2}$$

$$R^2 = 0.9958$$

$$RMSE = \sqrt{\frac{\sum(y_p - y_o)^2}{\text{no. of data}}}$$

$$RMSE = 1.0736$$



A.5 Kinetic Study of Apiol

$$\text{Equation: } C_t = C_e [1 - [F e^{-k_1 t}] - [(1 - F) e^{-k_2 t}]]$$

Time	Observed	Predicted	$[y - \text{mean}(y)]^2$	$(y_p - y_o)^2$
10	165.9531	165.8233	2946.3597	0.0168
15	217.3516	218.7022	8.3054	1.8243
20	232.8695	229.3131	159.6690	12.6479
30	229.2956	231.8696	82.1229	6.6252
40	237.2076	231.9725	288.1211	27.4060
50	228.9845	231.9767	76.5811	8.9529
60	229.9724	231.9768	94.8464	4.0179

Time: Sample heating time

Observed: Observed peak area

Predicted: Predicted peak area

$[y - \text{mean}(y)]^2$: [Observed peak area - mean value of observed peak area]²

$(y_p - y_o)^2$: (Predicted peak area - Observed peak area)²

By using Solver function in Microsoft Excel, the values of parameters can be obtained.


Parameter	Value
C_e	231.9768
F	-6.08222
k_1	135.4867
k_2	0.321225

$$R^2 = 1 - \frac{\sum(y_p - y_o)^2}{\sum[y - \text{mean}(y)]^2}$$

$$R^2 = 0.9832$$

$$RMSE = \sqrt{\frac{\sum(y_p - y_o)^2}{\text{no. of data}}}$$

$$RMSE = 2.9639$$



A.6 Kinetic Study of 3,3-dimethylhexane

$$\text{Equation: } C_t = C_e[1 - [Fe^{-k_1t}] - [(1 - F)e^{-k_2t}]]$$

Time	Observed	Predicted	[y-mean(y)] ²	(y _p -y _o) ²
5	120.3245	110.4987	10082.4868	96.5450
10	127.6258	143.5464	8669.5256	253.4680
15	174.5429	171.0385	2133.8078	12.2807
40	253.1402	252.8815	1050.0298	0.0669
50	274.2744	269.5733	2866.3563	22.1000
80	296.3638	294.6511	5719.5525	2.9333
110	298.8808	302.9627	6106.6071	16.6617

Time: Sample heating time

Observed: Observed peak area

Predicted: Predicted peak area

[y-mean(y)]²: [Observed peak area-mean value of observed peak area]²

(y_p-y_o)²: (Predicted peak area-Observed peak area)²

By using Solver function in Microsoft Excel, the values of parameters can be obtained.

Parameter	Value
C_e	307.0831
F	0.230468
k_1	135.4867
k_2	0.036811

$$R^2 = 1 - \frac{\sum(y_p - y_o)^2}{\sum[y - \text{mean}(y)]^2}$$

$$R^2 = 0.9890$$

$$RMSE = \sqrt{\frac{\sum(y_p - y_o)^2}{\text{no. of data}}}$$

$$RMSE = 7.5975$$

