

REMOVAL OF ASPIRIN AND PARACETAMOL FROM AQUEOUS SOLUTION  
BY USING TEA WASTE MODIFIED PEI ADSORBENT

HII KHENG WUONG

A dissertation submitted in partial fulfilment of the  
requirements for the award of the degree of  
Master of Engineering

Faculty of Chemical and Energy Engineering  
Universiti Teknologi Malaysia

MARCH 2018

*Specially thanks to  
my family,  
my friends,  
and  
to all that involve in this master project....*

## **ACKNOWLEDGEMENT**

First, I would like to express my appreciation to my supervisor, Dr Norzita Ngadi for her outstanding supervision, constant support, and the time she put up for assisting my project work.

I would like to express my thanks to my fellow friend for their encouragement during this master project. I also would like to appreciate those who were directly or indirectly contributed to this project

At last, I would like to have a special thanks to my dearest family who supported and encouraged me in terms of financial aid and spiritual. Without their full support and encouragement, my master project and my life would not be complete.

Thank you

## ABSTRACT

The water bodies were determined to have been polluted by high concentration of pharmaceutical substances such as aspirin and paracetamol which may harm the ecosystem and bring health hazard to human. Conventional water treatment such as activated carbon used to remove these substances were not cost effective to be implemented. This study presented the alternative to the removal of aspirin and paracetamol from aqueous solution by using PEI modified tea waste adsorbent. The characteristics such as surface areas, pore volume and functional group of PEI-TW were identified by using Fourier Transform Infrared Spectroscopy (FTIR) and Brunauer-Emmett- Teller (BET). The total surface areas and pore volume of PEI-TW were 4.6748 m<sup>2</sup>/g and 0.0249 cm<sup>3</sup>/g, respectively. The highest removal percentage for aspirin was 87.34% in the optimum condition of initial concentration, dosage adsorbent; pH and temperature were 20 mg/L, 0.1g, pH 3 and temperature 60 °C, respectively. Whereas removal of paracetamol achieved the percentage of 99.70% in the optimum condition of initial concentration, dosage adsorbent; pH and temperature were 20 mg/L, 0.1g, pH 7 and temperature 60 °C, respectively. To describe the equilibrium isotherm for the adsorption of aspirin and paracetamol, the experimental data were fitted into Langmuir and Freundlich isotherm models, and both obeyed the Freundlich isotherm model with the correlation coefficient of 0.9865 and 0.9957 respectively. The adsorption process of aspirin and paracetamol obeyed the Pseudo-second order kinetic. The thermodynamic study indicated that the adsorption process of aspirin and paracetamol onto PEI modified tea waste adsorbent was feasible, spontaneous and endothermic. It can conclude that PEI modified tea waste adsorbent has high potential to be low-cost adsorbent and effective in the removal of paracetamol from aqueous solution compared to aspirin.

## ABSTRAK

Sumber air didapati mempunyai kandungan farmaseutikal seperti aspirin dan paracetamol yang tinggi dimana ia boleh mengganggu ekosistem dan memberi kesan kepada kesihatan manusia. Rawatan air konvensional seperti karbon diaktifkan yang berfungsi menyingkirkan bahan-bahan tersebut tidak begitu efektif dari segi kos. Kajian ini mencadangkan alternative dalam penyingkiran aspirin dan paracetamol dari larutan akueus menggunakan sisa teh yang dimodifikasi dengan polyethylenimine. Ciri-ciri seperti luas permukaan, isipadu dan kumpulan berfungsi di PEI-TW telah dikenal pasti dengan menggunakan Fourier Transform Infrared Spectroscopy (FTIR) dan Brunauer-Emmett Teller (BET). Jumlah luas permukaan dan isipadu PEI-TW adalah  $4.6748 \text{ m}^2/\text{g}$  dan  $0.0249 \text{ cm}^3/\text{g}$ . Penyingkiran aspirin yang tertinggi adalah 87.34% dan keadaan ini dicapai semasa dalam kondisi kepekatan awal 20 mg/L, dos penjerap 0.1g, pH 3 dan suhu 60 °C. Manakala untuk penyingkiran paracetamol, peratus tertinggi yang diperolehi adalah 99.70% dengan kondisi kepekatan awal 20 mg/L, dos penjerap 0.1g, pH 7 dan suhu 60 °C. Data eksperimen telah dianalisis dengan menggunakan model isotherm penjerapan: Langmuir dan Freundlich dan ia selaras dengan isoterma Freundlich yang memberikan koefisien korelasi sebanyak 0.9865 dan 0.9957 untuk penyingkiran aspirin dan paracetamol. Kadar penjerapan didapati mematuhi kinetik pseudo-tertib kedua. Kajian termodinamik menunjukkan bahawa penjerapan aspirin dan paracetamol boleh dilaksanakan, spontan dan endothermik. Dengan kesuluruhannya, keputusan menunjukkan bahawa PEI-TW boleh digunakan sebagai alternatif berkos rendah untuk penyingkiran paracetamol daripada larutan akueus jika dibandingkan dengan penyingkiran aspirin.

## TABLE OF CONTENTS

<b>CHAPTE R</b>	<b>TITLE</b>	<b>PAGE</b>
	<b>DECLARATION</b>	ii
	<b>DEDICATION</b>	iii
	<b>ACKNOWLEDGEMENT</b>	iv
	<b>ABSTRACT</b>	v
	<b>ABSTRAK</b>	vi
	<b>TABLE OF CONTENTS</b>	vii
	<b>LIST OF TABLES</b>	xi
	<b>LIST OF FIGURES</b>	xiii
	<b>LIST OF ABBREVIATIONS</b>	xvi
	<b>LIST OF SYMBOLS</b>	xviii
	<b>LIST OF APPENDICES</b>	xx
<b>1</b>	<b>INTRODUCTION</b>	1
	1.1 Background Study	1
	1.2 Problem Statement	3
	1.3 Objectives	4
	1.4 Research Scope	4
<b>2</b>	<b>LITERATURE REVIEW</b>	6
	2.1 Water Pollution	6
	2.1.1 Paracetamol	7

2.1.2	Aspirin	8
2.2	Tea Waste	9
2.3	Surface Modification	11
2.3.1	Polyehtylenimine (PEI)	13
2.4	Adsorption	15
2.4.1	Parameters Affecting Adsorption Process	15
2.4.1.1	Effect of Contact Time	16
2.4.1.2	Effect of Initial Substances Concentration	16
2.4.1.3	Effect of Temperature	17
2.4.1.4	Effect of Solution pH	18
2.4.1.5	Effect of Adsorbent Dosage	19
2.4.1.6	Summarize Influence of Parameters Affecting Adsorption Process	20
2.4.2	Adsorption Isotherm Models	21
2.4.2.1	Langmuir Adsorption Isotherm	21
2.4.2.2	Freundlich Adsorption Isotherm	22
2.4.3	Adsorption Kinetics	23
2.4.4	Thermodynamics Analysis	25
<b>3</b>	<b>RESEARCH METHODOLOGY</b>	<b>27</b>
3.1	Introduction	27
3.2	Materials	29
3.3	Preparation of PEI-TW Adsorbent	30
3.3.1	Pretreatment of Tea Waste	30
3.3.2	Modification of Tea Waste with PEI	30
3.4	Characterization of TW and PEI-TW Adsorbent	31
3.5	Procedure for Aqueous APAP and ASA Adsorption Experiment	32
3.5.1	Preparation of Aqueous APAP and ASA Solution	32
3.5.2	Batch Adsorption Experiment	32
3.5.2.1	Effect of Contact Time	33
3.5.2.2	Effect of Initial Substances Concentration	33

3.5.2.3	Effect of Temperature	33
3.5.2.4	Effect of Solution pH	34
3.5.2.5	Effect of Adsorbent Dosage	34
3.6	Adsorption Analysis	34
3.6.1	UV-Vis Spectroscopy	34
3.6.2	Analytical Technique in Determination of Substances Concentration	35
3.6.3	Preparation of Calibration Curve	35
3.6.4	Calculation of Substances Removal Percentage and Adsorption Capacity	35
<b>4</b>	<b>RESULTS AND DISCUSSIONS</b>	<b>37</b>
4.1	Introduction	37
4.2	Ratio Screening	38
4.3	Sample Characterization	38
4.3.1	Fourier Transform Infrared Spectroscopy (FTIR)	39
4.3.2	Brunauer-Emmett-Teller (BET)	40
4.4	Pharmaceutical Compound Performances	41
4.4.1	Adsorption Parameters	41
4.4.1.1	Effect of Contact Time	42
4.4.1.2	Effect of Initial Concentration	43
4.4.1.3	Effect of Temperature	46
4.4.1.4	Effect of Solution pH	48
4.4.1.5	Effect of Adsorbent Dosage	50
4.4.2	Adsorption Equilibrium Isotherms	52
4.4.3	Adsorption Kinetics	56
4.4.4	Thermodynamics Study	61
<b>5</b>	<b>CONCLUSION AND RECOMMENDATION</b>	<b>64</b>
5.1	Conclusion	64
5.2	Recommendation	66

<b>REFERENCES</b>	67
Appendix A-C	74

**LIST OF TABLES**

<b>TABLE NO</b>	<b>TITLE</b>	<b>PAGE</b>
2.1	Properties of Paracetamol	8
2.2	Properties of Aspirin	9
2.3	Chemical Composition of Tea Waste	10
2.4	Previous Studies in Surface Modification of Waste Adsorbent	12
2.5	Structure and Physicochemical Properties of PEI	13
2.6	Previous Studies in Substances Removal using PEI-modified Adsorbent	14
2.7	Summarized Influence of Parameters Affecting Adsorption	20
3.1	Chemicals and Reagents Used According to Experimental Scopes	29
3.2	Materials and Equipment According to Experimental Scopes	29
4.1	Adsorption of Aspirin and Paracetamol by Using PEI Modified Tea Waste Adsorbent with Different Ratio	38
4.2	Pore Characteristic of PEI-TW	41
4.3	The Adsorption Isotherm Parameters for Aspirin and Paracetamol Removal using PEI-TW Adsorbent	55
4.4	Adsorption Kinetic Parameter for Aspirin and Paracetamol Adsorption	59
4.5	Thermodynamics Study for the Adsorption of Aspirin and Paracetamol onto PEI-TW Adsorbent at Various Temperature	62

## LIST OF FIGURES

FIGURE NO	TITLE	PAGE
2.1	Efficiency of Pharmaceutical Substances Removal using Tea Waste as Adsorbent	11
3.1	Flowchart of Research Methodology	28
3.2	Flowchart of Procedure for Pretreatment of Tea Waste	30
3.3	Flowchart of Procedure for Modification of Tea Waste PEI	31
4.1	FTIR Spectrum of PEI Modified Tea Waste with Ratio PEI:TW of 1:1, 1:2 and 2:1	39
4.2	Profile for the Surface Area of PEI-TW	40
4.3	Effect of Contact Time on Adsorption of Aspirin and Paracetamol. Experiment Condition: Volume of Substance Solution: 20 ml, C <sub>0</sub> = 100 mg/L of Aspirin and Paracetamol Solutions: Mass of Adsorbent: 0.1g; Contact Time: 180 min, Room Temperature; pH = 7 for Paracetamol, pH = 4 for Aspirin; Agitation Rate: 200rpm	43
4.4	Effect of Initial Concentration on Adsorption of Aspirin. Experiment Condition: C <sub>0</sub> = 20 to 100 mg/L of Aspirin Solution: Mass of Adsorbent: 0.1g; Contact Time: 60 mins, Room Temperature; pH = 7; Agitation Rate: 200 rpm	45
4.5	Effect of Initial Concentration on Adsorption of Paracetamol. Experiment Condition: C <sub>0</sub> = 20 to 100 mg/L of Paracetamol Solution: Mass of Adsorbent: 0.1g; Contact Time: 60 mins, Room Temperature; pH = 7; Agitation Rate: 200 rpm	45
4.6	Effect of Temperature on Adsorption of Aspirin. Experiment Condition: C <sub>0</sub> = 20 mg/L of Aspirin Solution: Mass of Adsorbent: 0.1g; Contact time:	47

	60 mins, Temperature = 30 to 60 °C; pH = 7; Agitation Rate: 200 rpm	
4.7	Effect of Temperature on Adsorption of Paracetamol. Experiment Condition: C <sub>0</sub> = 20 mg/L of Paracetamol Solution: Mass of adsorbent: 0.1g; Contact Time: 60 mins, Temperature = 30 to 60 °C; pH = 7; Agitation Rate: 200 rpm	47
4.8	Effect of pH on Adsorption of Aspirin. Experiment Condition: C <sub>0</sub> = 20 mg/L of Aspirin Solution: Mass of Adsorbent: 0.1g; Contact time: 60 mins, Temperature: 60 °C; pH = 3 to 7; agitation rate: 200 rpm	49
4.9	Effect of pH on Adsorption of Paracetamol. Experiment Condition: C <sub>0</sub> = 20 mg/L of Paracetamol Solution: Mass of Adsorbent: 0.1g; Contact time: 60 mins, Temperature: 60 °C; pH = 3 to 7; Agitation Rate: 200 rpm	49
4.10	Effect of Adsorbent Dosage on Adsorption of Aspirin. Experiment Condition: C <sub>0</sub> = 20 mg/L of Aspirin Solution: Mass of Adsorbent: 0.1 to 1.0g; Contact Time: 60 mins, Temperature: 60 °C; pH = 3; Agitation Rate: 200 rpm	51
4.11	Effect of Adsorbent Dosage on Adsorption of Paracetamol. Experiment Condition: C <sub>0</sub> = 20 mg/L of Paracetamol Solution: Mass of Adsorbent: 0.1 to 1.0g; Contact Time: 60 mins, Temperature: 60 °C; pH = 7; Agitation Rate: 200 rpm	51
4.12	Langmuir Adsorption Isotherm for Aspirin onto PEI-TW Adsorbent	53
4.13	Langmuir Adsorption Isotherm for Paracetamol onto PEI-TW Adsorbent	53
4.14	Freundlich Adsorption Isotherm for Aspirin onto PEI-TW Adsorbent	54
4.15	Freundlich Adsorption Isotherm for Paracetamol onto PEI-TW Adsorbent	54
4.16	Pseudo First Order for Adsorption of Aspirin onto PEI-TW Adsorbent	57
4.17	Pseudo First Order for Adsorption of Paracetamol onto PEI-TW Adsorbent	57
4.18	Pseudo Second Order for Adsorption of Aspirin onto PEI-TW Adsorbent	58
4.19	Pseudo Second Order for Adsorption of Paracetamol onto PEI-TW Adsorbent	58

4.20	Intraparticle Diffusion for Adsorption of Aspirin onto PEI-TW Adsorbent	60
4.21	Intraparticle Diffusion for Adsorption of Paracetamol onto PEI-TW Adsorbent	60
4.22	Thermodynamic Study for Adsorption of Aspirin onto PEI-TW Adsorbent	62
4.23	Thermodynamic Study for Adsorption of Paracetamol onto PEI-TW Adsorbent	63

**LIST OF ABBREVIATIONS**

AB45	-	Acid Blue 45
AO8	-	Acid Orange 8
APAP	-	Paracetamol
ARGs	-	Antibiotic Resistance Genes
ASA	-	Aspirin
BET	-	Brunauer-Emmett-Teller
C	-	Carbon
C <sub>8</sub> H <sub>8</sub> O <sub>4</sub>	-	Aspirin
C <sub>8</sub> H <sub>9</sub> NO <sub>2</sub>	-	Paracetamol
CBZ	-	Carbamazepine
CPC	-	Cetylpyridium Chloride
CR	-	Congo Red
CTAB	-	Cetyltrimethyl Ammonium Bromide
Cu	-	Copper
Fe/N-CNT/ $\beta$ -CD	-	Iron and Nitrogen Base Nanocomposite Polymer
Fe <sub>3</sub> O <sub>4</sub> NPs	-	Magnetite Nanoparticles
FTIR	-	Fourier Transform Infra-Red
H	-	Hydrogen
HCL	-	Hydrochloric Acid
IUPAC	-	International Union of Pure and Applied Chemistry
KNO <sub>3</sub>	-	Potassium Nitrate
MGEs	-	Mobile Genetic Elements
N	-	Nitrogen
NaOH	-	Sodium Hydroxide

NaOH	-	Sodium Hydroxide
N-CNT/ $\beta$ -CD	-	Nitrogen Base Nanocomposite Polymer
NO+SO <sub>2</sub>	-	Pyrolysed Waste Tea Char
O	-	Oxygen
OH	-	Hydroxyl Group
PACs	-	Pharmaceutically Active Compounds
Pb	-	Lead
PCA	-	Paracetamol
PEI	-	Polyethylenimine
PEI-TW	-	Tea Waste Modified using PEI
SMX	-	Sulfamethoxazole
TMP	-	Trimethoprim
TW	-	Tea Waste
UV	-	Ultra Violet
ZnCl <sub>2</sub>	-	Zinc Chloride

## LIST OF SYMBOLS

Ce	-	Equilibrium concentration
Cm <sup>3</sup> /g	-	Centimeter cubed per gram
Co	-	Initial concentration
g	-	Gram
g/L	-	Gram per litre
g/mol	-	Gram per mol
hr (s)	-	Hour (s)
µg/L	-	Microgram per litre
µm	-	Micro Meter
g/mol	-	Gram/mole
g/cm <sup>3</sup>	-	Gram per cubic centimetre
J	-	Joule
K	-	Kelvin
kJ	-	Kilo Joule
k1	-	Adsorption rate constant of first order adsorption
k2	-	Adsorption rate constant of second order adsorption
K <sub>F</sub>	-	Freundlich constant
kJ/mol	-	Kilo Joule per mol
K <sub>L</sub>	-	Langmuir constants related to the rate of adsorption
kp	-	Intraparticle diffusion rate constant
kV	-	Kilovolt
L	-	Liter
M	-	Molar
m <sup>2</sup> /g	-	Meter square per gram

mg	-	Milligram
mg/g	-	Milligram per gram
mL	-	Mililitre
mg/L	-	Milligram per litre
min	-	Minute
mm	-	Milimetre
ng/L	-	Nanograms/litre
nm	-	Nanometre
$q_e$	-	Amount of adsorbed on adsorbent at equilibrium
$q_m$	-	Maximum adsorption capacity
$q_t$	-	Amount of adsorbed on adsorbent at any time
R	-	Universal gas constant
$R^2$	-	Regression Coefficient
$R_L$	-	Separation factor
% w/v	-	weight/volume percent
$\Delta G^\circ$	-	Gibbs energy changes
$\Delta H^\circ$	-	Enthalpy change
$\Delta S^\circ$	-	Entropy change
$^\circ\text{C}$	-	Degree celcius
%	-	Percentage
<	-	Less than
>	-	Greater than

**LIST OF APPENDICES**

<b>APPENDIX</b>	<b>TITLE</b>	<b>PAGE</b>
A	Spectrum Profile of Aspirin and Paracetamol in Various Concentrations	84
B	Aspirin and Paracetamol Calibration Curve	85
C	Calculation	86

## CHAPTER 1

### INTRODUCTION

#### 1.1 Background Study

Water pollution is a major environmental concern that is caused by pollutants that are discharged to water stream from sources with inadequate water treatments. Industrial wastewater is apparently one of the main sources that released these harmful compounds, also known as pollutants. There are many types of pollutants that originate from industrial wastewater effluents and one of them is antibiotics. Antibiotics are extensively utilised in different type of industries, especially in pharmaceutical and personal care products. Up to 100  $\mu\text{g L}^{-1}$  of various pharmaceuticals were found in aqueous environment, soil and mostly in effluents from drug manufactures (Fakhri and Behrouz, 2015). The high concentration of antibiotics existing in industrial water is a troubling issue as it can cause several health effects to human and aquatic life.

Large amount of antibiotics such as antibiotic resistance genes (ARGs) and mobile genetic elements (MGEs) are harbour from pharmaceutical waste. It can increase the prevalence of single- and multiple-antibiotic resistance among *Acinetobacter* species in wastewater treatment plant, which may affect the efficiency of antibiotic removal by anaerobic digestion (Liu *et al.*, 2014). These antibiotics

present in water bodies are considered toxic and harmful as it can cause several health effects to human and aquatic life due to it assisting pathogen to be antibiotic-immune, making it hard to remove or destroy the pathogen (Romero, 2012).

Various low-cost materials especially natural or wasted materials have been extensively explored for adsorption removal of antibiotics from aqueous solution. Commonly used natural raw materials such as coal, wood, pit shells, coconut and nut shells are used to manufacture absorbents. One of the commonly known natural waste is tea waste. The large consumption of tea is found in Turkey, which is the world's largest black tea producing country (Gondogdu *et al.*, 2013). It leads to a significant amounts of tea waste being generated. Therefore, an efficient utilization of tea wastes has been gaining considerable attention and absorbent produce from tea waste will be used to remove antibiotics in industrial wastewater.

For improving the affinity of a material to pharmaceutical substances, various cationic surface modifications have been explored and studied. Polyethylenimine (PEI) is a relatively cheap cationic polymer that can be used for surface modification of an adsorbent. It has been proven in previous researches that it has the capability to enhance the removal of other anionic pollutant such as anionic metals and dyes (Deng and Ting, 2005; Won *et al.*, 2011). This polymer can readily bind with anionic substrate. Therefore, with the surface modification of tea waste, adsorption of pharmaceutical substances using tea waste can be widely implemented in industries in the future to remove various classes of pharmaceutical substances from wastewater.

## 1.2 Problem Statement

The presence of paracetamol and aspirin in water bodies could bring adverse effect to the human health and the environment, even in a low concentration (<0.1 mg/L). It could disrupt the biological function of micro-organism in wastewater treatment plant, reducing its efficiency in removal, furthermore promoting resistance development to the pathogen, which were harmful to human health (Larsson *et al.*, 2014).

Adsorption method was considered as the most effective wastewater treatments process to reduce the hazardous material in the wastewater. Nowadays, the adsorbent that was widely employed for adsorption in industry was commercial activated carbon. However, its production method was limited and costly due to the usage of strong acidic activating agent. Therefore, it was vital that an alternative material which was cost-effective as well as abundant to be used as the adsorbent for substances removal from wastewater (Yagub *et al.*, 2014).

Tea waste which was regarded as an inexpensive and abundant, obtainable at the amount of 30 000 tons from the extraction process of instant tea manufacturing (Malkoc and Nuhoglu, 2006) and the final residues originated from cafeteria which could act as a promising adsorbent for antibiotic removal (Gundogdu *et al.*, 2013). It has been proven that plant based materials appeared to be the best raw material and were suitable for large-scale production as they are abundant (Sebastian, 2017). Hence, the removal efficiency of antibiotics for tea waste adsorbent can be enhanced through a surface modification on the tea waste. Polyethylenimine (PEI) was a cationic polymer that was proved to be exemplary in its surface modification role for the removal of anionic pollutant such as anionic metals and dyes (Deng and Ting, 2005; Low *et al.*, 2008, Won *et al.*, 2011; Sadaf *et al.*, 2014). This was due to the significant amount of positively charged amine groups from PEI introduced to the surface of adsorbent in which it could readily bind to anionic substrate. Thus, tea waste impregnated with PEI could be used as a potential adsorbent for antibiotic removal from aqueous solution.

### 1.3 Objectives

The objectives of this research are:

- i) To synthesize and characterize PEI-modified tea waste as the adsorbent for paracetamol and aspirin removal.
- ii) To investigate the effect of the physicochemical parameters such as pH, temperature, contact time, initial paracetamol and aspirin concentration and adsorbent dosage on paracetamol and aspirin adsorption performance of PEI-modified tea waste.
- iii) To analyse the adsorption behaviour of PEI-modified tea waste using isotherm, kinetic, and thermodynamic analysis.

### 1.4 Research Scope

Paracetamol and aspirin for this study will be prepared in a solution and the adsorbent used were from natural waste material which was known to be inexpensive and abundant. In addition to that, the type of material used to modify the surface of the adsorbent was a cationic polymer. Moreover, the characterization of the adsorbent was conducted using analytical instrumentation, which was Fourier Transform Infrared (FTIR) Spectrophotometer and Brunauer-Emmett-Teller (BET).

Furthermore, this study was done under various parameters which were pH, temperature, contact time, initial substances concentration and adsorbent dosage. The range of the solution pH that the adsorption study was carried out is in the range of pH 2 to pH 9. As for operational temperature, the adsorption study was conducted in the range of 30°C to 60°C. For initial paracetamol and aspirin concentration, the adsorption process was conducted with initial concentration between 0.02 g/L to 0.1 g/L. Also, the adsorbent dosage that will be used in this study were in the range of 0.1 to 1.0 g. For the equilibrium study, the Langmuir and Freundlich isotherms were

applied. The Pseudo first-order, Pseudo second-order and intraparticle diffusion were applied for the kinetic study. The thermodynamics parameters of the adsorption of acetaminophen including the Gibbs free energy ( $\Delta G$ ), enthalpy ( $\Delta H$ ), entropy ( $\Delta S$ ) were calculated by using Van't Hoff equation.

## REFERENCES

- Ali, H. (2010). Biodegradation of Synthetic Dyes—A Review. *Water, Air, & Soil Pollution*, 213(1-4), 251-273.
- Al-Khateeb, L. A., Almotiry, S., & Salam, M. A. (2014). Adsorption of pharmaceutical pollutants onto graphene nanoplatelets. *Chemical Engineering Journal*, 248, 191-199. doi:10.1016/j.cej.2014.03.023
- Al-Khateeb, L. A., *et al.* 2014. Adsorption of Pharmaceutical Pollutants onto Graphene Nanoplatelets. *Chemical Engineering Journal* 248(191-199).
- Amarasinghe, B., & Williams, R. (2007). Tea waste as a low cost adsorbent for the removal of Cu and Pb from wastewater. *Chemical Engineering Journal*, 132(1-3), 299-309. doi:10.1016/j.cej.2007.01.016
- André, V., Piedade, M. F., & Duarte, M. T. (2012). Revisiting paracetamol in a quest for new co-crystals. *CrystEngComm*, 14(15), 5005. doi:10.1039/c2ce25307k
- Ansari, R., Seyghali, B., Mohammad-khah, A., & Zanjanchi, M. (2012). Highly Efficient Adsorption of Anionic Dyes from Aqueous Solutions Using Sawdust Modified by Cationic Surfactant of Cetyltrimethylammonium Bromide. *Journal Of Surfactants And Detergents*, 15(5), 557-565.
- Ayas, N., & Esen, T. (2016). Hydrogen production from tea waste. *International Journal of Hydrogen Energy*, 41(19), 8067-8072. doi:10.1016/j.ijhydene.2015.09.156
- Aziz, R., Chiban, M., Eddaoudi, E., Soudani, A., Zerbet, M., & Sinan, F. (2016). Factors controlling the adsorption of acid blue 113 dye from aqueous solution by dried *C. edulis* plant as natural adsorbent. *Arabian Journal Of Geosciences*, 9(15).

- Banerjee, S. & Chattopadhyaya, M. (2013). Adsorption characteristics for the removal of a toxic dye, tartrazine from aqueous solutions by a low cost agricultural by-product. *Arabian Journal Of Chemistry*.
- Chen, H., Zhao, J., Wu, J., & Dai, G. (2011). Isotherm, thermodynamic, kinetics and adsorption mechanism studies of methyl orange by surfactant modified silkworm exuviae. *Journal Of Hazardous Materials*.
- Clissold, S. P. (1986). Paracetamol and Phenacetin. *Drugs*,32(Supplement 4), 46-59. doi:10.2165/00003495-198600324-00005
- Demiral, I. & Samdan, C. A. 2016. Preparation and Characterisation of Activated Carbon from Pumpkin Seed Shell Using H<sub>3</sub>PO<sub>4</sub>. *Anadolu University Journal of Science and Technology-Applied Sciences and Engineering* 17(1): 125-138.
- Deng, S., & Ting, Y. (2005). Characterization of PEI-modified biomass and biosorption of Cu(II), Pb(II) and Ni(II). *Water Research*,39(10), 2167-2177. doi:10.1016/j.watres.2005.03.03
- Deng, S., & Ting, Y. (2005). Characterization of PEI-modified biomass and biosorption of Cu(II), Pb(II) and Ni(II). *Water Research*, 39(10), 2167-2177.
- Dieng, H., Zawawi, R. B., Nur Intan Saidaah Bt Mohamed Yusof, Ahmad, A. H., Abang, F., Ghani, I. A., . Noweg, G. T. (2016). Green tea and its waste attract workers of formicine ants and kill their workers—implications for pest management. *Industrial Crops and Products*, 89, 157-166. doi:10.1016/j.indcrop.2016.05.019
- Dutta, M., et al. 2015. Adsorption of Acetaminophen by Using Tea Waste Derived Activated Carbon. *International Journal of Environmental Sciences* 6(2):
- Fakhri and Behrouz, A. Fakhri, S. Behrouz (2015). Comparison studies of adsorption properties of MgO nanoparticles and ZnO–MgO nanocomposites for linezolid antibiotic removal from aqueous solution using response surface methodology, *Process Saf. Environ. Prot.*, 94 (2015), pp. 37–43
- Foroughi-dahr, M., Abolghasemi, H., Esmaili, M., Nazari, G., & Rasem, B. (2015). Experimental study on the adsorptive behavior of Congo red in cationic surfactant-modified tea waste. *Process Safety And Environmental Protection*, 95, 226-236.

- Granberg, R. A., & Rasmuson, Å C. (1999). Solubility of Paracetamol in Pure Solvents. *Journal of Chemical & Engineering Data*, 44(6), 1391-1395. doi:10.1021/je990124v
- Gulipalli, C. S., Prasad, B. & Wasewar, K. L. 2011. Batch Study, Equilibrium and Kinetics of Adsorption of Selenium Using Rice Husk Ash (Rha). *J. Eng. Sci. Technol* 6(5): 586-605.
- Gundogdu, A., Duran, C., Senturk, H. B., Soylak, M., Imamoglu, M., & Onal, Y. (2013). Physicochemical characteristics of a novel activated carbon produced from tea industry waste. *Journal of Analytical and Applied Pyrolysis*, 104, 249-259. doi:10.1016/j.jaap.2013.07.008
- Halling-Sørensen, B., Nors, S., Lanzky, P. F., Ingerslev, F., Holten, H. C., & Jørgensen, S. E. (1998, January). Occurrence, fate and effects of pharmaceutical substances in the environment, 36:357-393. Retrieved May 30, 2017
- Hameed, B. H. 2009. Spent Tea Leaves: A New Non-Conventional and Low Cost Adsorbent for Removal of Basic Dye from Aqueous Solution. *Journal of Hazardous Materials* 161(2-3): 753-759.
- Hari, A. C., Paruchuri, R. A., Sabatini, D. A., & Kibbey, T. C. (2005). Effects of pH and Cationic and Nonionic Surfactants on the Adsorption of Pharmaceuticals to a Natural Aquifer Material. *Environmental Science & Technology*, 39(8), 2592-2598. doi:10.1021/es048992m
- Humpola, P., Odetti, H., Fertitta, A., & Vicente, J. (2013). Thermodynamic analysis of adsorption models of Phenol in liquid phase on different activated carbons. *Journal Of The Chilean Chemical Society*, 58(1), 1541-1544.
- Kilic, M., *et al.* 2011. Adsorptive Removal of Phenol from Aqueous Solutions on Activated Carbon Prepared from Tobacco Residues: Equilibrium, Kinetics and Thermodynamics. *Journal of Hazardous Materials* 189(1-2): 397-403.
- Kozer, E., & Koren, G. (2001). Management of Paracetamol Overdose. *Drug Safety*, 24(7), 503-512. doi:10.2165/00002018-200124070-00003
- Lafi, R., & Hafiane, A. (2016). Removal of methyl orange (MO) from aqueous solution using cationic surfactants modified coffee waste (MCWs). *Journal Of The Taiwan Institute Of Chemical Engineers*, 58, 424-433.

- Lafi, R., Fradj, A., Hafiane, A., & Hameed, B. (2014). Coffee waste as potential adsorbent for the removal of basic dyes from aqueous solution. *Korean Journal Of Chemical Engineering*, 31(12), 2198-2206.
- Larsson, D. J., Pedro, C. D., & Paxeus, N. (2007). Effluent from drug manufactures contains extremely high levels of pharmaceuticals. *Journal of Hazardous Materials*, 148(3), 751-755. doi:10.1016/j.jhazmat.2007.07.008
- Li, N., Bai, R., 2005. A novel amine-shielded surface cross-linking of chitosan hydrogel beads for enhanced metal adsorption performance. *Ind. Eng. Chem.Res.* 44, 6692–6700.
- Lidstone, J. (2005). Aspirin: The Remarkable Story of a Wonder Drug. *Journal of Medical Marketing*, 5(2), 183-184. doi:10.1057/palgrave.jmm.5040227
- Liu, H., *et al.* 2013. Evaluation of Animal Hairs-Based Activated Carbon for Sorption of Norfloxacin and Acetaminophen by Comparing with Cattail Fiber-Based Activated Carbon. *Journal of Analytical and Applied Pyrolysis* 101(156-165).
- Liu, M., Ding, R., Zhang, Y., Gao, Y., Tian, Z., Zhang, T., & Yang, M. (2014). Abundance and distribution of Macrolide-Lincosamide-Streptogramin resistance genes in an anaerobic-aerobic system treating spiramycin production wastewater. *Water Research*, 63, 33-41. doi:10.1016/j.watres.2014.05.045
- Low, B., Ting, Y., & Deng, S. (2008). Surface modification of *Penicillium chrysogenum* mycelium for enhanced anionic dye removal. *Chemical Engineering Journal*, 141(1-3), 9-17.
- Malkoc, E., & Nuhoglu, Y. (2006). Removal of Ni(II) ions from aqueous solutions using waste of tea factory: Adsorption on a fixed-bed column. *Journal of Hazardous Materials*, 135(1-3), 328-336. doi:10.1016/j.jhazmat.2005.11.070
- Mashayekh-Salehi, A. & Moussave, G. 2015. Removal of Acetaminophen from the Contaminated Water Using Adsorption onto Carbon Activated with  $NH_4Cl$ . *Desalination and Water Treatment* 10(1-13).
- Mphahlele, K., Onyango, M. S., & Mhlanga, S. D. (2015). Adsorption of aspirin and paracetamol from aqueous solution using Fe/N-CNT/ $\beta$ -cyclodextrin nanocomposites synthesized via a benign microwave assisted method. *Journal of Environmental Chemical Engineering*, 3(4), 2619-2630. doi:10.1016/j.jece.2015.02.018

- Mukoko, T., *et al.* 2015. Preparation of Rice Hull Activated Carbon for the Removal of Selected Pharmaceutical Waste Compound in Hospital Effluent. *Environmental & Analytical Toxicology* 7
- Nielsen, L., & Bandosz, T. J. (2016). Analysis of the competitive adsorption of pharmaceuticals on waste derived materials. *Chemical Engineering Journal*, 287, 139-147. doi:10.1016/j.cej.2015.11.016
- Ofomaja, A. & Ho, Y. (2007). Equilibrium sorption of anionic dye from aqueous solution by palm kernel fibre as sorbent. *Dyes And Pigments*, 74(1), 60-66.
- Pan, J., Ge, X., Liu, R., Tang, H., 2006. Characteristic features of Bacillus cereus cell surfaces with biosorption of Pb(II) ions by AFM and FT-IR. *Colloid. Surf. B* 52, 89–95.
- Pirbazari, A. E., & Pargami, N. R. (2015). Surfactant-coated Tea Waste: Preparation, Characterization and its Application for Methylene Blue Adsorption from Aqueous Solution. *Journal of Environmental & Analytical Toxicology*, 05(05). doi:10.4172/2161-0525.1000310
- Pirbazari, A. E., Saberikhah, E., Badrouh, M., & Emami, M. S. (2014). Alkali treated Foumanat tea waste as an efficient adsorbent for methylene blue adsorption from aqueous solution. *Water Resources and Industry*, 6, 64-80. doi:10.1016/j.wri.2014.07.003
- Pradhan, S., Singh, S., Rai, L.C., 2007. Characterization of various functional groups present in the capsule of Microcystis and study of their role in biosorption of Fe, Ni and Cr. *Bioresour. Technol.* 98, 595–601.
- Qiu, H., Lv, L., Pan, B., Zhang, Q., Zhang, W., & Zhang, Q. (2009). Critical review in adsorption kinetic models. *Journal Of Zhejiang University-SCIENCE A*, 10(5), 716-724.
- R, S. D., & Sebastian Antony Selvan C. (2017). Greener synthesis and characterization of silver nanoparticles using murraya koenigii leaf extract and its antibacterial activity. *International Journal of Pharma and Bio Sciences*, 8(2). doi:10.22376/ijpbs.2017.8.2.p292-298
- Rao, P.S., Smitha, B., Sridhar, S., Krishnaiah, A., 2006. Preparation and performance of poly(vinyl alcohol)/ polyethyleneimine blend membranes for the dehydration of 1,4-dioxane by pervaporation: comparison with glutaraldehyde cross-linked membranes. *Sep. Purif. Technol.* 48, 244–254.

- Reddy, D. H., Sessaiah, K., Reddy, A., & Lee, S. (2012). Optimization of Cd(II), Cu(II) and Ni(II) biosorption by chemically modified *Moringa oleifera* leaves powder. *Carbohydrate Polymer* 88(3), 1077-1086. doi:10.1016/j.carbpol.2012.01.07
- Revest, P. (2009). Introduction. *Medical Sciences*, 1-14. doi:10.1016/b978-0-7020-2679-9.00001-7
- Romero, J., Gloria, C., & Navarrete, P. (2012). Antibiotics in Aquaculture – Use, Abuse and Alternatives. *Health and Environment in Aquaculture*. doi:10.5772/28157
- Sadaf, S., Bhatti, H., Arif, M., Amin, M., & Nazar, F. (2015). Adsorptive removal of direct dyes by PEI-treated peanut husk biomass: Box–Behnken experimental design. *Chemistry And Ecology*, 31(3), 252-264.
- Sajab, M., Chia, C., Zakaria, S., & Khiew, P. (2013). Cationic and anionic modifications of oil palm empty fruit bunch fibers for the removal of dyes from aqueous solutions. *Bioresource Technology*, 128, 571-577.
- Salleh, M., Mahmoud, D., Karim, W., & Idris, A. (2011). Cationic and anionic dye adsorption by agricultural solid wastes: A comprehensive review. *Desalination*, 280(1-3), 1-13.
- Shen, B., Tian, L., Li, F., Zhang, X., Xu, H., & Singh, S. (2017). Elemental mercury removal by the modified bio-char from waste tea. *Fuel*, 187, 189-196. doi:10.1016/j.fuel.2016.09.059
- Soni, H. (2014). Martindale: The Complete Drug Reference Brayfield Alison (Ed) Martindale: The Complete Drug Reference £459 4,688pp Pharmaceutical Press 9780857111395 0857111396. *Emergency Nurse*, 22(5), 12-12. doi:10.7748/en.22.5.12.s13
- Stan, M., Lung, I., Soran, M., Leostean, C., Popa, A., Stefan, M., Porav, A. S. (2017). Removal of antibiotics from aqueous solutions by green synthesized magnetite nanoparticles with selected agro-waste extracts. *Process Safety and Environmental Protection*, 107, 357-372. doi:10.1016/j.psep.2017.03.003
- Subbaiah, M. & Kim, D. (2016). Adsorption of methyl orange from aqueous solution by aminated pumpkin seed powder: Kinetics, isotherms, and thermodynamic studies. *Ecotoxicology And Environmental Safety*, 128, 109-117.
- Sumathi, T. & Alagumuthu, G. 2014. Adsorption Studies for Arsenic Removal Using Activated *Moringa Oleifera*. *International Journal of Chemical*

- Engineering 2014 Adams, C., Wang, Y., Loftin, K. & Meyer, M. 2002. Removal of Antibiotics from Surface and Distilled Water in Conventional Water Treatment Processes. *Journal of environmental engineering* 128(3): 253-260.
- Weber, W. J. & Morris, J. C. 1963. Kinetics of Adsorption on Carbon from Solution. *Journal of the Sanitary Engineering Division* 89(2): 31-60.
- Won, S., Park, J., Mao, J., & Yun, Y. (2011). Utilization of PEI-modified *Corynebacterium glutamicum* biomass for the recovery of Pd(II) in hydrochloric solution. *Bioresource Technology*, 102(4), 3888-3893.
- Yagub, M., Sen, T., Afroze, S., & Ang, H. (2014). Dye and its removal from aqueous solution by adsorption: A review. *Advances In Colloid And Interface Science*, 209, 172-184.
- Zhang, R., Zhang, J., Zhang, X., Dou, C., & Han, R. (2014). Adsorption of Congo red from aqueous solutions using cationic surfactant modified wheat straw in batch mode: Kinetic and equilibrium study. *Journal Of The Taiwan Institute Of Chemical Engineers*, 45(5), 2578-2583.
- Zhao, B., Xiao, W., Shang, Y., Zhu, H., & Han, R. (2014). Adsorption of light green anionic dye using cationic surfactant-modified peanut husk in batch mode. *Arabian Journal Of Chemistry*.