



# Draft Genome Sequence of *Roseovarius* sp. PS-C2, Isolated from Sekinchan Beach in Selangor, Malaysia

 Nurfatini Radzlin,<sup>a,b</sup>  Suhaila Mohd Omar,<sup>c,d</sup>  Kok Jun Liew,<sup>e</sup>  Kian Mau Goh,<sup>e</sup>  Iffah Izzati Zakaria,<sup>a</sup>  
 Ummirul Mukminin Kahar<sup>a</sup>

<sup>a</sup>Malaysia Genome Institute, National Institutes of Biotechnology Malaysia, Kajang, Selangor, Malaysia

<sup>b</sup>Department of Biochemistry, Faculty of Biotechnology and Biomolecular Sciences, Universiti Putra Malaysia, Serdang, Selangor, Malaysia

<sup>c</sup>Department of Biotechnology, Kulliyah of Science, International Islamic University Malaysia, Kuantan, Pahang, Malaysia

<sup>d</sup>Research Unit for Bioinformatics and Computational Biology, Kulliyah of Science, International Islamic University Malaysia, Kuantan, Pahang, Malaysia

<sup>e</sup>Faculty of Science, Universiti Teknologi Malaysia, Skudai, Johor, Malaysia

Iffah Izzati Zakaria and Ummirul Mukminin Kahar contributed equally to this work. Author order was determined on the basis of seniority.

**ABSTRACT** *Roseovarius* sp. PS-C2 is a bacterium that was isolated from Sekinchan Beach in Selangor, Malaysia, using an *ex situ* cultivation technique. Here, we present a high-quality annotated draft genome of strain PS-C2 and suggest potential applications of this bacterium.

*Roseovarius* spp. are halophilic or halotolerant bacteria that thrive in various marine habitats (1). These bacteria have been reported as potential candidates for bioremediation and quorum sensing (2, 3).

*Roseovarius* sp. PS-C2 was isolated using a modified *ex situ* cultivation technique (4). Wet sediment and mud (0 to 15 cm from the top layer) were collected, using a sterilized laboratory scoop, from Sekinchan Beach in Selangor, Malaysia (3.5029N, 101.0945E). A sterilized corn cob was inserted into a 3-liter bioreactor filled with the collected sediment. After 2 weeks of incubation at 30°C and 80 rpm, samples were taken from the corn cob and spread onto marine agar (Condalab, Madrid, Spain). The PS-C2 strain was isolated, and genomic DNA was extracted using the Monarch genomic DNA purification kit (New England Biolabs, Ipswich, MA, USA), according to the manufacturer's instructions. The 16S rRNA gene was amplified by PCR using 27F and 1492R primers (5), sequenced, and taxonomically identified using the NCBI BLASTn search and the EzBioCloud 16S database (6). The analyses revealed that PS-C2 showed greatest similarity (99.93%) to *Roseovarius pacificus* 81-2<sup>T</sup> (GenBank accession number [DQ120726.1](https://doi.org/10.1128/MRA.00673-21)). Here, we present the genome sequence and analysis of *Roseovarius* sp. PS-C2. The high-quality annotated draft genome of PS-C2 might provide insights into its potential biotechnological applications.

*Roseovarius* sp. PS-C2 was grown on marine agar (pH 6.5) at 30°C for 24 h. Subsequently, bacterial genomic DNA was extracted using the Monarch genomic DNA purification kit, according to the manufacturer's protocol. A paired-end library was prepared using the standard protocol of the NEBNext Ultra DNA library preparation kit for Illumina (New England BioLabs). Sequencing was performed using the NovaSeq 6000 system with 150-bp paired-end reads (Illumina, San Diego, CA, USA). Sequence adaptors and low-quality reads were filtered using Trimmomatic v0.40 (7). *De novo* assembly of the genome was performed using SOAPdenovo v2.04 (8). The genome was analyzed and annotated using the NCBI Prokaryotic Genome Annotation Pipeline (PGAP) v5.20 (9). Genome comparison between strain PS-C2 and all 75 available genomes of *Roseovarius* spp. in the NCBI genome database (August 2021) was carried out using digital DNA-DNA hybridization (dDDH) via the Genome-to-Genome Distance Calculator (GGDC) v2.1 (10) and the average nucleotide identity (ANI) function in the EzBioCloud server (11). Carbohydrate-active enzymes (CAZymes) present in the genome of the PS-C2 strain were mined using dbCAN2 (12). Default parameters were used for all software tools, unless otherwise specified.

**Citation** Radzlin N, Mohd Omar S, Liew KJ, Goh KM, Zakaria II, Kahar UM. 2021. Draft genome sequence of *Roseovarius* sp. PS-C2, isolated from Sekinchan Beach in Selangor, Malaysia. *Microbiol Resour Announc* 10:e00673-21. <https://doi.org/10.1128/MRA.00673-21>.

**Editor** Catherine Putonti, Loyola University Chicago

**Copyright** © 2021 Radzlin et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Iffah Izzati Zakaria, [iffahizzati@nibm.my](mailto:iffahizzati@nibm.my), or Ummirul Mukminin Kahar, [ummirul@nibm.my](mailto:ummirul@nibm.my).

**Received** 2 July 2021

**Accepted** 22 August 2021

**Published** 23 September 2021

The sequencer generated 1,748,493,900 bases from 5,828,313 paired-end reads. The *Roseovarius* sp. PS-C2 genome showed coverage of 362×. The genome was assembled into 39 contigs, with a total size of 3,958,657 bp, an  $N_{50}$  value of 411,343 bp, and a G+C content of 62.1%. The PS-C2 genome consisted of 3,920 predicted genes, of which 3,839 were protein-coding sequences, 31 were pseudogenes, 3 were rRNAs, 44 were tRNAs, and 3 were small RNAs. The genome comparison analyses showed that PS-C2 was closely related to *Roseovarius pacificus* 81-2<sup>T</sup> (dDDH, 60.7%; ANI, 95.0%). Because the dDDH value was <70% (13) and the ANI value was <96% (11), the results indicated that strain PS-C2 is likely a new species of *Roseovarius*. The PS-C2 genome encoded 56 CAZymes, including 13 glycoside hydrolases (GHs), 31 glycosyl transferases, 8 auxiliary activity enzymes, and 4 carbohydrate esterases. Among the GHs, 4 lytic transglycosylases (LTs) belonged to GH family 103. LTs play an essential role in bacterial cell wall metabolism and present an attractive new target for antibiotic development (14). To date, none of the *Roseovarius* LTs has been characterized. Collectively, the PS-C2 genome contributed to the body of knowledge and possible applications of *Roseovarius* spp.

**Data availability.** The whole-genome shotgun sequence of *Roseovarius* sp. PS-C2 has been deposited in NCBI GenBank under BioProject accession number [PRJNA716475](https://www.ncbi.nlm.nih.gov/bioproject/PRJNA716475), BioSample accession number [SAMN18388700](https://www.ncbi.nlm.nih.gov/biosample/SAMN18388700), and GenBank accession number [JAHKSQ000000000](https://www.ncbi.nlm.nih.gov/genbank/JAHKSQ000000000). The version described in this paper is the first version, [JAHKSQ000000000.1](https://www.ncbi.nlm.nih.gov/genbank/JAHKSQ000000000.1). The raw sequencing reads have been deposited in the NCBI Sequence Read Archive (SRA) with accession number [SRR15311899](https://www.ncbi.nlm.nih.gov/sra/SRR15311899). The 16S rRNA gene sequence of *Roseovarius* sp. PS-C2 has been deposited in NCBI GenBank with accession number [MW785753.1](https://www.ncbi.nlm.nih.gov/genbank/MW785753.1).

## ACKNOWLEDGMENTS

This work was supported by Eleventh Malaysia Plan 2016–2020 grant RMKe-11 (RP4) P30006059763005 and Malaysia Fundamental Research Grant Scheme (FRGS) grant FRGS/1/2020/WAB11/MESTECC/02/1 awarded to U.M.K. and I.I.Z. S.M.O. is thankful for International Islamic University Malaysia grant RIGS16-320-0484. K.M.G. is grateful for funding received from Malaysia FRGS grant 5F241. K.J.L. appreciates the support from Universiti Teknologi Malaysia.

## REFERENCES

- Lu L, Zhang Y, Peng X, Liu J, Qin K, Peng F. 2020. *Roseovarius arcticus* sp. nov., a bacterium isolated from arctic marine sediment. *Int J Syst Evol Microbiol* 70:2072–2078. <https://doi.org/10.1099/ijsem.0.004018>.
- Bruns H, Ziesche L, Taniwal NK, Wolter L, Brinkhoff T, Herrmann J, Müller R, Schulz S. 2018. *N*-Acylated amino acid methyl esters from marine *Roseobacter* group bacteria. *Beilstein J Org Chem* 14:2964–2973. <https://doi.org/10.3762/bjoc.14.276>.
- Chernikova TN, Bargiela R, Toshchakov SV, Shivaraman V, Lunev EA, Yakimov MM, Thomas DN, Golysheva PN. 2020. Hydrocarbon-degrading bacteria *Alcanivorax* and *Marinobacter* associated with microalgae *Pavlova lutheri* and *Nannochloropsis oculata*. *Front Microbiol* 11:572931. <https://doi.org/10.3389/fmicb.2020.572931>.
- Chaudhary DK, Khulan A, Kim J. 2019. Development of a novel cultivation technique for uncultured soil bacteria. *Sci Rep* 9:6666. <https://doi.org/10.1038/s41598-019-43182-x>.
- Chen YL, Lee CC, Lin YL, Yin KM, Ho CL, Liu T. 2015. Obtaining long 16S rDNA sequences using multiple primers and its application on dioxin-containing samples. *BMC Bioinformatics* 16(Suppl 18):S13. <https://doi.org/10.1186/1471-2105-16-S18-S13>.
- Yoon SH, Ha SM, Kwon S, Lim Y, Kim Y, Seo H, Chun J. 2017. Introducing EzBioCloud: a taxonomically united database of 16S rRNA gene sequences and whole-genome assemblies. *Int J Syst Evol Microbiol* 67:1613–1617. <https://doi.org/10.1099/ijsem.0.001755>.
- Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>.
- Luo R, Liu B, Xie Y, Li Z, Huang W, Yuan J, He G, Chen Y, Pan Q, Liu Y, Tang J, Wu G, Zhang H, Shi Y, Liu Y, Yu C, Wang B, Lu Y, Han C, Cheung DW, Yiu SM, Peng S, Xiaoqian Z, Liu G, Liao X, Li Y, Yang H, Wang J, Lam TW, Wang J. 2015. SOAPdenovo2: an empirically improved memory-efficient short-read de novo assembler. *GigaScience* 4:30. <https://doi.org/10.1186/s13742-015-0069-2>.
- Tatusova T, Dicuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res* 44:6614–6624. <https://doi.org/10.1093/nar/gkw569>.
- Auch AF, Von Jan M, Klenk H, Göker M. 2010. Digital DNA-DNA hybridization for microbial species delineation by means of genome-to-genome sequence comparison. *Stand Genomic Sci* 2:117–134. <https://doi.org/10.4056/sigs.531120>.
- Yoon SH, Ha S, Lim J, Kwon S, Chun J. 2017. A large-scale evaluation of algorithms to calculate average nucleotide identity. *Antonie Van Leeuwenhoek* 110:1281–1286. <https://doi.org/10.1007/s10482-017-0844-4>.
- Zhang H, Yohe T, Huang L, Entwistle S, Wu P, Yang Z, Busk PK, Xu Y, Yin Y. 2018. dbCAN2: a meta server for automated carbohydrate-active enzyme annotation. *Nucleic Acids Res* 46:W95–W101. <https://doi.org/10.1093/nar/gky418>.
- Meier-Kolthoff JP, Klenk HP, Göker M. 2014. Taxonomic use of DNA G+C content and DNA-DNA hybridization in the genomic age. *Int J Syst Evol Microbiol* 64:352–356. <https://doi.org/10.1099/ijms.0.056994-0>.
- Dik DA, Marous DR, Fisher JF, Mobashery S. 2017. Lytic transglycosylases: concinnity in concision of the bacterial cell wall. *Crit Rev Biochem Mol Biol* 52:503–542. <https://doi.org/10.1080/10409238.2017.1337705>.